



NBSIR 79-1777

A Compendium of Gas Phase Basicity and Proton Affinity Measurements

K. N. Hartman*, S. Lias, P. Ausloos, H. M. Rosenstock*,
S. S. Schroyer*, C. Schmidt*, D. Martinsen**, and G. W. A. Milne***

National Measurement Laboratory
National Bureau of Standards
Department of Commerce
Washington, D.C. 20234

*Ion Energetics Data Center
National Bureau of Standards
Department of Commerce
Washington, D.C. 20234

**Fein-Marquart Associates
Baltimore, MD 21212

***National Heart Lung and Blood Institute
National Institutes of Health
Bethesda, MD 20205

April 1979

Issued July 1979

This work was supported in part by the:

**National Heart, Lung, and Blood Institute
National Institutes of Health
Bethesda, MD 20205**

SEP 20 1979

NBSIR 79-1777

A COMPENDIUM OF GAS PHASE BASICITY AND PROTON AFFINITY MEASUREMENTS

K. N. Hartman*, S. Lias, P. Ausloos, H. M. Rosenstock*,
S. S. Schroyer*, C. Schmidt*, D. Martinsen**, and G. W. A. Milne***

National Measurement Laboratory
National Bureau of Standards
Department of Commerce
Washington, D.C. 20234

*Ion Energetics Data Center
National Bureau of Standards
Department of Commerce
Washington, D.C. 20234

**Fein-Marquart Associates
Baltimore, MD 21212

***National Heart Lung and Blood Institute
National Institutes of Health
Bethesda, MD 20205

April 1979

Issued July 1979

This work was supported in part by the:
National Heart, Lung, and Blood Institute
National Institutes of Health
Bethesda, MD 20205



U.S. DEPARTMENT OF COMMERCE, Juanita M. Kreps, *Secretary*

Jordan J. Baruch, *Assistant Secretary for Science and Technology*

NATIONAL BUREAU OF STANDARDS, Ernest Ambler, *Director*

THE UNIVERSITY OF CHICAGO

PHILOSOPHY DEPARTMENT

1100 S. MICHIGAN AVE.

CHICAGO, ILL. 60607

TEL: 773-936-5000

FAX: 773-936-5001

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

WWW.CHICAGOEDU.EDU

FOREWORD

The National Standard Reference Data System was established in 1963 for the purpose of promoting the critical evaluation and dissemination of numerical data of the physical sciences. The program is coordinated by the Office of Standard Reference Data of the National Bureau of Standards, but involves the efforts of many groups in universities, government laboratories, and private industry. The primary aim of the program is to provide compilations of critically evaluated numerical data. These tables are published in the Journal of Physical and Chemical Reference Data, the various publication series of the National Bureau of Standards, and through other appropriate channels.

The present report consists of tables of data on gas phase basicity and proton affinity of molecules. It represents a contribution from the Ion Energetics Data Center, prepared in collaboration with staff of NHLBI, NIH and Fein-Marquart Associates. Support for the preparation of these tables has been provided by the National Heart, Lung, and Blood Institute and the Office of Standard Reference Data.

David R. Lide, Jr.
Chief, Office of Standard
Reference Data

ABSTRACT

Numerical values of experimental measurements, calculations and estimations of gas phase basicities and proton affinities are presented. Where appropriate they are recalculated to common values for the proton affinity of ammonia and the heat of formation of the proton. The annotated information is ordered by molecular formula following the Hill System. Indices are given to permit location of information by numerical value of the measurement(s), the compound formula, name and synonym, and CAS Registry Number. The literature coverage is essentially complete through 1977.

TABLE OF CONTENTS

1. INTRODUCTION.....	iv
1.1 The Data Base.....	iv
1.2 Definitions and units.....	iv
1.3 Standard states, conventions, and reference values...	v
1.4 References.....	vii
2. THE MEASUREMENT TECHNIQUES.....	viii
2.1 Absolute measurements.....	viii
2.2 Equilibrium measurements.....	ix
2.3 Bracketing measurements.....	x
2.4 Calculations and estimates.....	xi
2.5 References.....	xi
3. ORGANIZATION OF THE INFORMATION.....	xiii
4. TABLE OF MEASUREMENTS IN MOLECULAR FORMULA ORDER.....	1
5. GAS PHASE BASICITY INDEX.....	476
6. PROTON AFFINITY INDEX.....	482
7. CAS REGISTRY NUMBER INDEX.....	489
8. MOLECULAR FORMULA INDEX.....	497
9. COMPOUND NAME INDEX.....	503


1. INTRODUCTION

1.1 The Data Base

The present compendium tabulates all experimental measurements of gas phase basicities and measurements or derived values of proton affinities available up to the end of 1977. This compilation was carried out for the NIH-EPA CIS project which is developing a comprehensive chemical data base for the use of workers concerned with research on environmental and health problems and related areas. The numerical information presented here is also accessible and searchable by computer through the NIH/EPA Chemical Information System (CIS)(1).

1.2 Definitions and units

The gas phase basicity (GB) of a molecule (M) is defined as the negative of the Gibbs free energy change for the process



and the proton affinity (PA) is defined as the negative of the enthalpy change for the same process.

$$PA(M) = \Delta H_f(M) + \Delta H_f(H^+) - \Delta H_f(MH^+)$$


For completeness, the temperature has to be specified. It is generally desired to evaluate the quantity at room temperature. Occasionally experiments are carried out at high temperatures (e.g. 600 K or at undefined temperatures (threshold measurements)), and for these it is necessary to apply the appropriate thermochemical corrections or threshold interpretations.

From the relation $\Delta G = \Delta H - T\Delta S$ the proton affinity can be determined from the gas phase basicity if the temperature and the entropy change of the process are known or can be estimated.

The unit in most widespread use is the kcal/mol and this unit is adopted throughout. It is related to the recommended SI unit by $1 \text{ kcal} = 4.1840 \text{ kJ}$, exactly.

1.3 Standard states, conventions, and reference values

The process



makes no reference to an electron and no difficulties arise if the heats and free energies of formation of all species are defined with the same ionic standard state. Unfortunately this has not always been the case.

Historically, the community determining ion heats of formation by threshold measurements have assumed the convention that the electron is at rest (as it surely is at threshold) and that it remains at rest even when deducing finite (or room) temperature heats of formation of the ions. This convention is also adopted in the two major compilations of gaseous ion thermochemistry (2,3).

The two principal compilations of neutral thermochemistry (4,5) assign to the electron a classical heat capacity, C_p , of $5/2 R$. Thus at room temperature the heats of formation of all ions will be larger than those in, for example Energetics of Gaseous Ions (GIE) (3) by an amount equal to

$$\int_0^{298.15} C_p dT = \frac{5}{2} RT = 1.481 \text{ kcal/mol}$$

Calculations of proton affinities using room temperature ion heats of formation from threshold measurements as compiled and evaluated in Energetics of Gaseous Ions (3) and the proton heat of formation as given in the NBS Tech Note 270 series (4) or the JANAF tables (5) will be in error (too large) by this amount. Wherever the original reference quoted proton affinity values were in error for this reason, we have made the appropriate correction. When the heat of formation of the ion M^+ , is defined according to the GIE convention, the value for the heat of formation of the proton at 298.15 K is 365.7 kcal/mol. This subject is discussed in detail in a recent review article (6).

A major part of the measurement base consists of relative measurements which can be put on an absolute basis by relating to a small number of absolute values. Unfortunately, the magnitude of some of the absolute values is still in doubt. For instance, many of the extensive scales of proton affinity ladders which have been published (7,8) have ultimately been related to the heat of formation of $t\text{-C}_4\text{H}_9^+$ as a primary standard, although the values are usually cited relative to the proton affinity of a secondary standard, ammonia. At the time the present compilation was carried out, the latter value was considered to be established at 202.3 kcal/mol (6), and it is this number which is used as the standard for the values cited here. More recent redeterminations of the heat of formation (9,10) and ionization potential (11) of the t -butyl radical, as well as additional information on the entropy change associated with the deprotonation of $t\text{-C}_4\text{H}_9^+$ (12), have resulted in a revised estimate that the proton affinity of ammonia is

greater than or equal to 207.0 kcal/mol (6). Thus, the absolute values of proton affinities given here based on relative ΔG measurements and expressed "relative to ammonia" are, according to the most recent information, too low by at least 5 kcal/mol. In this and other cases we have indicated the species to which the relative measurement is referred, so that readjustment can be made as additional information becomes available.

1.4 References

1. S. R. Heller, G. W. A. Milne, and R. J. Feldmann, "A Computer Based Chemical Information System," Science 195, 253 (1977).
2. J. L. Franklin, J. G. Dillard, H. M. Rosenstock, J. T. Herron, K. Draxl, and F. H. Field, "Ionization Potentials, Appearance Potentials, and Heats of Formation of Gaseous Positive Ions," NSRDS-NBS-26, June 1969.
3. H. M. Rosenstock, K. Draxl, B. W. Steiner, and J. T. Herron, "Energetics of Gaseous Ions," J. Phys. Chem. Ref. Data 6 Supplement 1 (1977).
4. D. D. Wagman, et al, "Selected Values of Chemical Thermodynamics Properties," NBS Technical Note 270-3, January 1968, and subsequent Tech Notes in this series.
5. D. R. Stull and H. Prophet, "JANAF Thermochemical Tables," Second Edition, NSRDS-NBS-37, June 1971 and supplements published in J. Phys. Chem. Ref. Data.
6. S. Lias, "Thermochemistry of Polyatomic Cations," in "Kinetics of Ion-Molecule Reactions," (P. Ausloos, ed.) Plenum Press, New York, pg. 233.
7. J. F. Wolf, R. H. Staley, I. Koppel, M. Taagepera, R.


- T. McIver, Jr., J. L. Beauchamp, and R. W. Taft, "Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements," J. Am. Chem. Soc. 99, 5417 (1977).
8. R. Yamdagni, and P. Kebarle, "Gas Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements," J. Am. Chem. Soc. 98, 1320 (1976).
9. W. Tsang, "Evidence for Strongly Temperature-Dependent A-factors in Alkane Decomposition and High Heats of Formation for Alkyl Radicals," Int. J. Chem. Kinetics 11, 820 (1978).
10. M. Rossi and D. M. Golden, J. Am. Chem. Soc. submitted for publication.
11. F. A. Houle and J. L. Beauchamp, "Photoelectron Spectroscopy of Methyl, Ethyl, Isopropyl, and Tert-Butyl Radicals. Implications for the Thermochemistry and Structures of the Radicals and their Corresponding Carbonium Ions," J. Am. Chem. Soc., in press.
12. P. Ausloos and S. G. Lias "Entropy Changes for the Protonation of Alkenes," J. Am. Chem. Soc. 100, 1953 (1978).

2. THE MEASUREMENT TECHNIQUES

2.1 Absolute measurements

These are generally based on measurement of the heat of formation of a proton-containing species by a threshold


process, such as



From the threshold energy of this process and the heats of formation of the neutral atomic and molecular species one can deduce heat of formation of HO_2^+ , and by further straightforward calculation the proton affinity of molecular oxygen. In actual practice considerable care has to be exercised to establish the temperature to which the heat of formation refers. Threshold measurements leading to more or less accurate absolute values have been carried out by photoionization (PI), monoenergetic electron impact (EM) and conventional electron impact (EI) techniques. The accuracy of these techniques has been comprehensively discussed in a recent review article (1).

2.2 Equilibrium measurements

From a determination of the proton transfer equilibrium constant for



one can determine the Gibbs free energy change of the reaction. This gives the basicity difference between the two compounds. In order to determine the proton affinity difference it is necessary to determine the entropy change of the reaction. The most direct and unambiguous method is to measure the equilibrium constant as a function of temperature, which yields the proton affinity directly from the Van't Hoff relation

$$\frac{d \ln K_{\text{eq}}}{d(1/T)} = \frac{-\Delta H}{R}$$

So far for proton transfer equilibria this has only been performed in a few instances, with varying degrees of success (2). Most estimates of the relative proton affinity are based on assumptions about the entropy change. These range from consideration of symmetry changes in the reaction, to reasoning by analogy, to the assumption that the entropy change is zero. The method or the assumptions made for obtaining a value for ΔS are indicated in this compendium for each determination.

Equilibrium constant measurements are carried out in three quite distinct experimental systems operating at rather different pressures. These are the high pressure ion source coupled to a mass spectrometer (MS), the flowing afterglow tube (FAG) whose equilibrium ion composition is sampled by a quadrupole mass spectrometer, and ions trapped for finite time in a cyclotron field (ICR) with subsequent ejection and analysis by rf absorption. These techniques have been discussed in detail in a number of reviews (2-6).

2.3 Bracketing measurements

The same techniques described immediately above can be used to examine the occurrence or non-occurrence of a proton transfer reaction, and applying the criterion that occurrence implies a negative free energy change and non-occurrence implies a positive free energy change. Accepting these assumptions, one could in principle bracket the range which the gas phase basicity of a molecule lies (7). Using different reagent ions many authors have reported such relative basicities as relative "proton affinities". Results

obtained from this approach should be used only with great caution; it is now well known that many endoergic reactions may occur with a rate constant high enough that they can easily be observed under the experimental conditions described above (3). Furthermore, many proton transfer reactions which are clearly exothermic are not observed to occur if more favorable reaction channels exist (8).

2.4 Calculations and estimates

In principle, proton affinities or ion heats of formation can be calculated quantum mechanically. We include here only the very high quality calculations carried out on H_3^+ (see Tables). Another type of calculation deals with the evaluation of the lattice energy of an ionic crystal (CLE) (9). For an ammonium halide, for example, this gives the energy of the crystal relative to the gas phase ammonium and halide ions. These can then be related to gas phase neutral species via a Born-Haber cycle, and will yield the proton affinity of $\text{NH}_3(\text{gas})$. Lastly, there are a number of empirical correlations of proton affinities with other physical parameters such as core ionization potentials (10).


2.5 References

1. H. M. Rosenstock, "The Measurement of Ionization and Appearance Potentials," Int. J. Mass Spectrom. Ion Phys. 20, 139 (1976).
2. S. G. Lias, "Thermodynamic Information from Ion-Molecule Equilibrium Constant Determinations," in "Ion Cyclotron Resonance Spectrometry" (H. Hartmann and K.-P. Wanczek, eds.), Springer-Verlag, Berlin (1978).

3. S. Lias, "Thermochemistry of Polyatomic Cations," in "Kinetics of Ion Molecule Reactions," (P. Ausloos, Ed.), Plenum Press, New York (1979), pg. 233.
4. P. Kebarle, "Ion Thermochemistry and Solvation from Gas Phase Ion Equilibria," Ann. Rev. Phys. Chem. 28, 445 (1977).
5. D. H. Aue and M. T. Bowers, "Stabilities of Positive Ions from Equilibrium Gas Phase Basicity Measurements," in "Gas Phase Ion Chemistry," (M. T. Bowers, ed.), Academic Press, New York (1979).
6. D. K. Bohme, "The Kinetics and Energetics of Proton Transfer," in "Interactions between Ions and Molecules," (P. Ausloos, ed.), Plenum Press, New York (1975).
7. J. Long and B. Munson, "Proton Affinities of some Oxygenated Compounds," J. Am. Chem. Soc. 95, 2427 (1973).
8. P. Ausloos and S. G. Lias, "Proton Affinity of Dichlorocarbene," J. Am. Chem. Soc. 100, 4594 (1978).
9. J. Sherman, "Crystal Energies of Ionic Compounds and Thermochemical Applications," Chem. Rev. 11, 93 (1932).
10. J. A. Jen and T. D. Thomas, "Core Ionization Potentials in Dimethyl Ether and Methylamine," J. Electron Spectrosc. 4, 43 (1974).

3. ORGANIZATION OF THE INFORMATION

3.1 A typical measurement entry is shown below:

C ₈ H ₁₀	CAS REGN : 108-38-3
	
Benzene, 1,3-dimethyl- (9CI) m-Xylene (8CI)	
GB (exp) : 182+/-3 kcal/mol	Temperature : 340 K
PA (exp) : 189.7+/-3 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Equilibrium relative to CH ₃ SH and (CH ₃) ₂ O Entropy change assumed zero	
GB (exp) : 187.2+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 194.6+/-2 kcal/mol	Temperature : 298 K
Devlin, J. L., III, Wolf, J. F., Taft, R. W., and Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to C ₆ H ₆ (Benzene)	

The entry gives successively the following information:

- a) The empirical formula in Hill System order.
- b) The Chemical Abstracts Service (CAS) Registry Number.
- c) The structural formula (see discussion below).
- d) The Chemical Abstracts Service (CAS) Name, as given in the 8th Collective Index (8CI) and the 9th Collective Index (9CI). If it is the same in both it is denoted with (8CI9CI).
- e) The gas phase basicity, GB, in kcal/mol.
- f) The proton affinity, PA, in kcal/mol.
- g) The temperature at which the measurement was carried out.
- h) The literature citation.
- i) The method used (see discussion below).
- j) Additional comments on the nature of the measurement.


Where there are several measurements on the same species, the numerical value, literature reference, method and associated comments are listed successively.

The structural diagrams used in this publication have been generated from connection tables by the Structure and Nomenclature Search System (SANSS), one of the components of the CIS. The following symbols have been used to represent the different types of chemical bonds:

- * - single bond
- + - double bond
- # - triple bond
- . - ring alternating bond (e.g., benzene)
- % - tautomeric bond (e.g., O% C% O in carboxylic acids)

Hydrogen atoms are not explicitly included in the diagrams but are understood to occupy any otherwise available bonds.

Tautomeric structures are recognized as equivalent and the percent sign used in the structural diagrams under the following conditions.



- when:
- a). Q = C, N, S, P, Sb, As, Se, Te, Br, Cl, or I with any allowed valence for the individual elements.
 - b). M and Z = any combination of trivalent N and/or bivalent O, S, Se or Te atoms.
 - c). The bonds which are involved in the tautomerization may be in an acyclic chain, a ring system, or partly in each.
 - d). The end points, M and Z, may be in adjacent rings of a fused ring system. However, a nitrogen atom at a fusion point in a fused ring system can not be involved in tautomerization.
 - e). A deuterium or tritium may replace the migrating hydrogen atom.
 - f). If more than one of the systems described above are linked through a common atom, the hydrogen may be free to migrate along the chain defining a larger tautomeric system.

The experimental methods were discussed in the preceding section. They are abbreviated as follows:

CLE - Crystal lattice energy
 EI - Threshold measurement by non-monoenergetic electron impact
 EM - Threshold measurement by monoenergetic electron impact
 FAG - Flowing afterglow
 FI - Field ionization threshold
 ICR - Ion cyclotron resonance
 MS - High pressure mass spectrometry
 PI - Threshold measurement by photoionization
 TC - Theoretical calculation

Following the table of measurements in molecular formula order, Section 4, there are a number of indexes:

- a). The gas phase basicity index in which the file is ordered in increasing order of gas phase basicity, with numerical values arbitrarily given to four significant figures for purposes of ordering. The page number refers to the main body of information, Section 4.
- b). The proton affinity index, structured in the same fashion.
- c,d,e). Additional indices are provided for molecular formula, CAS Registry Number and compound name. The compound name index includes a number of synonyms in addition to the names adopted in the 8th and 9th collective indices.



Ar	CAS REGN : 7440-37-1
AR	
Argon (8CI9CI)	
PA (exp) : <100.+/-5 kcal/mol	Temperature :
Roche, A. E., Sutton, M. M., Bohme, D. K., and Schiff, H. I., Determination of Proton Affinity from the Kinetics of Proton Transfer Reactions. I. Relative Proton Affinities, J. Chem. Phys. 55, 5480 (1971).	
Method used : FAG Bracketing relative to 02.	

AsH3	CAS REGN : 7784-42-1
AS	
Arsine (8CI9CI)	
GB (exp) : 172.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

BF2H	CAS REGN : 13709-83-6
F**B**F	
Borane, difluoro- (8CI9CI)	
PA (exp) : 115+/-14 kcal/mol	Temperature : 85 K
Pierce, R. C., and Porter, R. F., Ion-Molecule Chemistry of BF3 and HBF2 in Hydrogen, Inorg. Chem. 14, 1087 (1975).	
Method used : MS Bracketing relative to CH4 and H2.	

BF3	CAS REGN : 7637-07-2
F**B**F * * F	
Borane, trifluoro- (9CI) Boron fluoride (BF3) (8CI)	
PA (exp) : 115+/-14 kcal/mol	Temperature : 85 K
Pierce, R. C., and Porter, R. F., Ion-Molecule Chemistry of BF3 and HBF2 in Hydrogen, Inorg. Chem. 14, 1087 (1975).	
Method used : MS Bracketing relative to CH4 and H2.	

B2H6	CAS REGN : 19287-45-7
H**B * * * * B**H	
Diborane(6) (8CI9CI)	
PA (exp) : 147+/-4 kcal/mol	Temperature : 79-86 K
Pierce, R. C., and Porter, R. F., Low-Temperature Chemical Ionization Mass Spectrometry of Boron Hydrides. The Proton Affinities of Diborane and Tetraborane(10), J. Am. Chem. Soc. 95, 3849 (1973).	
Method used : MS Observed reaction $C_2H_9^+ + B_2H_6 \rightarrow 2CH_4 + B_2H_7^+$. Reaction $B_2H_7^+ \rightarrow B_2H_5^+ + H_2$ has delta H formation = 3.0+/-0.3. Based on delta H formation($B_2H_5^+$) = 232.	

B3H5N3	CAS REGN : Not available
<pre> B**N * * * * * * N B * * * * * * * * * * * * B**N </pre>	
.beta.-Boraziny1 radical	
PA (exp) : 193+/-2 kcal/mol	Temperature :
DeStefano, A. J., and Porter, R. F. Ion-Molecule Reactions of Cyclic Borazine Cations. Thermodynamic and Kinetic Considerations, J. Phys. Chem. 80, 2818 (1976).	
Method used : MS Bracketing relative to HC02C2H5 and (CH3)2CO. The reactant ions are generated by photon impact	

B3H6N3	CAS REGN : 6569-51-3
<pre> B**N * * * * * * N B * * * * * * B**N </pre>	
Borazine (8CI9CI)	
PA (exp) : 199+/-4 kcal/mol	Temperature :
Betowski, L. D., Solmon, J. J., and Porter, R. F., The Proton Affinity of Borazine, Inorg. Chem. 11, 424 (1972).	
Method used : MS Bracketing relative to C4H6(2-Butyne) and NH3	

B4H8	CAS REGN : 12007-71-5
B***B * * * * B***B	
Tetraborane(8)	
PA (exp) : 187+/-7 kcal/mol	Temperature : 375 K
Solmon, J. J., and Porter, R. F., Chemical Ionization Mass Spectrometry of Selected Boron Hydrides, J. Am. Chem. Soc. 94, 1443 (1972).	
Method used : MS Bracketing relative to C3D6 and (CH3)2CO.	

B4H10	CAS REGN : 18283-93-7
<pre> H*****B*****H * * * * * * * * B*****B * * * * * * * * H*****B*****H </pre>	
Tetraborane(10) (8CI9CI)	
PA (exp) : 144+/-5 kcal/mol	Temperature : 140-300 K
Pierce, R. C., and Porter, R. F., Low-Temperature Chemical Ionization Mass Spectrometry of Boron Hydrides. The Proton Affinities of Diborane and Tetraborane(10), J. Am. Chem. Soc. 95, 3849 (1973).	
Method used : MS Based on delta H formation(B2H5+) and estimating delta S. Equilibrium of B2H5+ + B2H6 -> B4H11+.	

B5H9	CAS REGN : 19624-22-7
<pre> H*****B*****H * *** * * * * * * B**B**B * * * * * * *** * H*****B*****H </pre>	
Pentaborane(9) (8CI9CI)	
PA (exp) : 172+/-2 kcal/mol	Temperature : 373~3 K
Solmon, J. J., and Porter, R. F., Chemical Ionization Mass Spectrometry of Selected Boron Hydrides, J. Am. Chem. Soc. 94, 1443 (1972).	
Method used : MS Based PA (H2O) = 170.3. Bracketing relative to H2O and H2S.	

BrH	CAS REGN : 10035-10-6
BR	
Hydrobromic acid (8CI9CI)	
PA (exp) : 141+/-3 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Absolute PA value estimated by assuming that for reaction $RH^+ + RH \rightarrow RH_2^+ + R$; the $\Delta H \sim 5X$ TC. TC = measured translational energy of the products.	

CBrN	CAS REGN : 506-68-3
N##C**BR	
Cyanogen bromide (8CI9CI)	
PA (exp) : 179+/-3 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to AsH3, HCO2H and F2CHCH2OH	

CC1N	CAS REGN : 506-77-4
N##C**CL	
Cyanogen chloride (8CI9CI)	
PA (exp) : 177+/-3 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Equilibrium relative to F2CH2CH, OH and HCHO	
GB (exp) : 169.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 177.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

CC12	CAS REGN : 1605-72-7
CL*C**CL	
Methylene, dichloro- (8CI9CI)	
PA (exp) : 194.7+/-4 kcal/mol	Temperature : 325 K
Ausloos, P., and Lias, S. G. The Proton Affinity of Dichlorocarbene, To be published in J. Am. Chem. Soc.	
Method used : ICR Bracketing relative to Me20 and Et20; scale based on PA (NH3) = 202.8	
PA (exp) : 200.4+/-1.5 kcal/mol	Temperature : 300 K
Lias, S. G., and Ausloos, P. Reactions of CC12H+ and CF2H+ with Organic and Inorganic Compounds: Proton Affinities and Heats of Formation of CC12 and CF2, Intern. J. Mass Spectrom. Ion Phys. 22, 135 (1976).	
Method used : ICR Bracketing relative to NH3 and (CH3)2S	

Continued on next page

GB not calculated for this reference	
PA (exp) : 208.3+/-2 kcal/mol	Temperature : 300 K
Levi, B. A., Taft, R. W., and Hehre, W. J. Dichlorocarbene, J. Am. Chem. Soc. 99, 8454 (1977).	
Method used : ICR Calculated entropy Bracketing relative to CF ₃ CH ₂ CH ₂ NH ₂ and C ₆ H ₅ NH ₂ ; scale based on NH ₃	
PA (exp) : 209+/-2 kcal/mol	Temperature : 300 K
Levi, B. A., Taft, R. W., and Hehre, W. J. Dichlorocarbene, J. Am. Chem. Soc. 99, 8454 (1977).	
Method used : ICR Calculated entropy The deuterium affinity determined by bracketing techniques relative to 2-CF ₃ Pyridine and CF ₃ (CH ₂) ₂ NH ₂ ; scale based on NH ₃	

CF2	CAS REGN : 2154-59-8
F**C**F	
Methylene, difluoro- (8CI9CI)	
PA (exp) : 178.5+/-1 kcal/mol	Temperature : 300 K
Lias, S. G., and Ausloos, P. Reactions of CCl₂H⁺ and CF₂H⁺ with Organic and Inorganic Compounds: Proton Affinities and Heats of Formation of CCl₂ and CF₂, Intern. J. Mass Spectrom. Ion Phys. 22, 135 (1976).	
Method used : ICR Bracketing relative to H₂S and HCOOH and also observing the following: CD₅⁺ + CF₂HCL -> CD₄ + DCl + CF₂H⁺ and CD₅⁺ + CF₃H -> CD₄ + DF + CF₂H⁺. Scale based on PA (H₂O) = 170.1 kcal/mol	

CF4	CAS REGN : 75-73-0
<pre> F * * F**C**F * * F </pre>	
Methane, tetrafluoro- (9CI) Carbon tetrafluoride (8CI)	
PA (exp) : 121+/-4 kcal/mol	Temperature : 300 K
Roche, A. E., Sutton, M. M., Bohme, D. K., and Schiff, H. I., Determination of Proton Affinity from the Kinetics of Proton Transfer Reactions. I. Relative Proton Affinities, J. Chem. Phys. 55, 5480 (1971).	
Method used : FAG Bracketing relative to CH4 and N2.	
PA (exp) : 121+/-5 kcal/mol	Temperature : 298 K
Blint, R. J., McMahon, T. B., and Beauchamp, J. L., Gas-Phase Ion Chemistry of Fluoromethanes by Ion Cyclotron Resonance Spectroscopy. New Techniques for the Determination of Carbonium Ion Stabilities, J. Am. Chem. Soc. 96, 1269 (1974).	
Method used : ICR Bracketing relative to CH4. Using IP of H and CF4 and also HA of CF4+.	

CHF3	CAS REGN : 75-46-7
F**C**F * * F	
Methane, trifluoro- (8CI9CI)	
PA (exp) : 147+/-5 kcal/mol	Temperature : 298 K
Blint, R. J., McMahon, T. B., and Beauchamp, J. L., Gas-Phase Ion Chemistry of Fluoromethanes by Ion Cyclotron Resonance Spectroscopy. New Techniques for the Determination of Carbonium Ion Stabilities, J. Am. Chem. Soc. 96, 1269 (1974).	
Method used : ICR Bracketing relative to CH3F. Using IP of H and CH3F, also HA (CH3F+).	

CHN	CAS REGN : 74-90-8
C##N	
Hydrocyanic acid (8CI9CI)	
GB (exp) : 166.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 174.5+/-2 kcal/mol	Temperature : 300 K
Freeman, C. G., Harland, P. W., and McEwan, M. J. The Equilibrium $\text{H}_3\text{S}^+ + \text{HCN} = \text{H}_2\text{CN}^+ + \text{H}_2\text{S}$ and the Relative Proton Affinities of HCN and H_2S , Intern. J. Mass Spectrom. Ion Phys. 27, 77 (1978).	
Method used : FAG Equilibrium relative to H_2S ; scale based on NH_3	
PA (exp) : 139 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Estimated $\Delta H = 5X$ translational energy of the product of reaction $\text{RH}^+ + \text{RH} \rightarrow \text{RH}_2^+ + \text{R}$	

Continued on next page

GB (exp) : 166.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 174.5+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

CH2F2	CAS REGN : 75-10-5
F**C**F	
Methane, difluoro- (8CI9CI)	
PA (exp) : 147+/-5 kcal/mol	Temperature : 298 K
Blint, R. J., McMahon, T. B., and Beauchamp, J. L., Gas-Phase Ion Chemistry of Fluoromethanes by Ion Cyclotron Resonance Spectroscopy. New Techniques for the Determination of Carbonium Ion Stabilities, J. Am. Chem. Soc. 96, 1269 (1974).	
Method used : ICR Bracketing relative to CH3F and CO. Using IP of H and CH2F2, also HA (CH2F2+).	

CH2N2	CAS REGN : 334-88-3
N++N++C	
Methane, diazo- (8CI9CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature : 300 K
Foster, M. S., and Beauchamp, J. L., Gas-Phase Ion Chemistry of Azomethane by Ion Cyclotron Resonance, J. Am. Chem. Soc. 94, 2425 (1972).	
Method used : ICR Bracketing relative to NH3 and CH3NH2	

CH2O	CAS REGN : 50-00-0
C++O	
Formaldehyde (8CI9CI)	
PA (exp) : 162+/-3 kcal/mol	Temperature :
Harrison, A. G., Ivko, A., and Van Raalte, D., Energetics of Formation of Some Oxygenated Ions and the Proton Affinities of Carbonyl Compounds, Can. J. Chem. 44, 1625 (1966).	
Method used : EI Electron impact fragmentation (various molecules).	

Continued on next page

PA (exp) : 169+/-1 kcal/mol	Temperature : 298 K
Refaey, K. M. A., and Chupka, W. A., Photoionization of the Lower Aliphatic Alcohols with Mass Analysis, J. Chem. Phys. 48, 5205 (1968).	
Method used : PI Appearance potential by IP fragmentation of CH ₃ OH	
PA (exp) : 170 kcal/mol	Temperature :
Lossing, F. P. Heats of Formation of Some Isomeric [C _n H _{2n} + 10] ⁺ Ions. Substitutional Effects on Ion Stability, J. Am. Chem. Soc. 99, 7526 (1977).	
Method used : EM Electron impact fragmentation (various molecules) using EM. $\Delta H_f(H^+) = 365.7$	
GB (exp) : 167.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 174.6+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

CH2O2	CAS REGN : 64-18-6
O=C=O	
Formic acid (8CI9CI)	
PA (exp) : 178+/-4 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH3OH and H2S.	
PA (exp) : 186.8 kcal/mol	Temperature :
Goldenfeld, I. V., Korostyshevsky, I. Z., and Mischanchuk, B. G., Analysis of Field Ion Energies in a Mass Spectrometer, Int. J. Mass Spectrom. Ion Phys. 13, 297 (1974).	
Method used : FI PA derived from field appearance potential. Correction for neutral particles potential energy	

Continued on next page

PA (exp) : 179+/-3 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Heats of Formation of Protonated Cyclopropane, Methylcyclopropane, and Ethane, J. Am. Chem. Soc. 94, 5347 (1972).	
Method used : MS Correction for path degeneracy. Equilibrium; scale based on CH ₃ CH=CH ₂ .	
GB (exp) : 162.5+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 178.9+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH ₃	

Continued on next page

GB (exp) : 171.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

CH3Br	CAS REGN : 74-83-9
BR*C	
Methane, bromo- (8CI9CI)	
PA (exp) : 163 kcal/mol	Temperature : 300 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also IP's of H and CH3Br and HA(CH3BR+).	

CH3Cl	CAS REGN : 74-87-3
C**CL	
Methane, chloro- (8CI9CI)	
PA (exp) : <170.3 kcal/mol	Temperature :
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds,. J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to H2O.	
PA (exp) : 164+/-3 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Absolute PA value estimated by assuming that for reaction $RH^+ + RH \rightarrow R + RH_2^+$ the $\Delta H \sim 5X \text{ TC}$. TC = measured translational energy of the products	

Continued on next page

PA (exp) : ~160 kcal/mol	Temperature : 300 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also IP's of H and CH₃Cl and HA(CH₃Cl⁺).	

CH ₃ F	CAS REGN : 593-53-3
C**F	
Methane, fluoro- (8CI9CI)	
PA (exp) : 151+/-5 kcal/mol	Temperature : 298 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Using also the IP's of H and CH₃F and HA of CH₃F⁺.	

CH3I	CAS REGN : 74-88-4
C**I	
Methane, iodo- (8CI9CI)	
PA (exp) : 174+/-3 kcal/mol	Temperature : 300 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also IP's of H and CH3I and HA(CH3I+).	

CH3NO2	CAS REGN : 75-52-5
O++N**C + + O	
Methane, nitro- (8CI9CI)	
PA (exp) : ~182 kcal/mol	Temperature : 300 K
McAllister, T., and Pitman, P. Ion-Molecule Reactions and Proton Affinities of Methyl Nitrite and Nitromethane, Intern. J. Mass Spectrom. Ion Phys. 19, 241 (1976).	
Method used : ICR Bracketing relative to CH3OH and (CH3)2CO	
PA (exp) : 182+/-2 kcal/mol	Temperature : 300 K
Kriemler, P., and Buttrill, S. E., Jr., Ion-Molecule Reactions and the Proton Affinities of the Nitroalkanes. I. Nitro methane and nitroethane, J. Am. Chem. Soc. 95, 1365 (1973).	
Method used : ICR Bracketing relative to CH3OH.	

CH3NO2	CAS REGN : 624-91-9
O++N**O**C	
Nitrous acid, methyl ester (8CI9CI) .	
PA (exp) : 190.3+/-3 kcal/mol	Temperature : 300 K
McAllister, T., and Pitman, P. Ion-Molecule Reactions and Proton Affinities of Methyl Nitrite and Nitromethane, Intern. J. Mass Spectrom. Ion Phys. 19, 241 (1976).	
Method used : ICR Bracketing relative to C2H5OH and (CH3)2CO	

CH4	CAS REGN : 74-82-8
C	
Methane (8CI9CI)	
PA (exp) : 127+/-2 kcal/mol	Temperature : 78,300 K
Chupka, W. A., and Berkowitz, J., Photoionization of Methane: Ionization Potential and Proton Affinity of CH4, J. Chem. Phys. 54, 4256 (1971).	
Method used : PI From PI-efficiency curve of CH5+ in CH4.	

Continued on next page

PA (exp) : 126+/-1 kcal/mol	Temperature : 483+/-10 K
Munson, M. S. B., and Field, F. H., Reactions of Gaseous Ions. XV. Methane + 1% Ethane and Methane + 1% Propane, J. Am. Chem. Soc. 87, 3294 (1965).	
Method used : MS Observed reactions: $\text{CH}_4^+ + \text{CH}_4 \rightarrow \text{CH}_5^+ + \text{CH}_3$, $\text{CH}_5^+ + \text{C}_2\text{H}_6 \rightarrow \text{C}_2\text{H}_5^+ + \text{H}_2 + \text{CH}_4$.	
PA (exp) : 131.4 kcal/mol	Temperature : 298 K
Schiff, H. I., and Bohme, D. K., Flowing Afterglow Studies at York University, Int. J. Mass Spectrom. Ion Phys. 16, 167 (1975).	
Method used : FAG Equilibrium relative to N_2O , ΔS estimated. Based on PA (N_2O) which is derived from calculated PA (CO).	
PA (exp) : 130.5 kcal/mol	Temperature : 340 K
Kasper, S. F., and Franklin, J. L., Ion-Molecule Reactions in the System $\text{CO}_2\text{-CH}_4$, J. Chem. Phys. 56, 1156 (1972).	
Method used : MS Entropy change assumed zero Equilibrium relative to CO_2 ; found ΔG $G = -1.35 \text{ Kcal/mol}$. Based on CO_2 with $\text{PA} = 129.1$. (Int. J. Mass Spectrom. Ion Phys. 16, 167 (1975)).	

CH40	CAS REGN : 67-56-1
C**0	
Methanol (8CI9CI)	
PA (exp) : 182+/-3 kcal/mol	Temperature :
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH3CHO and HCOOH.	
PA (exp) : 179.0+/-2 kcal/mol	Temperature : 299-329 K
Wei, L. Y., and Bone, L. I., Ion-Molecule Reactions in Methanol and Hydrogen Sulfide, J. Phys. Chem. 78, 2527 (1974).	
Method used : MS Equilibrium relative to H2S Van't Hoff plot	

Continued on next page

PA (exp) : 182 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Delta H estimated = 5X total translational energy of the products of $RH^+ + RH \rightarrow RH_2^+ + R$	
GB (exp) : 166.5+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 182.1+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 174.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 182.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

CH4S	CAS REGN : 74-93-1
C**S	
Methanethiol (8CI9CI)	
GB (exp) : 177.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 185.3+/-2 kcal/mol	Temperature : 300 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on CH3CHO Entropy change assumed zero	

Continued on next page

PA (exp) : 199+/-10 kcal/mol	Temperature :
Hobrock, B. G. and Kiser, R. W., Electron Impact Investigations of Sulfur Compounds. II. 3-Methyl-2-Thiabutane, 4-Thia-1-Pentene, and 3,4-Dithiahexane, J. Phys. Chem. 67, 648 (1963).	
Method used : EI Appearance potential by electron impact of sulfur compounds.	
GB (exp) : 178.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 185.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

CH5N	CAS REGN : 74-89-5
C**N	
Methanamine (9CI) Methylamine (8CI)	
GB (exp) : 202.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 211.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 211 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Estimated delta H = 5X translational energy of the product of reaction $RH^+ + RH \rightarrow RH_2^+ + R$	

Continued on next page

GB (exp) : 205.7+/-3 kcal/mol	Temperature : 298 K
PA (exp) : 213.7+/-3 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T. Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 195.9+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 213.1+/-2 kcal/mol	Temperature : 600 K
Briggs, J. P., Yamdagni, R., and Kebarle, P., Intrinsic Basicities of Ammonia, Methylamines, Anilines, and Pyridine from Gas-Phase Proton- Exchange Equilibria, J. Am. Chem. Soc. 94, 5128 (1972).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

CH5P	CAS REGN : 593-54-4
C**P	
Phosphine, methyl- (8CI9CI)	
GB (exp) : 193.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 201.8+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

CO	CAS REGN : 630-08-0
C##0	
Carbon monoxide (8CI9CI)	
PA (exp) : 143+/-2 kcal/mol	Temperature :
Matthews, C. S., and Warneck, P., Heats of Formation of CH0+ and C3H3+ by Photoionization, J. Chem. Phys. 51, 854 (1969).	
Method used : PI Appearance potential by photoionization. PA value derived from delta H formation(HCO+).	
PA (exp) : 142.7+/-3 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Excess Energies in Mass Spectra of some Oxygen- Containing Organic Compounds, Trans. Faraday Soc. 65, 1794 (1969).	
Method used : MS Appearance potential by electron impact from HCOOH, CH3CHO and C2H3CHO. Correction for excess energy.	

C02	CAS REGN : 124-38-9
0++C++0	
Carbon dioxide (8CI9CI)	
PA (exp) : 130.7+/-1 kcal/mol	Temperature :
Warneck, P. Heat of Formation of the HCO Radical, Z. Naturforsch. 29, 350 (1974).	
Method used : PI Appearance potential by photoionization. $\Delta H_f(H^+)$ = 365.7	
PA (exp) : 129.1 kcal/mol	Temperature : 298 K
Schiff, H. I., and Bohme, D. K., Flowing Afterglow Studies at York University, Int. J. Mass Spectrom. Ion Phys. 16, 167 (1975).	
Method used : FAG Calculated entropy. Equilibrium; scale based on CO VALUE	

Continued on next page

PA (exp) : 124+/-2 kcal/mol	Temperature : 300 K
Meot-Ner (Mautner), M., and Field, F. Proton Affinities and Cluster Ion Stabilities in CO ₂ and CS ₂ . Applications in Martian Ionspheric Chemistry, J. Chem. Phys. 66, 4527 (1977).	
Method used : MS Van't Hoff plot Equilibrium; scale based on PA (CH ₄) = 126.8	
GB (exp) : 123+/-3 kcal/mol	Temperature : 300 K
PA (exp) : 124+/-3 kcal/mol	Temperature : 300 K
Meot-Ner (Mautner), M., and Field, F. H. Proton Affinities and Cluster Ion Stabilities in CO ₂ and CS ₂ . Application in Martian Ionspheric Chemistry, J. Chem. Phys. 66, 4527 (1977).	
Method used : MS Calculated entropy The measurements done in the temperature range of 400-650K. Using the Hoff plot the delta H and delta S at 300K were derived. Using PA (CH ₄)-126.8	

Continued on next page

PA (exp) : 123+/-3 kcal/mol	Temperature : 800 K
Fehsenfeld, F. C., Lindinger, W., Schiff, H. I., Hemsworth, R. S., and Bohne, D. K. Determination of the Proton Affinity from the Kinetics of Proton Transfer Reactions. VI. The Relative Proton Affinities of N₂, Xe, and CO₂, J. Chem. Phys. 64, 4887 (1976).	
Method used : FAG Van't Hoff plot Equilibrium relative to N₂ and Xe; scale based on PA (N₂) = 112.6 Van't Hoff plot	

CS ₂	CAS REGN : 75-15-0
S++C++S	
Carbon disulfide (8CI9CI)	
GB (exp) : 165.7+/-2.5 kcal/mol	Temperature : 300 K
PA (exp) : 168+/-2.5 kcal/mol	Temperature : 300 K
Meot-Ner (Mautner), M., and Field, F. H. Proton Affinities and Cluster Ion Stabilities in CO₂ and CS₂. Applications in Martian Ionspheric Chemistry, J. Chem. Phys. 66, 4527 (1977).	
Method used : MS Calculated entropy The measurements done in the temperature range of 400-650K. Using the Hoff plot the delta H and delta S at 300K were derived. Equilibrium, scale based on NH₃	

C2C13N	CAS REGN : 545-06-2
CL * * CL*C**C##N * * CL	
Acetonitrile, trichloro- (8CI9CI)	
GB (exp) : 169.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 177.5+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2D60	CAS REGN : 17222-37-6
C**O**C	
Methane-d3, oxybis- (9CI) (Methyl ether)-d6 (8CI)	
GB (exp) : ~186 kcal/mol	Temperature : 374 K
Blair, A. S., and Harrison, A. G., Bimolecular Reactions of Trapped Ions. V. Ionic Chemistry of Simple Oxygen-containing Molecules, Can. J. Chem. 51, 703 (1972).	
Method used : MS Equilibrium relative to CH3CHO. Found delta G ~3.	

C2HF3	CAS REGN : 359-11-5
F**C++C**F * * F	
Ethene, trifluoro- (9CI) Ethylene, trifluoro- (8CI)	
PA (exp) : 172+/-2 kcal/mol	Temperature : 300 K
Ridge, D. P., Gas Phase Proton Affinities of Several Fluoroethylenes, J. Am. Chem. Soc. 97, 5670 (1975).	
Method used : ICR Bracketing relative to H2O and H2S.	

C2HF3O2	CAS REGN : 76-05-1
F * * F**C**C%O * % * % F O	
Acetic acid, trifluoro- (8CI9CI)	
PA (exp) : 172+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to H2S and H2O	
GB (exp) : 154.7+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 171.9+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

Continued on next page

GB (exp) : 165.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 173.3+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H2C1N	CAS REGN : 107-14-2
CL*C**C##N	
Acetonitrile, chloro- (8CI9CI)	
GB (exp) : 173.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H2F2	CAS REGN : 75-38-7
F*xC++C * * F	
Ethene, 1,1-difluoro- (9CI) Ethylene, 1,1-difluoro- (8CI)	
PA (exp) : 178+/-4 kcal/mol	Temperature : 300 K
Ridge, D. P., Gas Phase Proton Affinities of several Fluoroethylenes, J. Am. Chem. Soc. 97, 5670 (1975).	
Method used : ICR Bracketing relative to C2H5I and CH3OH	
PA (exp) : 174.8 kcal/mol	Temperature : 298 K
Williamson, A. D., LeBrenton, P. R., and Beauchamp, J. L., Photoionization Mass Spectrometry of 2-Fluoropropane and 2,2-Difluoropropane. A Novel Determination of the Proton Affinity of Vinyl Fluoride and 1,1-Difluoro- ethylene, J. Am. Chem. Soc. 98, 2705 (1976).	
Method used : PI Appearance potential by phototization of (CH3)2CF2.	

C2H2F2	CAS REGN : 1630-77-9
F**C++C**F	
Ethene, 1,2-difluoro-, (Z)- (9CI) Ethylene, 1,2-difluoro-, (Z)- (8CI)	
PA (exp) : 166+/-4 kcal/mol	Temperature : 300 K
Ridge, D. P., Gas Phase Proton Affinities of several Fluoroethylenes, J. Am. Chem. Soc. 97, 5670 (1975).	
Method used : ICR Bracketing relative to C2H5F and H2O.	

C2H2F2	CAS REGN : 1630-78-0
F**C++C**F	
Ethene, 1,2-difluoro-, (E)- (9CI) Ethylene, 1,2-difluoro-, (E)- (8CI)	
PA (exp) : 167+/-4 kcal/mol	Temperature : 300 K
Ridge, D. P., Gas Phase Proton Affinities of several Fluoroethylenes, J. Am. Chem. Soc. 97, 5670 (1975).	
Method used : ICR Bracketing relative to C2H5F and H2O.	

C2H2O	CAS REGN : 463-51-4
O++C++C	
Ethenone (9CI) Ketene (8CI)	
PA (exp) : 194.1+/-1 kcal/mol	Temperature : 350 K
Ausloos, P., and Lias, S. G. The Proton Affinity of Ketene and the Heat of Formation of CH3CO+, Chem. Phys. Letters 51, 53 (1977).	
Method used : ICR Equilibrium; scale based on PA (iso-C4H8) = 193.5+/-1	
PA (exp) : 195.9+/-2 kcal/mol	Temperature : 300 K
Vogt, J., Williamson, A. D., and Beauchamp, J. L. Properties and Reactions of Ketene in the Gas Phase by Ion Cyclotron Resonance Spectroscopy and Photoionization Mass Spectrometry. Proton Affinity, Site Specificity of Protonation, and Heat of Formation of Ketene, J. Am. Chem. Soc. 100, 3478 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H3ClO2	CAS REGN : 79-11-8
O=C(Cl)C=O	
Acetic acid, chloro- (8CI9CI)	
GB (exp) : 166.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 184+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C2H3Cl3O	CAS REGN : 115-20-8
CL * * CL* <i>C**C**O</i> * * CL	
Ethanol, 2,2,2-trichloro- (8CI9CI)	
GB (exp) : 171.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 179+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H3F	CAS REGN : 75-02-5
F**C++C	
Ethene, fluoro- (9CI) Ethylene, fluoro- (8CI)	
PA (exp) : 174+/-1 kcal/mol	Temperature : 300 K
Ridge, D. P., Gas Phase Proton Affinities of several Fluoroethylenes, J. Am. Chem. Soc. 97, 5670 (1975).	
Method used : ICR Bracketing relative to H2S and CH2O.	
PA (exp) : 173.6 kcal/mol	Temperature : 298 K
Williamson, A. D., LeBreton, P. R., and Beauchamp, J. L., Photoionization Mass Spectrometry of 2-Fluoro- propane and 2,2-Difluoropropane. A Novel Determination of Vinyl Fluoride and 1,1-Difluoro- ethylene, J. Am. Chem. Soc. 98, 2705 (1976).	
Method used : PI Appearance potential by photoionization of (CH3)2CHF.	

C2H3F02	CAS REGN : 144-49-0
0% C * * C * * F % % 0	
Acetic acid, fluoro- (8CI9CI)	
GB (exp) : 167.3+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 184.5+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C2H3F3O	CAS REGN : 75-89-8
F * * F**C**C**O * * F	
Ethanol, 2,2,2-trifluoro- (8CI9CI)	
GB (exp) : 164.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 172.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H3N	CAS REGN : 75-05-8
N##C**C	
Acetonitrile (8CI9CI)	
PA (exp) : 189.5+/-4.5 kcal/mol	Temperature : 298 K
Gray, G. A., Study of Ion-Molecule Reactions and Reaction Mechanisms in Acetonitrile by Ion Cyclotron Resonance, J. Am. Chem. Soc. 90, 6002 (1968).	
Method used : ICR Bracketing relative to CH3CHO and (CH3)2CO	
PA (exp) : 186+/-1 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Translational energy of RH2+ has been measured assuming $\Delta H = 5X$ translational energy for reaction $RH^+ + RH \rightarrow RH_2^+ + R$	

Continued on next page

GB (exp) : 179.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 187+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 171.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 187.4+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C2H3NS	CAS REGN : 556-61-6
S++C++N**C	
Methane, isothiocyanato- (9CI) Isothiocyanic acid, methyl ester (8CI)	
PA (exp) : 189+/-6 kcal/mol	Temperature : 300 K
McAllister, T., Ion-Molecule Reactions and Proton Affinities of Methyl Thio- and Isothiocyanate, Int. J. Mass Spectrom. Ion Phys. 15, 303 (1974).	
Method used : ICR Bracketing relative to (CH3)2CO and 1-C4H8.	

C2H3NS	CAS REGN : 556-64-9
N##C**S**C	
Thiocyanic acid, methyl ester (8CI9CI)	
PA (exp) : 189+/-6 kcal/mol	Temperature : 300 K
McAllister, T., Ion-Molecule Reactions and Proton Affinities of Methyl Thio- and Isothiocyanate, Int. J. Mass Spectrom. Ion Phys. 15, 303 (1974).	
Method used : ICR Bracketing relative to (CH3)2CO and 1-C4H8.	

C ₂ H ₄	CAS REGN : 74-85-1
C++C	
Ethene (9CI) Ethylene (8CI)	
PA (exp) : 160.6 kcal/mol	Temperature : 298 K
Lossing F. P., and Semeluk, G. P., Free Radicals by Mass Spectrometry. XLII. Ionization Potentials and Ionic Heats of Formation for C ₁ -C ₄ Alkyl Radicals, Can. J. Chem. 48, 955 (1970).	
Method used : EM Appearance potential of C ₂ H ₅ ⁺ using EM.	

C2H4F2O	CAS REGN : 359-13-7
F**C**C**O * * F	
Ethanol, 2,2-difluoro- (8CI9CI)	
GB (exp) : 170.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 177.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H4F3N	CAS REGN : 753-90-2
F * * F**C**C**N * * F	
Ethanamine, 2,2,2-trifluoro- (9CI) Ethylamine, 2,2,2-trifluoro- (8CI)	
GB (exp) : 188.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 197.9+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 191.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200.3+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C2H4N2	CAS REGN : 540-61-4
N**C**C##N	
Acetonitrile, amino- (8CI9CI)	
PA (exp) : ~197 kcal/mol	Temperature : 300 K
Taagepera, M., Hehre, W. J., Topsom, R. D., and Taft, R. W. Calculation of Polar Substituent Parameters by ab initio Molecular Orbital Methods. Proton Affinities of Substituted Primary Amines, J. Am. Chem. Soc. 98, 7438 (1976).	
Method used : ICR Equilibrium; scale based on NH3	

C2H4O	CAS REGN : 75-07-0
O++C**C	
Acetaldehyde (8CI9CI)	
PA (exp) : 184+/-3 kcal/mol	Temperature :
Harrison, A. G., Ivko, A., and Van Raalte, D., Energetics of Formation of Some Oxygenated Ions and the Proton Affinities of Carbonyl Compounds, Can. J. Chem. 44, 1625 (1966).	
Method used : EI Electron impact fragmentation	
PA (exp) : 186+/-1 kcal/mol	Temperature : 298 K
Refaey, K. M. A., and Chupka, W. A., Photoionization of the Lower Aliphatic Alcohols with Mass Analysis, J. Chem. Phys. 48, 5205 (1968).	
Method used : PI Appearance potential by IP fragmentation of C2H5OH.	

Continued on next page

PA (exp) : 189.4 kcal/mol	Temperature : 298 K
Lossing, F. P. Heats of Formation of Some Isomeric $[C_nH_{2n+10}]^+$ Ions. Substitutional Effects on Ion Stability, J. Am. Chem. Soc. 99, 7526 (1977).	
Method used : EM Electron impact fragmentation (various molecules) using EM. $\Delta H_f(H^+) = 365.7$	
GB (exp) : 177.2 $\pm$ 2 kcal/mol	Temperature : 300 K
PA (exp) : 185 $\pm$ 2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

Continued on next page

GB (exp) : 169.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 185.4+/-2 kcal/mol	Temperature : 600 K
Yamdaqni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C2H4O	CAS REGN : 75-21-8
<pre> 0 * * * * C*****C </pre>	
Oxirane (9CI) Ethylene oxide (8CI)	
PA (exp) : 185+/-4 kcal/mol	Temperature :
Beauchamp, J. L., and Dunbar, R. C., Identification of C2H5O+ Structural Isomers by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 92, 1477 (1970).	
Method used : ICR Bracketing relative to CH3OH and (CH3)2O	

C2H4O2	CAS REGN : 64-19-7
O=C*O % % O	
Acetic acid (8CI9CI)	
PA (exp) : 188+/-3 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Equilibrium; scale based on NH3	
GB (exp) : 171.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 188.2+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 179.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H4O2	CAS REGN : 107-31-3
O=C*O	
Formic acid, methyl ester (8CI9CI)	
PA (exp) : 188+/-3 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to (CH3)2O and CH3CHO	

Continued on next page

GB (exp) : 179.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 187+/-2 kcal/mol	Temperature : 300 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on CH ₃ CHO Entropy change assumed zero	
GB (exp) : 179.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 187.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

Continued on next page

GB (exp) : 172.2+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 187.8+/-2 kcal/mol	Temperature : 600 K
Yamdaqni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C2H5Br	CAS REGN : 74-96-4
C**C**BR	
Ethane, bromo- (8CI9CI)	
PA (exp) : 171+/-2 kcal/mol	Temperature : 300 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also IP's of H and C2H5BR and assuming HA(CH3BR+)=HA(C2H5BR+).	

C2H5Cl	CAS REGN : 75-00-3
CL*C**C	
Ethane, chloro- (8CI9CI)	
PA (exp) : 170+/-2 kcal/mol	Temperature :
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also IP's of H and C2H5Cl and HA(CH3Cl+)=HA(C2H5Cl+).	

C2H5F	CAS REGN : 353-36-6
F**C**C	
Ethane, fluoro- (8CI9CI)	
PA (exp) : 165+/-5 kcal/mol	Temperature : 300 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also the IP's of H and C2H5F and assuming HA(CH3F+) = HA(C2H5F+).	

C2H5F2N	CAS REGN : 430-67-1
F**C**C**N * * F	
Ethanamine, 2,2-difluoro- (9CI) Ethylamine, 2,2-difluoro- (8CI)	
GB (exp) : 197.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 205.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H5I	CAS REGN : 75-03-6
I**C**C	
Ethane, iodo- (8CI9CI)	
PA (exp) : 175 kcal/mol	Temperature : 300 K
Beauchamp, J. L., Holtz, D., Woodgate, S. D., and Patt, S. L., Thermochemical Properties and Ion-Molecule Reactions of the Alkyl Halides in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 94, 2798 (1972).	
Method used : ICR Bracketing and using also IP's of H and assuming HA(CH3I+)=HA(C2H5I+).	

C2H5N	CAS REGN : 151-56-4
<pre> N * * * * C*****C </pre>	
Aziridine (9CI) Ethylenimine (8CI)	
GB (exp) : 204.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 212.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 204.9+/-2 kcal/mol	Temperature : 298 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	

Continued on next page

PA (exp) : 213+/-2 kcal/mol	Temperature : 300 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Nitrogen and Oxygen Bases. Hybridization Effects, J. Am. Chem. Soc. 97, 4137 (1975).	
Method used : ICR Equilibrium; scale based on CH₃NH₂ Van't Hoff plot	

C ₂ H ₅ NO	CAS REGN : 60-35-5
N%*C**C % % 0	
Acetamide (8CI9CI)	
GB (exp) : 188.5+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 205.7+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on CH₃NH₂ Entropy change assumed zero	

C2H5NO	CAS REGN : 123-39-7
O%CC%N**C	
Formamide; N-methyl- (8CI9CI)	
GB (exp) : 195+/-4 kcal/mol	Temperature : 300 K
PA (exp) : 202.8+/-4 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H5NO2	CAS REGN : 79-24-3
O++N**C**C + + O	
Ethane, nitro- (8CI9CI)	
PA (exp) : 186+/-2 kcal/mol	Temperature : 300 K
Kriemler, P., and Buttrill, S. E., Jr., Ion-Molecule Reactions and the Proton Affinities of the Nitroalkanes, J. Am. Chem. Soc. 95, 1365 (1973).	
Method used : ICR Bracketing relative to CH3CN and CH3CHO.	

C2H5NO3	CAS REGN : 625-58-1
C**C**O**N**+O + + O	
Nitric acid, ethyl ester (8CI9CI)	
PA (exp) : 181+/-3 kcal/mol	Temperature : 300 K
Kriemler, P., and Buttrill, S. E., Jr., Positive and Negative Ion-Molecule Reactions and the Proton Affinity of Ethyl Nitrate, J. Am. Chem. Soc. 92, 1123 (1970).	
Method used : ICR Bracketing relative to cyclo-C6H10 and CH3NO2.	

C2H6	CAS REGN : 74-84-0
C**C	
Ethane (8CI9CI)	
PA (exp) : 128+/-1 kcal/mol	Temperature :
Chong, S. L., and Franklin, J. L., Heats of Formation of Protonated Cyclopropane, Methylcyclopropane, and Ethane, J. Am. Chem. Soc. 94, 6347 (1972).	
Method used : MS By applying path degeneracy correction found ΔH formation (C2H7+) = 218.8 Equilibrium relative to CH4; assume $\Delta S = 0$.	
PA (exp) : 139.6 (1) kcal/mol	Temperature : 300 K
Hiraoka, K., and Kebarle, P., Stabilities and Energetics of Pentacoordinated Carbonium Ions. The Isomeric C2H7+ Ions and Some Higher Analogues: C3H9+ and C4H11+, J. Am. Chem. Soc. 98, 6119 (1976).	
Method used : MS Based on ΔH formation(C2H5+) = 219. Equilibrium of reaction: C2H5+ + H2 -> C2H7+(C-C protonation) (1).	

Continued on next page

PA (exp) : 131.8 (2) kcal/mol	Temperature :
Hiraoka, K., and Kebarle, P., Stabilities and Energetics of Pentacoordinated Carbonium Ions. The Isomeric C₂H₇⁺ Ions and Some Higher Analogues: C₃H₉⁺ and C₄H₁₁⁺, J. Am. Chem. Soc. 98, 6119 (1976).	
Method used : MS Based on delta H formation(C₂H₅⁺) = 219 Equilibrium of reaction: C₂H₅⁺ + H₂ + H₂ → C₂H₇⁺ + H₂(CH protonation) (2).	

C ₂ H ₆ FN	CAS REGN : 406-34-8
F**C**C**N	
Ethanamine, 2-fluoro- (9CI) Ethylamine, 2-fluoro- (8CI)	
GB (exp) : 201.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 210.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH₃	

C2H6N2	CAS REGN : 4143-41-3
C**N++N**C	
Diazene, dimethyl-, (E)- (9CI) Azomethane, (E)- (8CI)	
GB (exp) : 197.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 204.5+/-2 kcal/mol	Temperature : 300 K
Foster, M. S., Williamson, A. D., and Beauchamp, J. L., Photoionization Mass Spectrometry of trans-Azomethane, Int. J. Mass Spectrom. Ion Phys. 15, 429 (1974).	
Method used : MS Calculated entropy Equilibrium relative to NH3	
PA (exp) : 206.8+/-5 kcal/mol	Temperature : 300 K
Foster, M. S., and Beauchamp, J. L., Gas-Phase Ion Chemistry of Azomethane by Ion Cyclotron Resonance, J. Am. Chem. Soc. 94, 2425 (1972).	
Method used : ICR Bracketing between NH3 and CH3NH2	

C2H6O	CAS REGN : 64-17-5
O**C**C	
Ethanol (9CI) Ethyl alcohol (8CI)	
PA (exp) : 186+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH3COOH and CH3CHO	
GB (exp) : 178.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 186.2+/-2 kcal/mol	Temperature : 300 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on CH3CHO Entropy change assumed zero	

Continued on next page

GB (exp) : 171.2+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 186.8+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 178.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 186.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H6O	CAS REGN : 115-10-6
C**O**C	
Methane, oxybis- (9CI) Methyl ether (8CI)	
PA (exp) : 190+/-4 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to I-C4H8 and CH3CHO; Scale based on NH3	
PA (exp) : 189 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Delta H assume = 5X translational energy of the products of reaction $RH + RH^+ \rightarrow RH_2^+ + R$.	

Continued on next page

GB (exp) : 181.3+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 189.1+/-2 kcal/mol	Temperature : 298 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on MECHO Entropy change assumed zero	
GB (exp) : 174.5+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 189.3+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 182.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 193.3 kcal/mol	Temperature : 300 K
Hiraoka, K., Grimsrud, E. P., and Kebarle, P., Gas Phase Ion Equilibria Studies of the Hydrogen Ion in Water-Dimethyl Ether and Methanol-Dimethyl, J. Am. Chem. Soc. 96, 3359 (1974).	
Method used : MS Van't Hoff plot Equilibrium relative to H2O. Delta G determined by means of thermodynamic cycles involving mixed clusters.	

C2H6OS	CAS REGN : 67-68-5
C**S**C + + 0	
Methane, sulfinylbis- (9CI) Methyl sulfoxide (8CI)	
GB (exp) : 200.9+/-2.5 kcal/mol	Temperature : 300 K
PA (exp) : 208.3+/-2.5 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 184+/-3 kcal/mol	Temperature : 300 K
McAllister, T. Ion Cyclotron Resonance Mass Spectroscopy of Dimethyl Sulfoxide, Intern. J. Mass Spectrom. Ion Phys. 25, 353 (1977).	
Method used : ICR Bracketing relative to CH3OH and EtOH.	

C2H6S	CAS REGN : 75-08-1
S**C**C	
Ethanethiol (8CI9CI)	
GB (exp) : 180.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H6S	CAS REGN : 75-18-3
C**S**C	
Methane, thiobis- (9CI) Methyl sulfide (8CI)	
GB (exp) : 190.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.6+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H7N	CAS REGN : 75-04-7
N**C**C	
Ethanamine (9CI) Ethylamine (8CI)	
GB (exp) : 205.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 214.0+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 205.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 214.3+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C2H7N	CAS REGN : 124-40-3
C**N**C	
Methanamine, N-methyl- (9CI) Dimethylamine (8CI)	
GB (exp) : 209.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 217.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 209.5+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 217.7+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

Continued on next page

GB (exp) : 203.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 220.6+/-2 kcal/mol	Temperature : 600 K
Briggs, J. P., Yamdagni, R., and Kebarle, P., Intrinsic Basicities of Ammonia, Methylamines, Anilines, and Pyridine from Gas-Phase Proton- Exchange Equilibria, J. Am. Chem. Soc. 94, 5128 (1972).	
Method used : MS Entropy change assumed zero Equilibrium; scale based on NH3	

C2H7P	CAS REGN : 676-59-5
C**P**C	
Phosphine, dimethyl- (8CI9CI)	
GB (exp) : 205.6+/-2 kcal/mol	Temperature : 298 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	

C2H8N2	CAS REGN : 57-14-7
C**N**C * * N	
Hydrazine, 1,1-dimethyl- (8CI9CI)	
GB (exp) : 209.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 217.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C2H8N2	CAS REGN : 107-15-3
N**C**C**N	
1,2-Ethanediamine (9CI) Ethylenediamine (8CI)	
GB (exp) : 213.60+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.50+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	
GB (exp) : 215.50+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 227.50+/-2 kcal/mol	Temperature : 298 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on (CH3)3N Van't Hoff plot	

C3D9N	CAS REGN : 13960-80-0
C**N**C * * C	
Methan-d3-amine, N,N-di(methyl-d3)- (9CI) Trimethylamine-d9. (8CI)	
GB (exp) : 214.00+/-2 kcal/mol	Temperature : 298 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	

C3H2N2	CAS REGN : 109-77-3
N##C**C**C##N	
Propanedinitrile (9CI) Malononitrile (8CI)	
GB (exp) : 169.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 177.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H3F3O2	CAS REGN : 431-47-0
<pre> F * * C**O**C**C**F + * + * O F </pre>	
Acetic acid, trifluoro-, methyl ester (8CI9CI)	
GB (exp) : 172.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.3+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H3F3O2	CAS REGN : 32042-38-9
F * * F**C**C**O**C++O * * F	
Ethanol, 2,2,2-trifluoro-, formate (9CI)	
GB (exp) : 173+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.8+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H3N	CAS REGN : 107-13-1
C++C**C##N	
2-Propenenitrile (9CI) Acrylonitrile (8CI)	
GB (exp) : 180.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 187.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 188.3+/-2.5 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to C3H6O2(1,3-Dioxolane) with its PA value = 189.8; relative to Et CHO and MeCN	

C3H4	CAS REGN : 74-99-7
C##C**C	
1-Propyne (9CI) Propyne (8CI)	
PA (exp) : 176+/-2 kcal/mol	Temperature : 300 K
Aue, D. H., Davidson, W. R., and Bowers, M. T., Heats of Formation of C3H5+ Ions. Allyl, Vinyl, and Cyclopropyl Cations in Gas-Phase Proton-Transfer Reactions, J. Am. Chem. Soc. 98, 6700 (1976).	
Method used : ICR Entropy change assumed zero Equilibrium; scale based on H2S.	

C3H4	CAS REGN : 463-49-0
C++C++C	
1,2-Propadiene (9CI) Allene (8CI)	
PA (exp) : 176+/-2 kcal/mol	Temperature : 300 K
Aue, D. H., Davidson, W. R., and Bowers, M. T., Heats of Formation of C3H5+ Ions. Allyl, Vinyl, and Cyclopropyl Cations in Gas-Phase Proton-Transfer Reactions, J. Am. Chem. Soc. 98, 6700 (1976).	
Method used : ICR Equilibrium; scale based on H2S. Entropy change assumed zero.	

C3H4	CAS REGN : 2781-85-3
<pre> C * + * + C*****C </pre>	
Cyclopropene (8CI9CI)	
PA (exp) : 194+/-3 kcal/mol	Temperature : 300 K
Aue, D. H., Davidson, W. R., and Bowers, M. T., Heats of Formation of C3H5+ Ions. Allyl, Vinyl, and Cyclopropyl Cations in Gas-Phase Proton-Transfer Reactions, J. Am. Chem. Soc. 98, 6700 (1976).	
Method used : ICR Bracketing relative to (CH3)2O and (C2H5)2O.	

C3H4ClN	CAS REGN : 542-76-7
N##C**C**C**CL	
Propanenitrile, 3-chloro- (9CI) Propionitrile, 3-chloro- (8CI)	
PA (exp) : 181.1+/-2 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to MeCN	
GB (exp) : 178.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 186.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H5N	CAS REGN : 107-12-0
C**C**C##N	
Propanenitrile (9CI) Propionitrile (8CI)	
GB (exp) : 182.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 190.2+/-2.5 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to n-PrCN, HCO2Et, i-PrCHO, Me2O, EtCHO and relative to 1,3-Dioxane using its PA = 189.8	

C3H5N	CAS REGN : 2450-71-7
C##C**C**N	
2-Propyn-1-amine (9CI) 2-Propynylamine (8CI)	
GB (exp) : 200.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 209.0+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6	CAS REGN : 75-19-4
<pre> C * * * * C*****C </pre>	
Cyclopropane (8CI9CI)	
PA (exp) : 180.6+/-4 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Heats of Formation of Protonated Cyclopropane, Methylcyclopropane, and Ethane, J. Am. Chem. Soc. 94, 6347 (1972).	
Method used : MS $\text{HC(OH)}_2^+ + \text{C-C}_3\text{H}_6 \rightarrow \text{HC(OH)}_2^+ + \text{C-C}_3\text{H}_7^+$ Equilibrium of reactions: $\text{CH}_3\text{OH}_2^+ + \text{C-C}_3\text{H}_6 \rightarrow \text{CH}_3\text{OH} + \text{C-C}_3\text{H}_7^+$; Apply correction for path degeneracy. Assuming $\Delta S = 0$.	

C3H6	CAS REGN : 115-07-1
C++C**C	
1-Propene (9CI) Propene (8CI)	
PA (exp) : 180 kcal/mol	Temperature : 298 K
Lossing F. P., and Semeluk, G. P. Free Radicals by Mass Spectrometry. XLII. Ionization Potentials and Ionic Heats of Formation for C1-C4 Alkyl Radicals, Can. J. Chem. 48, 955 (1970).	
Method used : EM Appearance potential of C3H7+.	
PA (exp) : 182.2+/-2 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Heats of Formation of Protonated Cyclopropane, Methylcyclopropane, and Ethane, J. Am. Chem. Soc. 94, 6347 (1972).	
Method used : MS Correction for path degeneracy. Equilibrium relative to CH3OH; assuming delta S = 0.	

Continued on next page

GB (exp) : 165.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 182.6+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C3H6F3N	CAS REGN : 460-39-9
<pre> F * * N**C**C**C**F * * F </pre>	
1-Propanamine, 3,3,3-trifluoro- (9CI) Propylamine, 3,3,3-trifluoro- (8CI)	
GB (exp) : 200.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 209.0+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6F3N	CAS REGN : 677-41-8
<pre> F C * * * * F**C**N**C * * F </pre>	
Methanamine, 1,1,1-trifluoro-N,N-dimethyl-	
PA (exp) : 192+/-3 kcal/mol	Temperature : 300 K
Staley, R. H., Taagepera, M., Henderson, W. G., Koppel, I., Beauchamp, J. L., and Taft, R. W. Effects of Alkyl and Fluoroalkyl Substitution on the Heterolytic and Homolytic Bond Dissociation Energies of Protonated Amines, J. Am. Chem. Soc. 99, 326 (1977).	
Method used : ICR Calculated entropy Equilibrium scale based on NH3	

C3H6F3N	CAS REGN : 2730-67-8
F x x C*xN*xC*xC*xF x x F	
Ethanamine, 2,2,2-trifluoro-N-methyl- (9CI) Ethylamine, 2,2,2-trifluoro-N-methyl- (8CI)	
GB (exp) : 199.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 207.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6N2	CAS REGN : 151-18-8
N##C**C**C**N	
Propanenitrile, 3-amino- (9CI) Propionitrile, 3-amino- (8CI)	
GB (exp) : 196.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 204.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6N2	CAS REGN : 5616-32-0
C**N**C**C##N	
Acetonitrile, (methylamino)- (8CI9CI)	
GB (exp) : 196.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 204.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6O	CAS REGN : 67-64-1
C**C**C + + O	
2-Propanone (9CI) Acetone (8CI)	
PA (exp) : 195.7+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to (C2H5)2O and CH2CO	
PA (exp) : 193.6+/-1 kcal/mol	Temperature : 350 K
Ausloos, P., and Lias, S. G. The Proton Affinity of Ketene and the Heat of Formation of CH3CO+, Chem. Phys. Letters 51, 53 (1977).	
Method used : ICR Equilibrium; scale based on PA (iso-C4H8) = 193.5+/-1	

Continued on next page

GB not calculated for this reference	
PA (exp) : 188 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Delta H estimated equal 5 times translational energy of the products.	
PA (exp) : 196 kcal/mol	Temperature : 298 K
Heats of Formation of Some Isomeric $[C_nH_{2n+10}]^+$ Ions. Substitutional Effects on Ion Stability, J. Am. Chem. Soc. 99, 7526 (1977).	
Method used : EM Electron impact fragmentation (various molecules) using EM. $\Delta H_f(H^+) = 365.7$ Lossing, F. P.	

Continued on next page

GB (exp) : 179+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 193.8+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 186.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 193.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6O	CAS REGN : 123-38-6
C**C**C++O	
Propanal (9CI) Propionaldehyde (8CI)	
PA (exp) : 190+/-3 kcal/mol	Temperature :
Harrison, A. G., Ivko, A., and Van Raalte, D., Energetics of Formation of Some Oxygenated Ions and the Proton Affinities of Carbonyl Compounds, Can. J. Chem. 44, 1625 (1966).	
Method used : EI Electron impact fragmentation of various molecules.	
PA (exp) : 187 kcal/mol	Temperature : 298 K
Lossing, F. P. Heats of Formation of Some Isomeric [C _n H _{2n} + 10] ⁺ Ions. Substitutional Effects on Ion Stability, J. Am. Chem. Soc. 99, 7526 (1977).	
Method used : EM Electron impact fragmentation (various molecules) using EM. $\Delta H_f(H^+) = 365.7$	

Continued on next page

GB (exp) : 172.3+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 187.9+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 180.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6O	CAS REGN : 503-30-0
C**O * * * * C**C	
Oxetane (9CI) Trimethylene oxide (8CI)	
PA (exp) : 194+/-1 kcal/mol	Temperature : 300 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Nitrogen and Oxygen Bases. Hybridization Effects, J. Am. Chem. Soc. 97, 4137 (1975).	
Method used : ICR Semiquantitative. Absolute value assign relative to CH3CHO and (CH3)2C=CH2.	

C3H6OS	CAS REGN : 1534-08-3
C**C**S**C + + 0	
Ethanethioic acid, S-methyl ester (9CI) Acetic acid, thio-, S-methyl ester (8CI)	
PA (exp) : ~197+/-2 kcal/mol	Temperature : 300 K
Grunwell, J. R., Foerst, D. L., Kaplan, F., and Siddiqui, J. The Relative Basicity of Sulfur containing Esters, Tetrahedron 33, 2781 (1977).	
Method used : ICR Bracketing relative to MeCOOMe and MeCONHCH3. Also found the reactions of Me2S with MeCOOMe and MeCOSMe are both thermoneutral concluded PA (CH3COSCH3) is within 2 kcal of that of PA (MeCOOMe)	

C3H6O2	CAS REGN : 79-09-4
O=C*CC* % % 0	
Propanoic acid (9CI) Propionic acid (8CI)	
GB (exp) : 173.9+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 191.1+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C3H6O2	CAS REGN : 79-20-9
C**C**O**C + + O	
Acetic acid, methyl ester (8CI9CI)	
PA (exp) : 194+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to ME2CO and CH2CO	
GB (exp) : 186.7+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 194.1+/-2 kcal/mol	Temperature : 298 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on (CH3)2CO Entropy change assumed zero	

Continued on next page

GB (exp) : 187.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 195.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 179.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 195.4+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C3H6O2	CAS REGN : 109-94-4
O++C**O**C**C	
Formic acid, ethyl ester (8CI9CI)	
PA (exp) : 194+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH2CO AND I-C4H8; scale based on NH3.	
GB (exp) : 183.7+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 191.1+/-2 kcal/mol	Temperature : 298 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on (CH3)2CO Entropy change assumed zero	

Continued on next page

GB (exp) : 175.6+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 191.2+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 183.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 191.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6O3	CAS REGN : 616-38-6
C**O**C**O**C + + O	
Carbonic acid, dimethyl ester (8CI9CI)	
GB (exp) : 189.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H6S2	CAS REGN : 2168-84-5
$\begin{array}{c} \text{S} \\ + \\ + \\ \text{C}^{\times}\text{S}^{\times}\text{C}^{\times}\text{C}^{\times}\text{C} \end{array}$	
Ethane(dithioic)acid, methyl ester	
PA (exp) : >197+/-2 kcal/mol	Temperature : 300 K
Grunwell, J. R., Forest, D. L., Kaplan, F., and Siddiqui, J. The Relative Basicity of Sulfur Containing Esters, Tetrahedron 33, 2781 (1977).	
Method used : ICR Double resonance technique shows PA (MeCSSMe) > PA (the COSMe)	

C3H7N	CAS REGN : 107-11-9
C++C**C**N	
2-Propen-1-amine (9CI) Allylamine (8CI)	
GB (exp) : 205.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 213.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H7N	CAS REGN : 503-29-7
C**N * * * * C**C	
Azetidine (8CI9CI)	
GB (exp) : 211.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 220.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 212.2+/-2 kcal/mol	Temperature : 290 K
PA (exp) : 220.+/-2 kcal/mol	Temperature : 290 K
Bowers, M. T., Aue, D. H., and Webb, H. M., Equilibrium Constants for Gas-Phase Ionic Reactions. Accurate Determination of Relative Proton Affinities, J. Am. Chem. Soc. 93, 4314 (1971).	
Method used : ICR Entropy change assumed zero Equilibrium; scale based on (CH3)3N	

C3H7N	CAS REGN : 1072-44-2
<pre> C * * * * N*****C * * C </pre>	
Aziridine, 1-methyl- (8CI9CI)	
PA (exp) : 219.4+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Nitrogen and Oxygen Bases. Hybridization Effects, J. Am. Chem. Soc. 97, 4137 (1975).	
Method used : ICR Equilibrium; scale based on CH3NH2	

C3H7NO	CAS REGN : 68-12-2
C**N**C++O * * C	
Formamide, N,N-dimethyl- (8CI9CI)	
GB (exp) : 201.2+/-3 kcal/mol	Temperature : 300 K
PA (exp) : 209.0+/-3 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H8	CAS REGN : 74-98-6
C**C**C	
Propane (8CI9CI)	
PA (exp) : 148.2 kcal/mol	Temperature : 163-213 K
Hiraoka, K., and Kebarle, P., Information on the Proton Affinity and Protolysis of Propane from Measurement of the Ion Cluster, Equilibrium: $C_2H_5^+ + CH_4 = C_3H_9^+$. Can. J. Chem. 53, 970 (1975).	
Method used : MS Equilibrium of reaction $C_2H_5^+ + 2CH_4 \rightarrow C_3H_9^+ + CH_4$ Van't Hoff plot	
PA (exp) : 143 kcal/mol	Temperature : 170 K
Field, F. H., and Beggs, D. P., Reversible Reactions of Gaseous Ions. III. Studies with Methane at 0.1-1.0 Torr and 77-300 K, J. Am. Chem. Soc. 93, 1585 (1971).	
Method used : MS Equilibrium of reaction: $C_2H_5^+ + CH_4 \rightarrow C_3H_9^+$. Based on delta H formation($C_2H_5^+$). Calculated entropy	

Continued on next page

GB not calculated for this reference	
PA (exp) : <152.8 kcal/mol	Temperature : -170 K
Hiraoka, K., and Kebarle, P., Stabilities and Energetics of Pentacoordinated Carbonium Ions. The Isomeric C ₂ H ₇ ⁺ Ions and Some Higher Analogues: C ₃ H ₉ ⁺ and C ₄ H ₁₁ ⁺ , J. Am. Chem. Soc. 98, 6119 (1976).	
Method used : MS Calculated entropy Equilibrium of reaction Sec-C ₃ H ₇ ⁺ + H ₂ -> C ₃ H ₉ ⁺ Based on delta H formation(C ₃ H ₇ ⁺); C-H protonation	

C ₃ H ₈ O	CAS REGN : 67-63-0
C**C**C * * 0	
2-Propanol (9CI) Isopropyl alcohol (8CI)	
PA (exp) : 192+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to I-C ₄ H ₈ and (CH ₃) ₂ O	

Continued on next page

GB not calculated for this reference	
PA (exp) : 190 kcal/mol	Temperature :
Hiraoka, K., and Kebarle, P. Condensation Reactions Involving Carbonium Ions and Lewis Bases in the Gas Phase. Hydration of the tert-Butyl Cation, J. Am. Chem. Soc. 99, 360 (1977).	
Method used : MS Equilibrium relative to known standard acids	

C3H8O	CAS REGN : 71-23-8
C**C**C**O	
1-Propanol (9CI) Propyl alcohol (8CI)	
GB (exp) : 181.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

PA (exp) : 189+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to C ₂ H ₅ OH and (CH ₃) ₂ O	

C ₃ H ₈ O	CAS REGN : 540-67-0
C**C**O**C	
Ethane, methoxy- (9CI) Ether, ethyl methyl (8CI)	
GB (exp) : 186.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 194+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

C3H8S	CAS REGN : 75-33-2
C**C**C * * S	
2-Propanethiol (8CI9CI)	
GB (exp) : 184.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 192+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H8S	CAS REGN : 107-03-9
C**C**C**S	
1-Propanethiol (8CI9CI)	
GB (exp) : 181.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H8S	CAS REGN : 624-89-5
C**C**S**C	
Ethane, (methylthio)- (9CI) Sulfide, ethyl methyl (8CI)	
GB (exp) : 192.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C3H9As	CAS REGN : 593-88-4
C**AS*C * * C	
Arsine, trimethyl- (8CI9CI)	
PA (exp) : 210.6 kcal/mol	Temperature : 300 K
Hodges, R. V., and Beauchamp, J. L., Basicity and Ion-Molecule Reactions of Trimethylarsine in the Gas Phase Determined by Ion Cyclotron Resonance Spectroscopy, Inorg. Chem. 14, 2887 (1975).	
Method used : ICR Equilibrium relative to CH3NH2 and also to CF3(CH2)3NH2.	

C3H9N	CAS REGN : 75-31-0
C**C**C * * N	
2-Propanamine (9CI) Isopropylamine (8CI)	
GB (exp) : 208.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 216.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 208.2+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.6+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C3H9N	CAS REGN : 75-50-3
C**N**C * * C	
Methanamine, N,N-dimethyl- (9CI) Trimethylamine (8CI)	
GB (exp) : 214.30+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 222.10+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 213.70+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 221.50+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

Continued on next page

GB (exp) : 208.40+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 225.60+/-2 kcal/mol	Temperature : 600 K
Briggs, J. P., Yamdagni, R., and Kebarle, P., Intrinsic Basicities of Ammonia, Methylamines, Anilines, and Pyridine from Gas-Phase Proton- Exchange Equilibria, J. Am. Chem. Soc. 94, 5128 (1972).	
Method used : MS Entropy change assumed zero Equilibrium; scale based on NH3	

C3H9N	CAS REGN : 107-10-8
C**C**C**N	
1-Propanamine (9CI) Propylamine (8CI)	
GB (exp) : 207.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 215.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 207.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 215.7+/-2 kcal/mol	Temperature : 300 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C3H9N	CAS REGN : 624-78-2
C**C**N**C	
Ethanamine, N-methyl- (9CI) Ethylamine, N-methyl- (8CI)	
GB (exp) : 212.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 220.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page


GB (exp) : 212+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 220.2+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C3H9NO	CAS REGN : 109-85-3
C**O**C**C**N	
Ethanamine, 2-methoxy- (9CI) Ethylamine, 2-methoxy- (8CI)	
GB (exp) : 209.3+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 221.0+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C3H9P	CAS REGN : 594-09-2
C**P**C * * C	
Phosphine, trimethyl- (8CI9CI)	
GB (exp) : 214.9+/-2 kcal/mol	Temperature : 298 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	
PA (exp) : 223.3+/-2 kcal/mol	
Temperature : 300 K	
Staley, P. H., and Beauchamp, J. L. Basicities and Ion-Molecule Reactions of the Methylphosphines in the Gas Phase by Ion Cyclotron Resonance Spectroscopy, J. Am. Chem. Soc. 96, 6252 (1974).	
Method used : ICR Bracketing relative to (CH3)3N and (CH3)2C2H5N	


C3H10N2	CAS REGN : 109-76-2
N**C**C**C**N	
1,3-Propanediamine (8CI9CI)	
GB (exp) : 219.10+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 231.60+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	
GB (exp) : 221.10+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 235.10+/-2 kcal/mol	Temperature : 298 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on (CH3)3N Van't Hoff plot	


C4H4F4O2	CAS REGN : 1683-88-1
F * * F**C**C**O**C**C**F * + * + F 0	
Acetic acid, trifluoro-, 2-fluoroethyl ester (8CI)	
GB (exp) : 172.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H4N2	CAS REGN : 289-80-5
	
Pyridazine (8CI9CI)	
GB (exp) : 206.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 213.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H4N2	CAS REGN : 289-95-2
	
Pyrimidine (8CI9CI)	
GB (exp) : 201.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 208.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H4N2	CAS REGN : 290-37-9
	
Pyrazine (8CI9CI)	
GB (exp) : 199.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 206.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H4N2O	CAS REGN : 557-01-7
	
2(1H)-Pyrimidinone (8CI9CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to NH3 and CH3NH2.	

C4H4N2O2	CAS REGN : 66-22-8
	
2,4(1H,3H)-Pyrimidinedione (9CI) Uracil (8CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to NH3 and CH3NH2.	


C4H4O	CAS REGN : 110-00-9
<pre> C*****O + * + * C C * + * + C </pre>	
Furan (8CI9CI)	
PA (exp) : 189.8+/-2 kcal/mol	Temperature : 300 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Entropy change assumed zero Equilibrium relative to (CH3)2O.	

C4H5F3O2	CAS REGN : 383-63-1
<pre> F * * F**C**C**O**C**C * + * + F O </pre>	
Acetic acid, trifluoro-, ethyl ester (8CI9CI)	
GB (exp) : 175.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 183.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H5N	CAS REGN : 109-97-7
<pre> C*****N + * + * C C * + * + C </pre>	
1H-Pyrrole (9CI) Pyrrole (8CI)	
GB (exp) : 197.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 205.9+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	
GB (exp) : 192.2+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 209.4+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C4H5N	CAS REGN : 5500-21-0
<pre> C * * * * * * C*****C * * N##C </pre>	
Cyclopropanecarbonitrile (8CI9CI)	
PA (exp) : 193.2+/-2 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to HC02-n-C4H9	
GB (exp) : 185.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 193.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H5NO2	CAS REGN : 623-49-4
N##C**C**O**C**C + + 0	
Carbonocyanidic acid, ethyl ester (9CI) Formic acid, cyano-, ethyl ester (8CI)	
GB (exp) : 173.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 180.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H5N3O	CAS REGN : 71-30-7
	
2(1H)-Pyrimidinone, 4-amino- (9CI) Cytosine (8CI)	
PA (exp) : 220+/-2 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)2NH and (CH3)3N.	

C4H6N2	CAS REGN : 616-47-7
<pre> C+++++N * * * * C**N C * + * + C </pre>	
1H-Imidazole, 1-methyl- (9CI) Imidazole, 1-methyl- (8CI)	
GB (exp) : 217.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 225.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H7F3O	CAS REGN : 461-24-5
F * * F**C**C**O**C**C * * F	
Ethane, 2-ethoxy-1,1,1-trifluoro- (9CI) Ether, ethyl 2,2,2-trifluoroethyl (8CI)	
GB (exp) : 177.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 184.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H7N	CAS REGN : 78-82-0
C**C**C##N * * C	
Propanenitrile, 2-methyl- (9CI) Isobutyronitrile (8CI)	
PA (exp) : 192.2+/-2 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to HC02C2H5	
GB (exp) : 184.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 192.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H7N	CAS REGN : 109-74-0
N##C**C**C**C	
Butanenitrile (9CI) Butyronitrile (8CI)	
PA (exp) : 191.2+/-2.5 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to HC02Et, ME2O and EtCN	
GB (exp) : 183.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 191.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8	CAS REGN : 106-98-9
C**C**C++C	
1-Butene (8CI9CI)	
PA (exp) : 183.2 kcal/mol	Temperature : 298 K
Lossing F. P., and Semeluk, G. P. Free Radicals by Mass Spectrometry. XLII. Ionization Potentials and Ionic Heats of Formation for C1-C4 Alkyl Radicals, Can. J. Chem. 48, 955 (1970).	
Method used : EM Appearance potential of C4H9+ by EM	

C4H8	CAS REGN : 115-11-7
C**C**C + + C	
1-Propene, 2-methyl- (9CI) Propene, 2-methyl- (8CI)	
PA (exp) : 192.6+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Using $\Delta H_f(t-C_4H_9^+) = 169 \pm 1.8$ derived from, $\Delta H_f(t-C_4H_9) = 9.3$ and $IP(t-C_4H_9) = 159.9$	
PA (exp) : 193.7 kcal/mol	Temperature :
Lossing F. P., and Semeluk, G. P. Free Radicals by Mass Spectrometry. XLII. Ionization Potentials and Ionic Heats of Formation for C1-C4 Alkyl Radicals, Can. J. Chem. 48, 955 (1970).	
Method used : EM Appearance potential of $C_4H_9^+$	

Continued on next page

GB (exp) : 177.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 194.2+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 185.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 193.5+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8	CAS REGN : 590-18-1
C**C++C**C	
2-Butene, (Z)- (8CI9CI)	
PA (exp) : 181.9 kcal/mol	Temperature : 298 K
Lossing F. P., and Semeluk, G. P. Free Radicals by Mass Spectrometry. XLII. Ionization Potentials and Ionic Heats of Formation for C1-C4 Alkyl Radicals, Can. J. Chem. 48, 955 (1970).	
Method used : EM Appearance potential of C4H9+	

C4H8	CAS REGN : 594-11-6
<pre> C * * * * C*****C * * C </pre>	
Cyclopropane, methyl- (8CI9CI)	
PA (exp) : 181+/-2 kcal/mol	Temperature :
Chong, S. L., and Franklin, J. L., Heats of Formation of Protonated Cyclopropane, Methylcyclopropane, and Ethane, J. Am. Chem. Soc. 94, 6347 (1972).	
Method used : MS Equilibrium relative to HCOOH based on PA(HCOOH)=180.1+/-2 assuming delta S=0 Correction for path degeneracy.	

C4H8	CAS REGN : 624-64-6
C**C++C**C	
2-Butenè, (E)- (8CI9CI)	
PA (exp) : 180.9 kcal/mol	Temperature : 298 K
Lossing F. P., and Semeluk, G. P. Free Radicals by Mass Spectrometry. XLII. Ionization Potentials and Ionic Heats of Formation for C1-C4 Alkyl Radicals, Can. J. Chem. 48, 955 (1970).	
Method used : EM Appearance potential of C4H9+	

C4H8F3N	CAS REGN : 819-06-7
<pre> F * * C**N**C**C**F * * * * C F </pre>	
Ethanamine, 2,2,2-trifluoro-N,N-dimethyl- (9CI) Ethylamine, 2,2,2-trifluoro-N,N-dimethyl- (8CI)	
GB (exp) : 204.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 212.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8F3N	CAS REGN : 819-46-5
F * * F**C**C**C**C**N * * F	
1-Butanamine, 4,4,4-trifluoro- (9CI) Butylamine, 4,4,4-trifluoro- (8CI)	
GB (exp) : 203.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 212.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8N2	CAS REGN : 926-64-7
N##C**C**N**C * * C	
Acetonitrile, (dimethylamino)- (8CI9CI)	
GB (exp) : 200.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 208.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8O	CAS REGN : 78-84-2
C**C**C++O * * C	
Propanal, 2-methyl- (9CI) Isobutyraldehyde (8CI)	
GB (exp) : 182.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.6+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8O	CAS REGN : 78-93-3
C**C**C**C + + 0	
2-Butanone (8CI9CI)	
GB (exp) : 189.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 193.7+/-1 kcal/mol	Temperature : 293 K
Lossing, F. P. Heats of Formation of Some Isomeric $[C_nH_{2n} + 10]^+$ Ions. Substitutional Effects on Ion Stability, J. Am. Chem. Soc. 99, 7526 (1977).	
Method used : EM Electron impact fragmentation (various molecules) using EM. $\Delta H_f(H^+) = 365.7$	

C4H8O	CAS REGN : 109-99-9
<pre> C*****O * * * * C C * * * * C </pre>	
Furan, tetrahydro- (8CI9CI)	
GB (exp) : 189+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 196.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8O	CAS REGN : 123-72-8
O++C**C**C**C	
Butanal (9CI) Butyraldehyde (8CI)	
GB (exp) : 181.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8O2	CAS REGN : 110-74-7
C**C**C**O**C++O	
Formic acid, propyl ester (8CI9CI)	
PA (exp) : 194+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH2CO AND I-C4H8; scale based on NH3.	
GB (exp) : 184.7+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 192.1+/-2 kcal/mol	Temperature : 298 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on (CH3)2CO Entropy change assumed zero	

Continued on next page

GB (exp) : 184.5 kcal/mol	Temperature : 300 K
PA (exp) : 192.3+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 176.1+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 191.7+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C4H8O2	CAS REGN : 123-91-1
<pre> O**C * * * * C C * * * * C**O </pre>	
1,4-Dioxane (9CI) p-Dioxane (8CI)	
GB (exp) : 183.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8O2	CAS REGN : 141-78-6
C**C**O**C**C + + O	
Acetic acid ethyl ester (8CI9CI)	
PA (exp) : 197+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to (C2H5)2O and CH3CO2ME.	
GB (exp) : 188.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 196+/-2 kcal/mol	Temperature : 300 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on (CH3)2CO Entropy change assumed zero	

Continued on next page

GB (exp) : 190.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 198.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 182.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 198+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C4H8O2	CAS REGN : 554-12-1
C**O**C**C**C + + O	
Propanoic acid, methyl ester (9CI) Propionic acid, methyl ester (8CI)	
GB (exp) : 189.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 197+/-2 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to (C2H5)2O and CH3COOMe	

C4H8O2	CAS REGN : 625-55-8
O++C**O**C**C * * C	
Formic acid, 1-methylethyl ester (9CI) Formic acid, isopropyl ester (8CI)	
GB (exp) : 185.8+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 193.2+/-2 kcal/mol	Temperature : 298 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on (CH3)2CO Entropy change assumed zero	

C4H8O2S	CAS REGN : 38103-96-7
C**O**C**S**C**C + + O	
Carbonothioic acid, S-ethyl O-methyl ester (9CI)	
GB (exp) : 190.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 198.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H8O3	CAS REGN : 623-53-0
C**O**C**O**C**C + + O	
Carbonic acid, ethyl methyl ester (8CI9CI)	
GB (exp) : 192.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 199.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H9N	CAS REGN : 123-75-1
<pre> C*****N * * * * C C * * * * * * C </pre>	
Pyrrolidine (8CI9CI)	
GB (exp) : 213.70+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 221.90+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 213.30+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 221.50+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C4H9N	CAS REGN : 1190-79-0
C**C++N**C**C	
Ethanamine, N-ethylidene- (9CI) Ethylamine, N-ethylidene- (8CI)	
PA (exp) : 220.5+/-2 kcal/mol	Temperature :
Aue, D. H., Webb, H. M., and Bowers, M. T., Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Nitrogen and Oxygen Bases. Hybridization Effects, J. Am. Chem. Soc. 97, 4137 (1975).	
Method used : ICR Equilibrium; scale based on CH3NH2	

C4H9NO	CAS REGN : 110-91-8
<pre> O**C * * * * * * C C * * * * * * C**N </pre>	
Morpholine (8CI9CI)	
GB (exp) : 208.6+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.8+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C4H9NO	CAS REGN : 127-19-5
C**C**N**C + * + * O C	
Acetamide, N,N-dimethyl- (8CI9CI)	
GB (exp) : 205.4+/-2.5 kcal/mol	Temperature : 300 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	

C4H9NO	CAS REGN : 563-83-7
C**C**C**N * % * % C 0	
Propanamide, 2-methyl- (9CI) Isobutyramide (8CI)	
GB (exp) : 206.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 213.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H10	CAS REGN : 106-97-8
C**C**C**C	
Butane (8CI9CI)	
PA (exp) : 166.8 (2) kcal/mol	Temperature : 291 K
Hiraoka, K., and Kebarle, P., Stabilities and Energetics of Pentacoordinated Carbonium Ions. The Isomeric C2H7+ Ions and Some Higher Analogues: C3H9+ and C4H11+, J. Am. Chem. Soc. 98, 6119 (1976).	
Method used : MS Van't Hoff plot C-H protonation (2) Equilibrium of reaction t-C4H9+ + H2 -> C4H11+ (2)	
PA (exp) : 164 (1) kcal/mol	Temperature : 291 K
Hiraoka, K., and Kebarle, P., Stabilities and Energetics of Pentacoordinated Carbonium Ions. The Isomeric C2H7+ Ions and Some Higher Analogues: C3H9+ and C4H11+, J. Am. Chem. Soc. 98, 6119 (1976).	
Method used : MS Van't Hoff plot C-C protonation (1) Equilibrium of reaction C3H7+ + CH4 -> C4H11+ (1)	

C4H10N2	CAS REGN : 110-85-0
<pre> N**C * * * * C C * * * * C**N </pre>	
Piperazine (8CI9CI)	
GB (exp) : 210.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 218.9+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C4H100	CAS REGN : 60-29-7
C**C**O**C**C	
Ethane, 1,1'-oxybis- (9CI) Ethyl ether (8CI)	
PA (exp) : 198+/-4 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to (CH3)2CO and NH3.	
GB (exp) : 188.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 196.1+/-2 kcal/mol	Temperature : 300 K
Harrison, A. G., Lin, P. H., and Tsang, C. W., Proton Transfer Reactions by Trapped-Ion Mass Spectrometry, Int. J. Mass Spectrom. Ion Phys. 19, 23 (1976).	
Method used : MS Equilibrium; scale based on (CH3)2CO Entropy change assumed zero	

Continued on next page

GB (exp) : 182.5+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 197.3+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 190+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taaqepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H100	CAS REGN : 71-36-3
O**C**C**C**C	
1-Butanol (9CI) Butyl alcohol (8CI)	
GB (exp) : 181.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H100	CAS REGN : 75-65-0
O * * C**C**C * * C	
2-Propanol, 2-methyl- (9CI) tert-Butyl alcohol (8CI)	
PA (exp) : 194+/-1 kcal/mol	Temperature :
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH2CO and I-C5H10	
PA (exp) : 192 kcal/mol	Temperature :
Hiraoka, K., and Kebarle, P. Condensation Reactions Involving Carbonium Ions and Lewis Bases in the Gas Phase. Hydration of the tert-Butyl Cation, J. Am. Chem. Soc. 99, 360 (1977).	
Method used : MS Equilibrium relative to known standard acids	

C4H10S	CAS REGN : 75-66-1
S * * C**C**C * * C	
2-Propanethiol, 2-methyl- (8CI9CI)	
GB (exp) : 186.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 194.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H10S	CAS REGN : 352-93-2
C**C**S**C**C	
Ethane, 1,1'-thiobis- (9CI) Ethyl sulfide (8CI)	
GB (exp) : 195.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 202.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C4H11N	CAS REGN : 75-64-9
<pre> N * * C**C**C * * C </pre>	
2-Propanamine, 2-methyl- (9CI) tert-Butylamine (8CI)	
GB (exp) : 210.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 218.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 210.2+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 218.6+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on CH3NH2	

C4H11N	CAS REGN : 78-81-9
C**C**C**N * * C	
1-Propanamine, 2-methyl- (9CI) Isobutylamine (8CI)	
GB (exp) : 207.7+/-2 kcal/mol	Temperature : 298 K
Taft, R. W., Proton-Transfer Reactions, Ed, E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	
GB (exp) : 208.3+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.7+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C4H11N	CAS REGN : 109-73-9
N**C**C**C**C	
1-Butanamine (9CI) Butylamine (8CI)	
GB (exp) : 207.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 216+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 207.8+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.2+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C4H11N	CAS REGN : 109-89-7
C**C**N**C**C	
Ethanamine, N-ethyl- (9CI) Diethylamine (8CI)	
GB (exp) : 214.50+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 222.70+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 214.10+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 222.30+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C4H11N	CAS REGN : 598-56-1
C**N**C**C * * C	
Ethanamine, N,N-dimethyl- (9CI) Ethylamine, N,N-dimethyl- (8CI)	
GB (exp) : 216.70+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 224.50+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 216.00+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 223.80+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C4H11N	CAS REGN : 13952-84-6
C**C**C**C * * N	
2-Butanamine (9CI) sec-Butylamine (8CI)	
GB (exp) : 208.9 kcal/mol	Temperature : 300 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	
GB (exp) : 209.3+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 217.7+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C4H12N2	CAS REGN : 110-60-1
N**C**C**C**C**N	
1,4-Butanediamine (8CI9CI)	
GB (exp) : 221.70+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 236.60+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C5Fe05	CAS REGN : 13463-40-6
<pre> O # # C C##O * * ** O##C**FE*C##O * * C # # O </pre>	
Iron carbonyl (Fe(CO)5), (TB-5-11)- (9CI) Iron carbonyl (Fe(CO)5) (8CI)	
PA (exp) : 200+/-3 kcal/mol	Temperature : 300 K
Foster, M. S., and Beauchamp, J. L., Gas-Phase Ion Chemistry of Iron Pentacarbonyl by Ion Cyclotron Resonance Spectroscopy. New Insights into the Properties and Reactions of Transition Metal Complexes in the Absence of Complicating Solvation Phenomena, J. Am. Chem. Soc. 97, 4808 (1975).	
Method used : ICR Bracketing relative to (CH3O)2CO and NH3.	


C5H3ClN4	CAS REGN : 87-42-3
<pre> C . . . N N . . CLxC C . . . % C % * % * % N% % % % C </pre>	
1H-Purine, 6-chloro- (9CI) Purine, 6-chloro- (8CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to NH3 and CH3NH2.	

C5H4ClN	CAS REGN : 109-09-1
	
Pyridine, 2-chloro- (8CI9CI)	
GB (exp) : 204.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 211.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H4ClN	CAS REGN : 626-60-8
	
Pyridine, 3-chloro- (8CI9CI)	
GB (exp) : 204.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 212.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H4ClN	CAS REGN : 626-61-9
	
Pyridine, 4-chloro- (8CI9CI)	
GB (exp) : 207.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 215+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H4FN	CAS REGN : 372-47-4
	
Pyridine, 3-fluoro- (8CI9CI)	
GB (exp) : 203.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 211.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C5H4FN	CAS REGN : 372-48-5
	
Pyridine, 2-fluoro- (8CI9CI)	
GB (exp) : 200.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 208.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H4FN	CAS REGN : 694-52-0
<pre> C . . . C N . . F**C C . . C </pre>	
Pyridine, 4-fluoro- (8CI9CI)	
GB (exp) : 206.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 214.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H4N2O2	CAS REGN : 1122-61-8
<pre> C . . . C N . . O++N**C C + . + . O C </pre>	
Pyridine, 4-nitro- (8CI9CI)	
GB (exp) : 198.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 206.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 201+/-2 kcal/mol	Temperature : 300 K
Taagepera, M., Henderson, W. G., Brownlee, R. T. C., Beauchamp, J. L., Holtz, D., and Taft, R. W., Gas-Phase Basicities and Pyridine Substituent Effects. J. Am. Chem. Soc. 94, 1369 (1972).	
Method used : ICR Equilibrium; scale based on Pyridine	

C5H4N4O

CAS REGN : 68-94-0




6H-Purin-6-one, 1,7-dihydro- (9CI)
Hypoxanthine (8CI)

PA (exp) : 215+/-4 kcal/mol

Temperature :

Wilson, M. S., and McCloskey, J. A.,
Chemical Ionization Mass Spectrometry of Nucleosides.
Mechanisms of Ion Formation and Estimations of
Proton Affinity,
J. Am. Chem. Soc. 97, 3436 (1975).

Method used : MS
Bracketing relative to CH3NH2 and (CH3)2NH.

C5H5N	CAS REGN : 110-86-1
	
Pyridine (8CI9CI)	
GB (exp) : 210.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 218.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : >209.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to (CH3)2NH	

Continued on next page

GB (exp) : 209.8+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 217.6+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH₃NH₂	
GB (exp) : 203.7+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 220.9+/-2 kcal/mol	Temperature : 600 K
Briggs, J. P., Yamdagni, R., and Kebarle, P., Intrinsic Basicities of Ammonia, Methylamines, Anilines, and Pyridine from Gas-Phase Proton-Exchange Equilibria, J. Am. Chem. Soc. 94, 5128 (1972).	
Method used : MS Equilibrium; scale based on NH₃ Entropy change assumed zero	

C5H5N5	CAS REGN : 73-24-5
<pre> C . N . N . N% % C C % . . % C N * % * % N% % % % C </pre>	
1H-Purin-6-amine (9CI) Adenine (8CI)	
PA (exp) : 220+/-2 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)2NH and (CH3)3N	

C5H5N5O	CAS REGN : 73-40-5
<pre> C%%N % % % % % C..N % . % . N**C C%%N . . . C..N . % . % . 0 </pre>	
6H-Purin-6-one, 2-amino-1,7-dihydro- (9CI) Guanine (8CI)	
PA (exp) : 220+/-2 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)2NH and (CH3)3N.	

C5H6N2O2	CAS REGN : 65-71-4
<pre> C..N . . C**C . . C%%O . . C..N % % O </pre>	
2,4(1H,3H)-Pyrimidinedione, 5-methyl- (9CI) Thymine (8CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to NH3 and CH3NH2	

C5H7F3O2	CAS REGN : 383-66-4
F * * F**C**C**O**C**C**C * + * + F 0	
Acetic acid, trifluoro-, propyl ester (8CI9CI)	
GB (exp) : 176.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 184.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H8	CAS REGN : 693-86-7
<pre> C * * * * C*****C * * C++C </pre>	
Cyclopropane, ethenyl- (9CI) Cyclopropane, vinyl- (8CI)	
GB (exp) : 187.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 195.2+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H8O	CAS REGN : 765-43-5
<pre> C * * * * * * * * * * * C*****C * * C**C + + 0 </pre>	
Ethanone, 1-cyclopropyl- (9CI) Ketone, cyclopropyl methyl (8CI)	
GB (exp) : 194.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 202.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H8O2	CAS REGN : 123-54-6
C**C**C**C**C + + + + 0 0	
2,4-Pentanedione (8CI9CI)	
GB (exp) : 197.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 204.6+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia From Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H8O2	CAS REGN : 2868-37-3
<pre> C * * * * * * C*****C * * C**O**C + + 0 </pre>	
Cyclopropanecarboxylic acid, methyl ester (8CI9CI)	
GB (exp) : 192.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H9N	CAS REGN : 110-59-8
C**C**C**C**C##N	
Pentanenitrile (9CI) Valeronitrile (8CI)	
PA (exp) : 191.9+/-2 kcal/mol	Temperature : 300 K
Staley, R. H., Kleckner, J. E., and Beauchamp, J. L. Relationship between Orbital Ionization Energies and Molecular Properties. Proton Affinities and Photoelectron Spectra of Nitriles, J. Am. Chem. Soc. 98, 2081 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to HC02C2H5	
GB (exp) : 184.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 191.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H10	CAS REGN : 513-35-9
C**C++C**C * * C	
2-Butene, 2-methyl- (8CI9CI)	
PA (exp) : 194.4+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Van't Hoff plot Equilibrium of hydride transfer reaction between (t-C4H9+) + Me2CHEt. This delta H of the reaction was determined from van't Hoff plot. PA (M) derived from the obtained delta Hf(MH+); scale based on delta Hf(t-C4H9+) = 169	

C5H10	CAS REGN : 563-46-2
C**C**C**C + + C	
1-Butene, 2-methyl- (8CI9CI)	
PA (exp) : 193.5+/-2 kcal/mol	Temperature :
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to Iso-C4H8	
PA (exp) : 197.32 kcal/mol	Temperature : 300 K
Solomon, J. J., and Field, F. H., Reversible Reactions of Gaseous Ions. IX. The Stability of C4-C7 Tertiary Alkyl Carbonium Ions, J. Am. Chem. Soc. 97, 2625 (1975).	
Method used : MS Equilibrium scale Based on delta H formation(T-C4H9+) = 169 Kcal/mol.	

Continued on next page

GB not calculated for this reference	
PA (exp) : 195.9+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Equilibrium of hydride transfer reaction between (t-C ₄ H ₉ ⁺) + Me ₂ CH ₂ Et. This delta H of the reaction was determined from van't Hoff plot. PA (M) derived from the obtained delta Hf(MH ⁺); scale based on delta Hf(t-C ₄ H ₉ ⁺) = 169	

C5H10O	CAS REGN : 96-22-0
C**C**C**C**C + + O	
3-Pentanone (8CI9CI)	
GB (exp) : 191.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 198.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

C5H100	CAS REGN : 96-47-9
<pre> O*****C * * * * C**C C * * * * C </pre>	
Furan, tetrahydro-2-methyl- (8CI9CI)	
GB (exp) : 192.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H100	CAS REGN : 110-62-3
C**C**C**C**C++0	
Pentanal (9CI) Valeraldehyde (8CI)	
GB (exp) : 182.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.6+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H100	CAS REGN : 142-68-7
<pre> C**O * * * * * * C * C * * * * * * C**C </pre>	
2H-Pyran, tetrahydro- (8CI9CI)	
GB (exp) : 189.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H10O	CAS REGN : 563-80-4
C**C**C**C * + * + C O	
2-Butanone, 3-methyl- (8CI9CI)	
GB (exp) : 190.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 198.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H10O2	CAS REGN : 109-60-4
C**C**O**C**C**C + + O	
Acetic acid, propyl ester (8CI9CI)	
PA (exp) : 200+/-3 kcal/mol	Temperature :
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to NH3 and CH3COOC2H5.	
GB (exp) : 182.9+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 200.1+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds Between Water and Ammonia and Substituted Benzenes from A Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C5H10O2	CAS REGN : 547-63-7
C**O**C**C**C + * + * O C	
Propanoic acid, 2-methyl-, methyl ester (9CI) Isobutyric acid, methyl ester (8CI)	
GB (exp) : 191.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 198.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H10O2	CAS REGN : 592-84-7
O++C**O**C**C**C**C	
Formic acid, butyl ester (8CI9CI)	
GB (exp) : 184.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 192.5+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H11N	CAS REGN : 110-89-4
<pre> C**N * * * * C * C * * * * * * C**C </pre>	
Piperidine (8CI9CI)	
GB (exp) : 214.90+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 223.10+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 214.40+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 222.60+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

Continued on next page

GB (exp) : 209.60+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 226.80+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C5H11N	CAS REGN : 120-94-5
<pre> C*****C * * * * * * C**N C * * * * * * C </pre>	
Pyrrolidine, 1-methyl- (8CI9CI)	
GB (exp) : 217.80+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 225.60+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 217.30+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 225.10+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C5H11NO2	CAS REGN : 687-48-9
C**N**C**O**C**C * + * + C 0	
Carbamic acid, dimethyl-, ethyl ester (8CI9CI)	
GB (exp) : 203.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 211.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H12O	CAS REGN : 625-54-7
C**C**O**C**C * * C	
Propane, 2-ethoxy- (9CI) Ether, ethyl isopropyl (8CI)	
GB (exp) : 192.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H120	CAS REGN : 1634-04-4
C * * C**C**O**C * * C	
Propane, 2-methoxy-2-methyl- (9CI) Ether, tert-butyl methyl (8CI)	
GB (exp) : 191.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 199.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H13N	CAS REGN : 110-58-7
C**C**C**C**C**N	
1-Pentanamine (9CI) Pentylamine (8CI)	
GB (exp) : 208+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.4+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C5H13N	CAS REGN : 594-39-8
N * * C**C**C**C * * C	
2-Butanamine, 2-methyl- (9CI) tert-Pentylamine (8CI)	
GB (exp) : 211.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 220.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 211.1+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 219.5+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C5H13N	CAS REGN : 616-39-7
<chem>C**C**N**C**C</chem> * * C	
Ethanamine, N-ethyl-N-methyl- (9CI) Diethylamine, N-methyl- (8CI)	
GB (exp) : 218.90+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 226.70+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 218.40+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 226.20+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C5H13N	CAS REGN : 996-35-0
C**C**N**C * * * * C C	
2-Propanamine, N,N-dimethyl- (9CI) Ethylamine, N,N,1-trimethyl- (8CI)	
GB (exp) : 218.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 226.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H13N	CAS REGN : 5813-64-9
C * * C**C**C**N * * C	
1-Propanamine, 2,2-dimethyl- (9CI) Neopentylamine (8CI)	
GB (exp) : 208.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 217.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 209.0+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 217.4+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C5H13N	CAS REGN : 19961-27-4
C**C**N**C**C * * C	
2-Propanamine, N-ethyl- (9CI) Diethylamine, 1-methyl- (8CI)	
GB (exp) : 216+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 224.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C5H14N2	CAS REGN : 109-55-7
C**N**C**C**C**N * * C	
1,3-Propanediamine, N,N-dimethyl- (8CI9CI)	
GB (exp) : 225.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Equilibrium; scale based on NH3	

C5H14N2	CAS REGN : 462-94-2
N**C**C**C**C**C**N	
1,5-Pentanediamine (8CI9CI)	
GB (exp) : 220.00+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 234.60+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	
GB (exp) : 221.40+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 235.10+/-2 kcal/mol	Temperature : 298 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on (CH3)3N Van't Hoff plot	

C6F6	CAS REGN : 392-56-3
<pre> F F * * * * C..C . . F**C C**F . . C..C * * * * F F </pre>	
Benzene, hexafluoro- (8CI9CI)	
GB (exp) : 171.2 kcal/mol	Temperature : 373 K
PA (exp) : 178.5+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

C6HF5	CAS REGN : 363-72-4
<pre> C . . . F**C . C**F . . F**C . C**F C * * F </pre>	
Benzene, pentafluoro- (8CI9CI)	
GB (exp) : 172.7+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 180.3+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

C6H2F4	CAS REGN : 327-54-8
<pre> F * * C..C . . F**C C**F . . C..C * * F </pre>	
Benzene, 1,2,4,5-tetrafluoro- (8CI9CI)	
GB (exp) : 173.1+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 180.3+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

C6H2F4	CAS REGN : 551-62-2
<pre> C . . F**C . C . . F**C . C**F . . C * * F </pre>	
Benzene, 1,2,3,4-tetrafluoro- (8CI9CI)	
GB (exp) : 173.4+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 180.4+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

C6H2F4	CAS REGN : 2367-82-0
<pre> C . . . F**C C**F . . F**C C C * * F </pre>	
Benzene, 1,2,3,5-tetrafluoro- (8CI9CI)	
GB (exp) : 173.4+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 180.5+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	


C6H3F3	CAS REGN : 367-23-7
<pre> C..C . . F**C . C**F . . C..C x x F </pre>	
Benzene, 1,2,4-trifluoro- (8CI9CI)	
GB (exp) : 173.7+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 181.35+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

C6H3F3	CAS REGN : 372-38-3
<pre> F x x C . . . C C . . FxxC CxxF . . C </pre>	
Benzene, 1,3,5-trifluoro- (8CI9CI)	
GB (exp) : 174.1+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 181.15+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

Continued on next page

GB (exp) : 165.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 179.8+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C6H4F2	CAS REGN : 367-11-3
<pre> C..C . . F**C C . . C..C * * F </pre>	
Benzene, 1,2-difluoro- (9CI) Benzene, o-difluoro- (8CI)	
GB (exp) : 174.36+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 181.35+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

C6H4F2	CAS REGN : 372-18-9
	
Benzene, 1,3-difluoro- (9CI) Benzene, m-difluoro- (8CI)	
GB (exp) : 174.42+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 171.67+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

Continued on next page


GB (exp) : 165.9+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 181.6+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	


C6H4F2	CAS REGN : 540-36-3
<pre> C . . . C C**F . . F**C C . . C . </pre>	
Benzene, 1,4-difluoro- (9CI) Benzene, p-difluoro- (8CI)	
GB (exp) : 173.9+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 180.6+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	


C6H4F3N	CAS REGN : 368-48-9
<pre> C . . F N C * . . * . . F**C**C C * . . * . . F C </pre>	
Pyridine, 2-(trifluoromethyl)- (8CI9CI)	
GB (exp) : 201.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 209.0+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C6H4F3N	CAS REGN : 3796-23-4
<pre> N . . . F C . C * . . * . . F**C**C C * . . * . . F . C </pre>	
Pyridine, 3-(trifluoromethyl)- (8CI)	
GB (exp) : 202.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 210.0+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H4F3N	CAS REGN : 3796-24-5
<pre> C . . . F C N * . * . F**C**C * . * . F C </pre>	
Pyridine, 4-(trifluoromethyl)- (8CI9CI)	
GB (exp) : 202.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 210.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 207.1+/-2 kcal/mol	Temperature : 298 K
Taagepera, M., Henderson, W. G., Brownlee, R. T. C., Beauchamp, J. L., Holtz, D., and Taft, R. W., Gas-Phase Basicities and Pyridine Substituent Effects, J. Am. Chem. Soc. 94, 1369 (1972).	
Method used : ICR Equilibrium; scale based on C5H5N(Pyridine)	

C6H4N2	CAS REGN : 100-48-1
	
4-Pyridinecarbonitrile (9CI) Isonicotinonitrile (8CI)	
GB (exp) : 199.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 207.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C6H4N2	CAS REGN : 100-54-9
	
3-Pyridinecarbonitrile (9CI) Nicotinonitrile (8CI)	
GB (exp) : 199.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 206.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H4N2	CAS REGN : 100-70-9
	
2-Pyridinecarbonitrile (9CI) Picolinonitrile (8CI)	
GB (exp) : 197.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 205.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H5Cl	CAS REGN : 108-90-7
	
Benzene, chloro- (8CI9CI)	
GB (exp) : 174.24+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 181.9+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

Continued on next page


GB (exp) : 166.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 182.3+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB and PA values respectively 167.8+/-2 and 181.2+/-2 kcal/mol	
GB (exp) : 166.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 182.5+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C6H5F	CAS REGN : 462-06-6
	
Benzene, fluoro- (8CI9CI)	
GB (exp) : 174.5+/-2 kcal/mol	Temperature : 373 K
PA (exp) : 182.2+/-2 kcal/mol	Temperature : 373 K
Hartman, K. G., and Lias, S. G. Proton Transfer Equilibria in Halobenzene Systems: Entropy Changes and Relative Proton Affinities, Intern. J. Mass Spectrom. Ion Phys. 28, 213 (1978).	
Method used : ICR Calculated entropy Equilibrium; scale based on C6H6	

Continued on next page

GB (exp) : 167+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 182.5+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB and PA values respectively 167.8+/-2 and 181.2+/-2 kcal/mol	
GB (exp) : 166.9+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 182.6+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	

C6H5NO2	CAS REGN : 98-95-3
<pre> C . . . C C . . O++N**C C + . + . O C </pre>	
Benzene, nitro- (8CI9CI)	
GB (exp) : 176.7+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 192.2+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB (C6H6) = 167.8+/-2 and PA (C6H6) = 181.2+/-2	

C6H6	CAS REGN : 71-43-2
	
Benzene (8CI9CI)	
GB (exp) : 172+/-2 kcal/mol	Temperature : 340 K
PA (exp) : 180.1+/-2 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Equilibrium relative to HCOOH Entropy change assumed zero	
GB (exp) : 174.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 181.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 167.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 181.2+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	


C6H6ClN	CAS REGN : 106-47-8
<pre> C . . . C C**N . . CLxC C C </pre>	
Benzenamine, 4-chloro- (9CI) Aniline, p-chloro- (8CI)	
GB (exp) : 198.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 206.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H6ClN	CAS REGN : 108-42-9
<pre> C . . C C . . CLxC C*N . . C C . . </pre>	
Benzenamine, 3-chloro- (9CI) Aniline, m-chloro- (8CI)	
GB (exp) : 197.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 205.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H6FN	CAS REGN : 371-40-4
<pre> C . C . C**N . F**C C . C </pre>	
Benzenamine, 4-fluoro- (9CI) Aniline, p-fluoro- (8CI)	
GB (exp) : 198.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 206.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H6FN	CAS REGN : 372-19-0
<pre> C . . . C C . . F**C C**N . . C </pre>	
Benzenamine, 3-fluoro- (9CI) Aniline, m-fluoro- (8CI)	
GB (exp) : 197.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 205.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H6N4	CAS REGN : 2004-03-7
<pre> C . . . N N . . C**C C . . . % . % C N * % * % N%%%%C </pre>	
1H-Purine, 6-methyl- (9CI) Purine, 6-methyl- (8CI)	
PA (exp) : 220+/-2 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)2NH and (CH3)3N	

C ₆ H ₆ O	CAS REGN : 108-95-2
	
Phenol (8CI9CI)	
PA (exp) : ~180 kcal/mol	Temperature :
DeFrees, D. J. McIver, R. T., and Hehre, W. J. The Proton Affinities of Phenol, J. Am. Chem. Soc. 99, 3853 (1977).	
Method used : ICR Bracketing relative to various molecules. Oxygen protonation	
GB (exp) : 179.1+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 194.6+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to C ₆ H ₆ . Using GB (C ₆ H ₆) = 167.8+/-2 and PA (C ₆ H ₆) = 181.2+/-2	

C6H6O	CAS REGN : 6921-27-3
C##C**C**O**C**C##C	
1-Propyne, 3,3'-oxybis- (9CI) 2-Propynyl ether (8CI)	
GB (exp) : 181.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C6H7N	CAS REGN : 62-53-3
<pre> C . . . C C . . N**C C . . C </pre>	
Benzenamine (9CI) Aniline (8CI)	
GB (exp) : 200.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 208.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 193.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 208.9+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB (C6H6) = 167.8+/-2 and PA (C6H6) = 181.2	


Continued on next page


GB (exp) : >193.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to NH3	
GB (exp) : 194.0+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 211.2+/-2 kcal/mol	Temperature : 600 K
Briggs, J. P., Yamdagni, R., and Kebarle, P., Intrinsic Basicities of Ammonia, Methylamines, Anilines, and Pyridine from Gas-Phase Proton- Exchange Equilibria, J. Am. Chem. Soc. 94, 5128 (1972).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

Continued on next page

GB (exp) : 199.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 203.9+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH ₃ NH ₂	


C6H7N	CAS REGN : 108-89-4
	
Pyridine, 4-methyl- (9CI) 4-Picoline (8CI)	
GB (exp) : 214.30+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 222.10+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H7N	CAS REGN : 108-99-6
	
Pyridine, 3-methyl- (9CI) 3-Picoline (8CI)	
GB (exp) : 213+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 220.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H7N	CAS REGN : 109-06-8
	
Pyridine, 2-methyl- (9CI) 2-Picoline (8CI)	
GB (exp) : 213.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 221.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H7N	CAS REGN : 6921-28-4
C##C**C**N**C**C##C	
2-Propyn-1-amine, N-2-propynyl- (9CI) Di-2-propynylamine (8CI)	
GB (exp) : 205.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 213.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C6H7NO	CAS REGN : 620-08-6
<pre> C . . . C N . . C**O**C C C </pre>	
Pyridine, 4-methoxy- (8CI9CI)	
GB (exp) : 217.00+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 224.80+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H7NO	CAS REGN : 694-85-9
	
2(1H)-Pyridinone, 1-methyl- (9CI) 2(1H)-Pyridone, 1-methyl- (8CI)	
PA (exp) : 217.2+/-2 kcal/mol	Temperature :
Cook, M. J., Katritzky, A. R., Taagepera, M., Singh, I. D., and Taft, R. W. Application of Ion Cyclotron Resonance Spectro- scopic Gas-Phase Basicities to the Study of Tautomeric Equilibria, J. Am. Chem. Soc. 98, 6048 (1976).	
Method used : ICR Equilibrium relative to pyridine. Use value of PA (pyridine) = 210.1 Entropy change assumed zero	

C6H7NO	CAS REGN : 1628-89-3
<pre> C N C . . C**O**C C C </pre>	
Pyridine, 2-methoxy- (8CI9CI)	
GB (exp) : 211.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 219.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page


PA (exp) : 219.8+/-2 kcal/mol	Temperature :
Cock, M. J., Katritzky, A. R., Taagepera, M., Singh, T. D., and Taft, R. W. Application of Ion Cyclotron Resonance Spectro- scopic Gas-Phase Basicities to the Study of Tautomeric Equilibria, J. Am. Chem. Soc. 98, 6048 (1976).	
Method used : ICR Equilibrium relative to t-Butylamine. Use value of PA (t-C4H9NH2) = 218.3 Entropy change assumed zero	

C6H7NO	CAS REGN : 7295-76-3
	
Pyridine, 3-methoxy- (8CI9CI)	
GB (exp) : 213.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 221.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H7N5O	CAS REGN : 23662-75-1
<pre> C * * C%%N % % % % % % C..N % % % % N**C C%%N . C..N % % O </pre>	
6H-Purin-6-one, 2-amino-1,7-dihydro-8-methyl- (9CI) Guanine, 8-methyl- (8CI)	
PA (exp) : >222 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)3N	

C6H8N2	CAS REGN : 95-54-5
<pre> C..C . . N**C . . C . . C..C * * N </pre>	
1,2-Benzenediamine (9CI) o-Phenylenediamine (8CI)	
GB (exp) : >193.7 kcal/mol	Temperature : 424-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to NH3	

C6H8N2	CAS REGN : 106-50-3
<pre> C . . . C C**N . . N**C C C </pre>	
1,4-Benzenediamine (9CI) p-Phenylenediamine (8CI)	
GB (exp) : >193.7 kcal/mol	Temperature : 424-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to NH3	

C6H8N2	CAS REGN : 108-45-2
	
1,3-Benzenediamine (9CI) m-Phenylenediamine (8CI)	
GB (exp) : >202.9 kcal/mol	Temperature : 424-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to CH3NH2	


C6H9F3O2	CAS REGN : 367-64-6
F * * F**C**C**O**C**C**C**C * + * + F 0	
Acetic acid, trifluoro-, butyl ester (8CI9CI)	
GB (exp) : 176.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 184.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H10	CAS REGN : 693-89-0
<pre> C*****C + * + * C**C C * * * * C </pre>	
Cyclopentene, 1-methyl- (8CI9CI)	
GB (exp) : 186.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 195+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H10	CAS REGN : 1528-30-9
<pre> C*****C * * * * C++C C * * * * * C </pre>	
Cyclopentane, methylene- (8CI9CI)	
GB (exp) : 190.7+/-2 kcal/mol	Temperature : 298 K
Pollack, S. K., Wolf, J. F., Levi, B. A., Taft, R. W., and Hehre W. J. Kinetic Detection of Common Intermediates in Gas Phase Ion-Molecule Reactions, J. Am. Chem. Soc. 99, 1350 (1977).	
Method used : ICR Equilibrium; scale based on NH3	

C6H10	CAS REGN : 4663-22-3
<pre> C * * * * C*****C * * C**C + + C </pre>	
Cyclopropane, (1-methylethenyl)- (9CI) Propene, 2-cyclopropyl- (8CI)	
GB (exp) : 198.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 207.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H10F3NO	CAS REGN : 400-59-9
F * * F**C**C**N**C**C**C**C * % * % F 0	
Acetamide, N-butyl-2,2,2-trifluoro- (8CI9CI)	
GB (exp) : 193.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 201.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H100	CAS REGN : 108-94-1
	
Cyclohexanone (3CI9CI)	
PA (exp) : 204 kcal/mol	Temperature :
Jelus, B. L., Murray, R. K., Jr., Munson, B., Studies in Chemical Ionization Mass Spectrometry. Secondary Alcohols with Isobutane, J. Am. Chem. Soc. 97, 2362 (1975).	
Method used : MS The (M-H)+ ion for cyclo-hexanol is assumed to be the same as O-protonated cyclohexanone.	

C6H100	CAS REGN : 557-40-4
C++C**C**O**C**C++C	
1-Propene, 3,3'-oxybis- (9CI) Allyl ether (8CI)	
GB (exp) : 190+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H11N	CAS REGN : 124-02-7
C++C**C**N**C**C++C	
2-Propen-1-amine, N-2-propenyl- (9CI) Diallylamine (8CI)	
GB (exp) : 213.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 221.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H11NO	CAS REGN : 931-20-4
<pre> C**C * * * * C**N * C * * * * C**C + + O </pre>	
2-Piperidinone, 1-methyl- (9CI) 2-Piperidone, 1-methyl- (8CI)	
PA (exp) : 216.8+/-2 kcal/mol	Temperature :
Cook, M. J., Katritzky, A. R., Taagepera, M., and Singh, T. D., and Taft, R. W. Application of Ion Cyclotron Resonance Spectroscopic Gas-Phase Basicities to the Study of Tautomeric Equilibria, J. Am. Chem. Soc. 98, 6048 (1976).	
Method used : ICR Equilibrium relative to i-C3H7NH2. Use value of PA (i-C3H7NH2) = 216.5 Entropy change assumed zero	

C6H11NO	CAS REGN : 53687-79-9
<pre> N + * + * C C * * * * C**O***C C * * * * C </pre>	
Piperidine, 2-methoxy-	
PA (exp) : 226.3+/-2 kcal/mol	Temperature :
Cook, M. J., Katritzky, A. R., Taagepera, M., and Singh, T. D., and Taft, R. W. Application of Ion Cyclotron Resonance Spectroscopic Gas-Phase Basicities to the Study of Tautomeric Equilibria, J. Am. Chem. Soc. 98, 6048 (1976).	
Method used : ICR Equilibrium relative to (i-C3H7)2NH. Use value of PA(i-C3H7)2NH = 266.6 Entropy change assumed zero	

C6H12	CAS REGN : Not available
C * * C * * C**C**C++C	
1-Butene, 2-ethyl-	
PA (exp) : 197.7+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Equilibrium of hydride transfer reaction between (t-C4H9+) + Et2CHMe and between (t-C5H11+) + Et2CHMe. This delta H of the reaction was determined from van't Hoff plot. PA (M) derived from the obtained delta Hf(MH+); scale based on delta Hf(t-C4H9+) = 169	

C6H12	CAS REGN : 616-12-6
C**C++C**C**C * * C	
2-Pentene, 3-methyl-, (E)- (8CI9CI)	
PA (exp) : 196.1+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Equilibrium of hydride transfer reaction between (t-C4H9+) + Et2CHMe and between (t-C5H11+) + Et2CHMe. This delta H of the reaction was determined from van't Hoff plot. PA (M) derived from the obtained delta Hf(MH+); scale based on delta Hf(t-C4H9+) = 169	

C6H12	CAS REGN : 625-27-4
C**C**C++C**C * * C	
2-Pentene, 2-methyl- (8CI9CI)	
PA (exp) : 196.4+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Equilibrium of hydride transfer Reaction between (t-C4H9+) + Me2CHC3H7 and between (t-C5H11+) + Me2CHC3H7. This delta H of the reaction was determined from van't Hoff plot. PA (M) derived from the obtained delta Hf(MH+); scale based on delta Hf(t-C4H9+) = 169	

C6H12	CAS REGN : 763-29-1
C**C**C**C**C + + C	
i-Pentene, 2-methyl- (8CI9CI)	
PA (exp) : 198.2+/-2 kcal/mol	Temperature : 363-553 K
Goren, A., and Munson, B. Thermochemistry of Alkyl Ions, J. Phys. Chem. 80, 2848 (1976).	
Method used : MS Equilibrium of hydride transfer reaction between (t-C4H9+) + Me2CHC3H7 and between (t-C5H11+) + Me2CHC3H7. This delta H of the reaction was determined from van't Hoff plot. PA (M) derived from the obtained delta Hf(MH+); scale based on delta Hf(t-C4H9+) = 169	

C6H12N2	CAS REGN : 280-57-9
<pre> C**N * ** * C * C * C * C * * * * N**C </pre>	
1,4-Diazabicyclo[2.2.2]octane (8CI9CI)	
GB (exp) : 217.20+/-2 kcal/mol	Temperature :
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	
GB (exp) : 217.90+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 225.90+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

Continued on next page

PA (exp) : 225.50+/-2 kcal/mol	Temperature :
Staley, R., and Beauchamp, J. L., Relationship of Nitrogen Lone Pair Interactions to Thermodynamic Parameters Associated with Amine Basicities, J. Am. Chem. Soc. 96, 1604 (1974).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C6H12O	CAS REGN : 75-97-8
<pre> C * * C**C**C**C * + * + C O </pre>	
2-Butanone, 3,3-dimethyl- (8CI9CI)	
GB (exp) : 191.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 199.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H12O2	CAS REGN : 598-98-1
<pre> C * * * C**O**C**C**C + * + * O C </pre>	
Propanoic acid, 2,2-dimethyl-, methyl ester (9CI) Pivalic acid, methyl ester (8CI)	
GB (exp) : 192.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H13N	CAS REGN : 108-91-8
<pre> C * * * * * * C C * * * * N**C C * * * * C C </pre>	
Cyclohexanamine (9CI) Cyclohexylamine (8CI)	
GB (exp) : 210.10+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 218.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 210.1+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 218.5+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Dowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

Continued on next page

GB (exp) : 204.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 222+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	

C6H13N	CAS REGN : 626-67-5
	
Piperidine, 1-methyl- (8CI9CI)	
GB (exp) : 218.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 226.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 218.3+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 226.1+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C6H14O	CAS REGN : 108-20-3
C**C**O**C**C * * * * C C	
Propane, 2,2'-oxybis- (9CI) Isopropyl ether (8CI)	
GB (exp) : 195.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 203+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H14O	CAS REGN : 111-43-3
C**C**C**O**C**C**C	
Propane, 1,1'-oxybis- (9CI) Propyl ether (8CI)	
GB (exp) : 191.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 199.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H14O	CAS REGN : 637-92-3
<pre> C * * C**C**O**C**C * * C </pre>	
Propane, 2-ethoxy-2-methyl- (9CI) Ether, tert-butyl ethyl (8CI)	
GB (exp) : 195.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 203.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H14S	CAS REGN : 111-47-7
C**C**C**S**C**C**C	
Propane, 1,1'-thiobis- (9CI) Propyl sulfide (8CI)	
GB (exp) : 196.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 204+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H14S	CAS REGN : 625-80-9
C**C**S**C**C * * * * C C	
Propane, 2,2'-thiobis- (9CI) Isopropyl sulfide (8CI)	
GB (exp) : 199.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 206.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C6H15N	CAS REGN : 108-18-9
C**C**N**C**C * * * * C C	
2-Propanamine, N-(1-methylethyl)- (9CI) Diisopropylamine (8CI)	
GB (exp) : 218.40+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 226.60+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 217.90+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 226.1+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C6H15N	CAS REGN : 111-26-2
N**C**C**C**C**C	
1-Hexanamine (9CI) Hexylamine (8CI)	
GB (exp) : 208.1+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.5+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C6H15N	CAS REGN : 121-44-8
C**C**N**C**C * * C * * C	
Ethanamine, N,N-diethyl- (9CI) Triethylamine (8CI)	
GB (exp) : 221.20+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 229.00+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 220.60+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 228.40+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C6H15N	CAS REGN : 142-84-7
C**C**C**N**C**C**C	
1-Propanamine, N-propyl- (9CI) Dipropylamine (8CI)	
GB (exp) : 216.50+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 224.70+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 216.10+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 224.30+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C6H15N	CAS REGN : 918-02-5
<pre> C * * C**C**N**C * * * * C C </pre>	
2-Propanamine, N,N,2-trimethyl- (9CI) Ethylamine, N,N,1,1-tetramethyl- (8CI)	
GB (exp) : 220.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 228.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C6H16N2	CAS REGN : 110-18-9
C**N**C**C**N**C * * * * C C	
1,2-Ethanediamine, N,N,N',N'-tetramethyl- (9CI) Ethylenediamine, N,N,N',N'-tetramethyl- (8CI)	
GB (exp) : 224.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 232.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taaqepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 224+/-2 kcal/mol	Temperature : 298 K
Taft, R. W., Proton-Transfer Reactions, Ed. E. F. Caldin and V. Gold, (John Wiley and Sons, New York, 1975) Ch. 2.	
Method used : ICR Equilibrium; scale based on NH3	


C6H16N2	CAS REGN : 124-09-4
N**C**C**C**C**C**C**N	
1,6-Hexanediamine (8CI9CI)	
GB (exp) : 220.00+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 235.40+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	
GB (exp) : 221.20+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 235.00+/-2 kcal/mol	Temperature : 298 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on (CH3)3N Van't Hoff plot	

C6H17NSi	CAS REGN : 18182-40-6
<pre> C * * C**N**C**SI*C * * * * C C </pre>	
Methanamine, N,N-dimethyl-1-(trimethylsilyl)- (9CI) Trimethylamine, (trimethylsilyl)- (8CI)	
GB (exp) : 219.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 227.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C7H5ClO	CAS REGN : 104-88-1
<pre> C . . . C C**CL O++C**C C . . . C </pre>	
Benzaldehyde, 4-chloro- (9CI) Benzaldehyde, p-chloro- (8CI)	
GB (exp) : 189.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C7H5D3	CAS REGN : 1124-18-1
<pre> C D . * . * . * C C**C**D . * . * . * C C D . . C </pre>	
Benzene, methyl-d3-	
GB (exp) : 180.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188.7+/-2 kcal/mol	Temperature : 300 K
Ausloos, P., and Lias, S. G. Equilibrium Isotope Effects on the Proton Transfer Reactions of Methylbenzenes, J. Am. Chem. Soc. 99, 4198 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C7H5FO	CAS REGN : 456-48-4
	
Benzaldehyde, 3-fluoro- (9CI) Benzaldehyde, m-fluoro- (8CI)	
GB (exp) : 186.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 194.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C7H5FO	CAS REGN : 459-57-4
	
Benzaldehyde, 4-fluoro- (9CI) Benzaldehyde, p-fluoro- (8CI)	
GB (exp) : 183.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 196.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H5N	CAS REGN : 100-47-0
<pre> C . . . C C . . N##C**C C . . C </pre>	
Benzonitrile (8CI9CI)	
GB (exp) : 179.2+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 194.7+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB (C6H6) = 167.8+/-2 and PA (C6H6) = 181.2+/-2	

C7H6O	CAS REGN : 100-52-7
	
Benzaldehyde (8CI9CI)	
GB (exp) : 190.4 kcal/mol	Temperature : 300 K
PA (exp) : 198.2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 183.2+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 198.7+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB (C6H6) = 167.8+/-2 and PA (C6H6) = 181.2+/-2	

C7H7NO	CAS REGN : 350-03-8
	
Ethanone, 1-(3-pyridinyl)- (9CI) Ketone, methyl 3-pyridyl (8CI)	
GB (exp) : 206.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 214.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H7NO	CAS REGN : 1122-54-9
	
Ethanone, 1-(4-pyridinyl)- (9CI) Ketone, methyl 4-pyridyl (8CI)	
GB (exp) : 206.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 214.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H8	CAS REGN : 108-88-3
	
Benzene, methyl- (9CI) Toluene (8CI)	
GB (exp) : 179.6+/-2 kcal/mol	Temperature : 340 K
PA (exp) : 187.5+/-2 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Equilibrium relative to CH3SH Entropy change assumed zero	
PA (exp) : 187.4+/-2 kcal/mol	Temperature : 300 K
Devlin, J. L., III., Wolf, J. F., Taft, R. W., Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Equilibrium relative to C6H6(Benzene) Protonation position 2	

Continued on next page


PA (exp) : 183.8+/-2 kcal/mol	Temperature : 300 K
Devlin, J. L., III., Wolf, J. F., Taft, R. W., Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Equilibrium relative to C6H6(Benzene) Protonation position 3	
PA (exp) : 182.6+/-2 kcal/mol	Temperature : 300 K
Devlin, J. L., III., Wolf, J. F., Taft, R. W., Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Equilibrium relative to C6H6(Benzene) Protonation position 1	
GB (exp) : 174.1+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 139.6+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB and PA values respectively 167.8+/-2 and 181.2+/-2 kcal/mol	


Continued on next page


GB (exp) : 180.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188.6+/-2 kcal/mol	Temperature : 300 K
Devlin, J. L., III., Wolf, J. F., Taft, R. W., Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Protonation position 4 Calculated entropy Equilibrium relative to C6H6(Benzene)	
GB (exp) : 180.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 188.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

Continued on next page

GB (exp) : 180.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 187.9+/-2 kcal/mol	Temperature : 300 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on C6H6(Benzene)	

C7H8O	CAS REGN : 100-66-3
	
Benzene, methoxy- (9CI) Anisole (8CI)	
GB (exp) : 183.5+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 199+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy . Equilibrium relative to C6H6. Using GB (C6H6) = 167.8+/-2 and PA (C6H6) = 181.2+/-2	

C7H9N	CAS REGN : 100-46-9
	
Benzenemethanamine (9CI) Benzylamine (8CI)	
GB (exp) : 206.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 214.5+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C7H9N	CAS REGN : 100-61-8
	
Benzenamine, N-methyl- (9CI) Aniline, N-methyl- (8CI)	
GB (exp) : 200.2+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 217.4+/-2 kcal/mol	Temperature : 600 K
Briggs, J. P., Yamdagni, R., and Kebarle, P., Intrinsic Basicities of Ammonia, Methylamines, Anilines, and Pyridine from Gas-Phase Proton- Exchange Equilibria, J. Am. Chem. Soc. 94, 5128 (1972).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	


C7H9N	CAS REGN : 106-49-0
<pre> C . . . C C**N . . C**C C . . . C . . C </pre>	
Benzenamine, 4-methyl- (9CI) p-Toluidine (8CI)	
GB (exp) : 203.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 211.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H9N	CAS REGN : 108-44-1
<pre> C . . . C C . . C**C C**N . . C </pre>	
Benzenamine, 3-methyl- (9CI) m-Toluidine (8CI)	
GB (exp) : 203.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 211.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 211.6+/-2 kcal/mol	Temperature : 298 K
Pollack, S. K., Devlin, J. L., III, Summerhays, K. D., Taft, R. W., and Hehre, W J. The Site of Protonation in Aniline, J. Am. Chem. Soc. 99, 4583 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3. Ring protonation	

Continued on next page

PA (exp) : 211.6+/-2 kcal/mol	Temperature : 300 K
Summerhays, K. D., Pollack, S. K., Taft, R. W., and Hehre, W. J. Gas-Phase Basicities of Substituted Anilines. Inferences about the Role of Solvent in Dictating Site of Protonation, J. Am. Chem. Soc. 99, 4585 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale base on NH3. Ring protonation	

C7H9N	CAS REGN : 108-47-4
	
Pyridine, 2,4-dimethyl- (9CI) 2,4-Lutidine (8CI)	
PA (exp) : 225.1+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., Bowers, M. T., Liotta, C. L., Alexander, C. J., and Hopkins, H. P., Jr. A Quantitative Comparison of Gas- and Solution-Phase Basicities of Substituted Pyridines, J. Am. Chem. Soc. 98, 854 (1976).	
Method used : ICR Calculated Entropy Equilibrium; scale based on NH3	


C7H9N	CAS REGN : 108-48-5
	
Pyridine, 2,6-dimethyl- (9CI) 2,6-Lutidine (8CI)	
PA (exp) : 226.1+/-2 kcal/mol	Temperature : 298 K
Ave, D. H., Webb, H. M., Bowers, M. T., Liotta, C. L., Alexander, C. J., and Hopkins, H. P., Jr. A Quantitative Comparison of Gas- and Solution-Phase Basicities of Substituted Pyridines, J. Am. Chem. Soc. 98, 854 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H9N	CAS REGN : 589-93-5
Pyridine, 2,5-dimethyl- (9CI) 2,5-Lutidine (8CI)	
PA (exp) : 224.5+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., Bowers, M. T., Liotta, C. L., Alexander, C. J., and Hopkins, H. P., Jr. A Quantitative Comparison of Gas- and Solution-Phase Basicities of Substituted Pyridines, J. Am. Chem. Soc. 98, 854 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H9N	CAS REGN : 591-22-0
<pre> N C C . . C**C C**C C </pre>	
Pyridine, 3,5-dimethyl- (9CI) 3,5-Lutidine (8CI)	
PA (exp) : 223.5 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., Bowers, M. T., Liotta, C. L., Alexander, C. J., and Hopkins, H. P., Jr. A Quantitative Comparison of Gas- and Solution-Phase Basicities of Substituted Pyridines, J. Am. Chem. Soc. 98, 854 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H9NO

CAS REGN : 90-04-0



Benzenamine, 2-methoxy- (9CI)
o-Anisidine (8CI)

GB (exp) : 200.2+/-2 kcal/mol

Temperature : 600 K

PA (exp) : 217.4+/-2 kcal/mol

Temperature : 600 K

Yamdagni, R., and Kebarle, P.,
Gas-Phase Basicities of Amines. Hydrogen Bonding
in Proton-Bound Amine Dimers and Proton-Induced
Cyclization of alpha,omega-Diamines,
J. Am. Chem. Soc. 95, 3504 (1973).

Method used : MS
Entropy change assumed zero
Equilibrium; scale based on CH3NH2

C7H9NO	CAS REGN : 104-94-9
<pre> C . . . C C**N . . C**O**C C . . C </pre>	
Benzenamine, 4-methoxy- (9CI) p-Anisidine (8CI)	
GB (exp) : 203.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 212.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H9NO	CAS REGN : 536-90-3
<pre> C . . . C C . . C**O**C C**N . . C </pre>	
Benzenamine, 3-methoxy- (9CI) m-Anisidine (8CI)	
PA (exp) : 214.0+/-2 kcal/mol	Temperature : 300 K
Summerhays, K. D., Pollack, S. K., Taft, R. W., and Hehre, W. J. Gas-Phase Basicities of Substituted Anilines. Inferences about the Role of Solvent in Dictating Site of Protonation, J. Am. Chem. Soc. 99, 4585 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale base on NH3. Ring protonation	

C7H10	CAS REGN : 498-66-8
<pre> C*****C * * * * C*C+C*C * * * * C </pre>	
Bicyclo[2.2.1]hept-2-ene (9CI) 2-Norbornene (8CI)	
PA (exp) : 198.8+/-2 kcal/mol	Temperature : 300 K
Staley, R. H., Wieting, R. D., and Beauchamp, J. L. Carbenium Ion Stabilities in the Gas Phase and Solution. An Ion Cyclotron Resonance Study of Bromide Transfer Reactions Involving Alkali Ions, Alkyl Carbenium Ions, Acyl Cations, and Cyclic Halonium Ions, J. Am. Chem. Soc. 99, 5964 (1977).	
Method used : ICR Calculated entropy Equilibrium relative to Et20; scale based on NH3	

C7H10N2	CAS REGN : 1122-58-3
<pre> C . . / \ C N C**N**C C * * * * C C </pre>	
4-Pyridinamine, N,N-dimethyl- (9CI) Pyridine, 4-(dimethylamino)- (8CI)	
GB (exp) : 224.90+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 232.70+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	
GB (exp) : >209.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to (CH ₃) ₂ NH	

C7H12	CAS REGN : 591-49-1
<pre> C * * * * C C + * + * C**C C * * * * C </pre>	
Cyclohexene, 1-methyl- (8CI9CI)	
GB (exp) : 188.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 196.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, Mciver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H12	CAS REGN : 765-47-9
<pre> C*****C * * * * C**C C + * + * C * * C </pre>	
Cyclopentene, 1,2-dimethyl- (8CI9CI)	
GB (exp) : 188.4+/-2 kcal/mol	Temperature : 298 K
Pollack, S. K., Wolf, J. F., Levi, B. A., Taft, R. W., and Hehre, W. J. Kinetic Detection of Common Intermediates in Gas Phase Ion-Molecule Reactions, J. Am. Chem. Soc. 99, 1350 (1977).	
Method used : ICR	

C7H13N	CAS REGN : 100-76-5
<pre> C**N * ** * C * C * C * C * * * * C**C </pre>	
1-Azabicyclo[2.2.2]octane (9CI) Quinuclidine (8CI)	
GB (exp) : 220.80+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 228.60+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 220.80+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 228.60+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

Continued on next page

GB (exp) : 209.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to (CH ₃) ₂ NH	

C7H17N	CAS REGN : 111-68-2
C**C**C**C**C**C**N	
1-Heptanamine (9CI) Heptylamine (8CI)	
GB (exp) : 208.2+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 216.5+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Evaluation of Intramolecular Strong Hydrogen Bonding in the Gas Phase, J. Am. Chem. Soc. 95, 2669 (1973).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH ₃ NH ₂	

C7H17N	CAS REGN : 4458-31-5
C**C**N**C**C**C * * C * * C	
1-Propanamine, N,N-diethyl- (9CI) Propylamine, N,N-diethyl- (8CI)	
GB (exp) : 221.3+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 229.1+/-2 kcal/mol	Temperature : 298 K
Ave, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C7H17N	CAS REGN : 10076-31-0
<pre> C * * C**N**C**C**C * * * * C C </pre>	
1-Propanamine, N,N,2,2-tetramethyl- (9CI) Propylamine, N,N,2,2-tetramethyl- (8CI)	
GB (exp) : 218.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 226.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C7H18N2	CAS REGN : 110-95-2
C**N**C**C**C**N**C * * * * C C	
1,3-Propanediamine, N,N,N',N'-tetramethyl- (8CI9CI)	
GB (exp) : 227.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Equilibrium; scale based on NH3	

C8H4D6	CAS REGN : 25493-13-4
<pre> D C * . * . D**C**C C D * . * * . * D C C**C**D . . * . . * C D </pre>	
Benzene, 1,4-(dimethyl-d3)-	
GB (exp) : 182.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.8+/-2 kcal/mol	Temperature : 300 K
Ausloos, P., and Lias, S. G. Equilibrium Isotope Effects on the Proton Transfer Reactions of Methylbenzenes, J. Am. Chem. Soc. 99, 4198 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C8H5F3O	CAS REGN : 455-19-6
<pre> C . . . F C . C**C++O * . . * . . F**C**C C * . . * . . F C </pre>	
Benzaldehyde, 4-(trifluoromethyl)- (9CI) p-Tolualdehyde, .alpha.,.alpha.,.alpha.-trifluoro- (8CI)	
GB (exp) : 181.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H5NO	CAS REGN : 105-07-7
<pre> C . C . C**C++O . N##C**C C . C </pre>	
Benzonitrile, 4-formyl- (9CI) Terephthalaldehydonitrile (8CI)	
GB (exp) : 177.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 185.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H8	CAS REGN : 100-42-5
	
Benzene, ethenyl- (9CI) Styrene (8CI)	
GB (exp) : 191.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 199.3+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H8O	CAS REGN : 98-86-2
<pre> C . . . C C . . C**C**C + . + . O C </pre>	
Ethanone, 1-phenyl- (9CI) Acetophenone (8CI)	
GB (exp) : 195.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 203.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H8O	CAS REGN : 104-87-0
<pre> C . . . C C**C . . O++C**C C C </pre>	
Benzaldehyde, 4-methyl- (9CI) p-Tolualdehyde (8CI)	
GB (exp) : 194+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 201.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H8O2	CAS REGN : 93-58-3
	
Benzoic acid, methyl ester (8CI9CI)	
GB (exp) : 193.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 200.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H8O2	CAS REGN : 123-11-5
<pre> C . . . C C**O**C . . O++C**C C . . C </pre>	
Benzaldehyde, 4-methoxy- (9CI) p-Anisaldehyde (8CI)	
GB (exp) : 203.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 210.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C8H10	CAS REGN : 95-47-6
<pre> C..C . . C**C . C . . C..C * * C </pre>	
Benzene, 1,2-dimethyl- (9CI) o-Xylene (8CI)	
GB (exp) : 183+/-3 kcal/mol	Temperature : 340 K
PA (exp) : 190+/-3 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Equilibrium relative to CH3SH and (CH3)2O Entropy change assumed zero	

Continued on next page

GB (exp) : 183.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 191.1+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia From Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H10

CAS REGN : 100-41-4



Benzene, ethyl- (8CI9CI)

GB (exp) : 181.8+/-2 kcal/mol

Temperature : 300 K

Hehre, W. J., McIver, R. T., Jr., Pople, J. A.,
and Schleyer, P. v. R.,
Alkyl Substituent Effects on the Stability of
Protonated Benzene,
J. Am. Chem. Soc. 96, 7162 (1974).


Method used : ICR
Equilibrium; scale based on C6H5CH3(Toluene)


Continued on next page


GB (exp) : 175.1+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 190.6+/-2 kcal/mol	Temperature : 600 K
Lau, Y. K., and Kebarle, P. Substituent Effects on the Intrinsic Basicity of Benzene: Proton Affinities of Substituted Benzenes, J. Am. Chem. Soc. 98, 7452 (1976).	
Method used : MS Calculated entropy Equilibrium relative to C6H6. Using GB (C6H6) = 167.8+/-2 and PA (C6H6) = 181.2+/-2	
GB (exp) : 180.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.1+/-2 kcal/mol	Temperature : 300 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on C6H6(Benzene)	

Continued on next page

GB (exp) : 181.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.6+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H10	CAS REGN : 106-42-3
	
Benzene, 1,4-dimethyl- (9CI) p-Xylene (8CI)	
GB (exp) : 182+/-3 kcal/mol	Temperature : 340 K
PA (exp) : 189+/-3 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Equilibrium; scale based on CH3SH and (CH3)2O Entropy change assumed zero	
GB (exp) : 182.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 189.8+/-2 kcal/mol	Temperature : 300 K
Devlin, J. L., III, Wolf, J. F., Taft, R. W., and Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to C6H6(Benzene)	

C8H10	CAS REGN : 108-38-3
	
Benzene, 1,3-dimethyl- (9CI) m-Xylene (8CI)	
GB (exp) : 182+/-3 kcal/mol	Temperature : 340 K
PA (exp) : 189.7+/-3 kcal/mol	Temperature : 340 K
Chong, S. L., and Franklin, J. L., Proton Affinities of Benzene, Toluene, and the Xylenes, J. Am. Chem. Soc. 94, 6630 (1972).	
Method used : MS Equilibrium relative to CH3SH and (CH3)2O Entropy change assumed zero	
GB (exp) : 187.2+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 194.6+/-2 kcal/mol	Temperature : 298 K
Devlin, J. L., III, Wolf, J. F., Taft, R. W., and Hohre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to C6H6(Benzene)	

C8H11N	CAS REGN : 103-69-5
 Chemical structure diagram of N-ethylbenzenamine (9CI) and its SMILES representation: C**C**N**C.	
Benzenamine, N-ethyl- (9CI) Aniline, N-ethyl- (8CI)	
GB (exp) : 203.6+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 220.8+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : ICR Equilibrium; scale based on NH3 Entropy change assumed zero	

C8H11N	CAS REGN : 121-69-7
<pre> C C C . . . C**N**C C * * . * * . C C </pre>	
Benzenamine, N,N-dimethyl- (9CI) Aniline, N,N-dimethyl- (8CI)	
GB (exp) : 212.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 220.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
PA (exp) : 220+/-2 kcal/mol	Temperature : 298 K
Pollack, S. K., Devlin, J. L., III, Summerhays, K. D., Taft, R. W., and Hehre, W. J. The Site of Protonation in Aniline, J. Am. Chem. Soc. 99, 4583 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3. Protonation on the nitrogen	

Continued on next page

GB (exp) : 206.8+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 224+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities of Amines. Hydrogen Bonding in Proton-Bound Amine Dimers and Proton-Induced Cyclization of alpha,omega-Diamines, J. Am. Chem. Soc. 95, 3504 (1973).	
Method used : MS Equilibrium; scale based on NH3 Entropy change assumed zero	
GB (exp) : >209.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to (CH3)2NH	

C8H18S	CAS REGN : 107-47-1
<pre> C C * * * * C**C**S**C**C * * * * C C </pre>	
Propane, 2,2'-thiobis[2-methyl- (9CI) tert-Butyl sulfide (8CI)	
GB (exp) : 203.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 210.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H19N	CAS REGN : 110-96-3
C**C**C**N**C**C**C * * * * C C	
1-Propanamine, 2-methyl-N-(2-methylpropyl)- (9CI) Diisobutylamine (8CI)	
GB (exp) : 217.7+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 225.9+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	


C8H19N	CAS REGN : 111-92-2
C**C**C**C**N**C**C**C**C	
1-Butanamine, N-butyl- (9CI) Dibutylamine (8CI)	
GB (exp) : 217.20+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 225.40+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C8H19N	CAS REGN : 626-23-3
C**C**C**N**C**C**C * * * * C C	
2-Butanamine, N-(1-methylpropyl)- (9CI) Di-sec-butylamine (8CI)	
GB (exp) : 219.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 228.1+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C8H19N	CAS REGN : 7087-68-5
C**C**N**C**C * * * * * * C C C * * C	
2-Propanamine, N-ethyl-N-(1-methylethyl)- (9CI) Triethylamine, 1,1'-dimethyl- (8CI)	
GB (exp) : 223.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 231.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H19N	CAS REGN : 21981-37-3
<pre> C C * * * * C**C**N**C**C * * * * C C </pre>	
2-Propanamine, N-(1,1-dimethylethyl)-2-methyl- (9CI) Di-tart-butylamine (8CI)	
GB (exp) : 221.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 229.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C8H20N2	CAS REGN : 111-51-3
C**N**C**C**C**C**N**C * * * * C C	
1,4-Butanediamine, N,N,N',N'-tetramethyl- (8CI9CI)	
GB (exp) : 229.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Equilibrium; scale based on NH3	

C9H9Cl	CAS REGN : 1712-70-5
	
Benzene, 1-chloro-4-(1-methylethenyl)- (9CI) Styrene, p-chloro-.alpha.-methyl- (8CI)	
GB (exp) : 195.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 203+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C9H9F	CAS REGN : 350-40-3
<pre> C . . . C C**F . . C**C**C C + . + . C C </pre>	
Benzene, 1-fluoro-4-(1-methylethenyl)- (9CI) Styrene, p-fluoro-.alpha.-methyl- (8CI)	
GB (exp) : 196.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 204.7+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C9H9N	CAS REGN : 6921-29-5
C##C**C**N**C**C##C * * C * * C # # C	
2-Propyn-1-amine, N,N-di-2-propynyl- (9CI) Tri-2-propynylamine (8CI)	
GB (exp) : 209.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 217.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C9H10	CAS REGN : 98-83-9
<pre> C . . . C C . . C**C**C + . + . C C </pre>	
Benzene, (1-methylethenyl)- (9CI) Styrene, .alpha.-methyl- (8CI)	
GB (exp) : 197.0+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 205.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C9H12	CAS REGN : 98-82-8
<pre> C . . . C C . . C**C**C * . * . C C </pre>	
Benzene, (1-methylethyl)- (9CI) Cumene (8CI)	
GB (exp) : 183+/-2 kcal/mol	Temperature : 300 K
Hehre, W. J., McIver, R. T., Jr., Pople, J. A., and Schleyer, P. v. R., Alkyl Substituent Effects on the Stability of Protonated Benzene, J. Am. Chem. Soc. 96, 7162 (1974).	
Method used : ICR Equilibrium; scale based on C6H5CH3(Toluene)	

Continued on next page

GB (exp) : 182.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.4+/-2 kcal/mol	Temperature : 300 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on C6H6(Benzene)	


C9H12	CAS REGN : 103-65-1
	
Benzene, propyl- (8CI9CI)	
GB (exp) : 182.6+/-2 kcal/mol	Temperature : 300 K
Hehre, W. J., McIver, R. T., Jr., Pople, J. A., and Schleyer, P. v. R., Alkyl Substituent Effects on the Stability of Protonated Benzene, J. Am. Chem. Soc. 96, 7162 (1974).	
Method used : ICR Equilibrium; scale based on C6H5CH3(Toluene)	

Continued on next page

GB (exp) : 181.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190+/-2 kcal/mol	Temperature : 300 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on C6H6(Benzene)	
GB (exp) : 182.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C9H12	CAS REGN : 108-67-8
<pre> C * * C . . . C C . . C**C C**C C </pre>	
Benzene, 1,3,5-trimethyl- (9CI) Mesitylene (8CI)	
GB (exp) : 191.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 198.4+/-2 kcal/mol	Temperature : 300 K
Devlin, J. L., III, Wolf, J. F., Taft, R. W., and Hehre, W. J., The Proton Affinities of Toluene, J. Am. Chem. Soc. 98, 1990 (1976).	
Method used : ICR Calculated entropy Equilibrium relative to benzene	

C9H12N2O6	CAS REGN : 58-96-8
<pre> O * * C * * C++C O**C * * * * * * * * 0%%C N*****C * * % * * * % * * * N%%C * * * % * * * % * * * O O O O </pre>	
Uridine (8CI9CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to NH3 and CH3NH2.	

C9H13N	CAS REGN : 121-72-2
	
Benzenamine, N,N,3-trimethyl- (9CI) m-Toluidine, N,N-dimethyl- (8CI)	
PA (exp) : 221+/-2 kcal/mol	Temperature : 298 K
Pollack, S. K., Devlin, J. L., III, Summerhays, K. D. Taft, R. W., and Hehre, W. J. The Site of Protonation in Aniline, J. Am. Chem. Soc. 99, 4583 (1977).	
Method used : ICR Equilibrium; scale based on NH3. Protonation on the nitrogen	

C9H13N

CAS REGN : 613-97-8



Benzenamine, N-ethyl-N-methyl- (9CI)
Aniline, N-ethyl-N-methyl- (8CI)

GB (exp) : 209.40+/-2 kcal/mol

Temperature : 600 K

PA (exp) : 226.60+/-2 kcal/mol

Temperature : 600 K


Yamdagni, R., and Kebarle, P.,
Gas-Phase Basicities of Amines. Hydrogen Bonding
in Proton-Bound Amine Dimers and Proton-Induced
Cyclization of alpha,omega-Diamines,
J. Am. Chem. Soc. 95, 3504 (1973).

Method used : MS
Entropy change assumed zero
Equilibrium; scale based on NH3

C9H15N	CAS REGN : 102-70-5
C++C**C**N**C**C++C * * C * * C + + C	
2-Propen-1-amine, N,N-di-2-propenyl- (9CI) Triallylamine (8CI)	
GB (exp) : 219+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 226.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C9H21N	CAS REGN : 102-69-2
C**C**C**N**C**C**C * * C * * C * * C	
1-Propanamine, N,N-dipropyl- (9CI) Tripropylamine (8CI)	
GB (exp) : 223.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 230.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 222.8+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 230.60+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., A Thermodynamic Analysis of Solvation Effects on the Basicities of Alkylamines. An Electrostatic Analysis of Substituent Effects, J. Am. Chem. Soc. 98, 318 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C9H21N	CAS REGN : 2085-66-7
<pre> C C * * * * C**C**N**C**C**C * * * * C C </pre>	
Propylamine, N-tert-butyl-1,1-dimethyl- (8CI)	
GB (exp) : 221.9+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 230.1+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C ₁₀ H ₈	CAS REGN : 275-51-4
	
Azulene (8CI9CI)	
GB (exp) : 215.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 223.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

C10H9F3	CAS REGN : 55186-75-9
<pre> C . F C C**C**C * . + * . + F**C**C C C * . . * . . F C </pre>	
Benzene, 1-(1-methylethenyl)-4-(trifluoromethyl)- (9CI)	
GB (exp) : 189.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 197.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C10H10Fe	CAS REGN : 102-54-5
<pre> C+++C C+++C * * * * * * * * * * * * * * * * * * * * * * * * C+++C C+++C </pre>	
Ferrocene (8CI9CI)	
PA (exp) : 208+/-4 kcal/mol	Temperature : 300 K
Foster, M. S., and Beauchamp, J. L., Ion-Molecule Reactions and Gas-Phase Basicity of Ferrocene, J. Am. Chem. Soc. 97, 4814 (1975).	
Method used : ICR Bracketing relative to CH3N=NCH3 and CH3NH2.	

C10H12	CAS REGN : 1195-32-0
<pre> C . . . C C**C . . C**C**C C + . + . C C </pre>	
Benzene, 1-methyl-4-(1-methylethenyl)- (9CI) Styrene, p,.alpha.-dimethyl- (8CI)	
GB (exp) : 200.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 208.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C10H12O	CAS REGN : 1712-69-2
<pre> C . . . C C**O**C . . C**C**C C + . + . C C </pre>	
Benzene, 1-methoxy-4-(1-methylethenyl)- (9CI) Anisole, p-isopropenyl- (8CI)	
GB (exp) : 206.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 214.6+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C10H13N	CAS REGN : 4096-21-3
<pre> C . . . C C C*****C . . * * . . * * C C C*****N C . . * * . * * C C </pre>	
Pyrrolidine, 1-phenyl- (8CI9CI)	
GB (exp) : 213.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 221.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C10H13N5	CAS REGN : 715-91-3
<pre> C . N . N . N% C C C*****C . . * * * * . * * * * C N*****C C * * * * * * * * N+++++C C </pre>	
9H-Purin-1-amine, 9-cyclopentyl- (9CI) Adenine, 9-cyclopentyl- (8CI)	
PA (exp) : >222.1 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)3N	

C10H13N5O4	CAS REGN : 58-61-7
<pre> O * * * C * N N++C % * * % * * C..C * O**C . . * * . . * * N C**N*****C * . . * * . . * * C..N C**C * * * * * O O </pre>	
Adenosine (8CI9CI)	
PA (exp) : >222 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to (CH3)3N.	

C10H14	CAS REGN : 98-06-6
<pre> C . . . C C C * . . * . . C**C**C C * . . * . . C C </pre>	
Benzene, (1,1-dimethylethyl)- (9CI) Benzene, tert-butyl- (8CI)	
GB (exp) : 183.2+/-2 kcal/mol	Temperature : 300 K
Hahre, W. J., McIver, R. T., Jr., Pople, J. A., and Schleyer, P. v. R., Alkyl Substituent Effects on the Stability of Protonated Benzene, J. Am. Chem. Soc. 96, 7162 (1974).	
Method used : ICR Equilibrium; scale based on C6H5CH3(Toluene)	

Continued on next page

GB (exp) : 182.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.6+/-2 kcal/mol	Temperature : 300 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on C6H6(Benzene)	

C10H14	CAS REGN : 104-51-8
<pre> C . . . C C . . C**C**C**C**C . . C C . C </pre>	
Benzene, butyl- (8CI9CI)	
GB (exp) : 183+/-2 kcal/mol	Temperature : 300 K
Hehre, W. J., McIver, R. T., Jr., Pople, J. A., and Schleyer, P. v. R., Alkyl Substituent Effects on the Stability of Protonated Benzene, J. Am. Chem. Soc. 96, 7162 (1974).	
Method used : ICR Equilibrium; scale based on C6H5CH3(Toluene)	


Continued on next page

GB (exp) : 183+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.8+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 182.2+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 190.4+/-2 kcal/mol	Temperature : 300 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on C6H6(Benzene)	

C10H14N2O5	CAS REGN : 50-89-5
<pre> O * * C**C * * * * * 0**C**C * * * * * * * O**C*****N * * * * * * C**N % % O C * * C++C * * C%O * * % % O </pre>	
Thymidine (8CI9CI)	
PA (exp) : 207+/-5 kcal/mol	Temperature :
Wilson, M. S., and McCloskey, J. A., Chemical Ionization Mass Spectrometry of Nucleosides. Mechanisms of Ion Formation and Estimations of Proton Affinity, J. Am. Chem. Soc. 97, 3436 (1975).	
Method used : MS Bracketing relative to NH3 and CH3NH2.	

C10H15N

CAS REGN : 91-66-7



Benzenamine, N,N-diethyl- (9CI)
Aniline, N,N-diethyl- (8CI)

GB (exp) : 212.10+/-2 kcal/mol

Temperature : 600 K

PA (exp) : 229.30+/-2 kcal/mol

Temperature : 600 K

Yamdagni, R., and Kebarle, P.,
Gas-Phase Basicities of Amines. Hydrogen Bonding
in Proton-Bound Amine Dimers and Proton-Induced
Cyclization of alpha,omega-Diamines,
J. Am. Chem. Soc. 95, 3504 (1973).


Method used : MS
Equilibrium; scale based on NH3
Entropy change assumed zero

C ₁₀ H ₁₉ N	CAS REGN : 31023-92-4
<pre> C**C * * * * C *N * *C* * * *C * C*C C * * * * * C**C </pre>	
1-Azabicyclo[3.3.3]undecane (9CI)	
PA (exp) : 227.9+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Photoelectron Spectrum and Gas-Phase Basicity of Manxine. Evidence for a Planar Bridgehead Nitrogen, J. Am. Chem. Soc. 97, 4136 (1975).	
Method used : ICR Equilibrium; scale based on CH ₃ NH ₂ Van't Hoff plot	

C10H23N	CAS REGN : 2016-57-1
N**C**C**C**C**C**C**C**C**C	
1-Decanamine (9CI) Decylamine (8CI)	
GB (exp) : >209.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to (CH3)2NH	

C10H24N2	CAS REGN : 111-18-2
C*N**C**C**C**C**C**C*N**C * * * * C C	
1,6-Hexanediamine, N,N,N',N'-tetramethyl- (8CI9CI)	
GB (exp) : 226.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Equilibrium; scale based on NH3	


C11H13N	CAS REGN : 4363-25-1
<pre> N**C * * * * C * C..C * C . . C * . . C * C . ** C**C . . C..C </pre>	
2H-1,4-Ethanoquinoline, 3,4-dihydro- (8CI9CI)	
GB (exp) : 220.5+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 228.3+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C11H15N	CAS REGN : 4096-20-2
	
Piperidine, 1-phenyl- (8CI9CI)	
GB (exp) : 216.1+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 223.9+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C12H11N	CAS REGH : 122-39-4
<pre> C C C . C . C . C C . C*****N**C . C C C </pre>	
Benzenamine, N-phenyl- (9CI) Diphenylamine (8CI)	
GB (exp) : >193.7 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to NH3	

C12H16N2O6

CAS REGN : 362-43-6



Uridine, 2',3'-O-(1-methylethylidene)- (9CI)
Uridine, 2',3'-O-isopropylidene- (8CI)

PA (exp) : 207+/-5 kcal/mol

Temperature :

Wilson, M. S., and McCloskey, J. A.,
Chemical Ionization Mass Spectrometry of Nucleosides.
Mechanisms of Ion Formation and Estimations of
Proton Affinity,
J. Am. Chem. Soc. 97, 3436 (1975).

Method used : MS
Bracketing relative to CH3NH2 and NH3.

C12H18	CAS REGN : 87-85-4
<pre> C C * * * * C..C . . C**C C**C . . C..C * * * * C C </pre>	
Benzene, hexamethyl- (8CI9CI)	
GB (exp) : 197.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 205.0+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C12H19N	CAS REGN : 22025-87-2
<pre> C C..C * . * . C**N**C C..C * * * C**C**C * * C </pre>	
Benzenamine, 2-(1,1-dimethylethyl)-N,N-dimethyl- (9CI) Aniline, o-tert-butyl-N,N-dimethyl- (8CI)	
GB (exp) : 218+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 225.8+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	


C12H27N	CAS REGN : 102-82-9
C**C**C**C**N**C**C**C**C * * C * * C * * C * * C	
1-Butanamine, N,N-dibutyl- (9CI) Tributylamine (8CI)	
GB (exp) : 224.1+/-2 kcal/mol	Temperature : 298 K
PA (exp) : 232+/-2 kcal/mol	Temperature : 298 K
Aue, D. H., Webb, H. M., and Bowers, M. T., Quantitative Proton Affinities, Ionization Potentials, and Hydrogen Affinities of Alkylamines, J. Am. Chem. Soc. 98, 311 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on CH3NH2	

C13H100	CAS REGN : 119-61-9
<pre> C C . . C . . C C . . C C C*****C**C C C 0 C </pre>	
Methanone, diphenyl- (9CI) Benzophenone (8CI)	
GB (exp) : ~200.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : ~203.1+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C13H13OP	CAS REGN : 2129-89-7
<pre> C C C C O C C . . + ... + . . C C C*****P**C C . . x x . . C C C </pre>	
Phosphine oxide, methyldiphenyl- (8CI9CI)	
PA (exp) : 220+/-3 kcal/mol	Temperature : ~400 K
Goff, S. D., Jelus, B. L., and Schweizer, E. E. Electron Impact and Chemical Ionization Mass Spectra of Alkyldiphenylphosphine Oxides, Org. Mass Spectrom. 12, 33 (1977).	
Method used : MS Bracketing relative to pyridine and Me3N scale based on NH3	

C13H17N5O5

CAS REGN : 362-76-5



Guanosine, 2',3'-O-(1-methylethylidene)- (9CI)
 Guanosine, 2',3'-O-isopropylidene- (8CI)

PA (exp) : >222 kcal/mol

Temperature :

Wilson, M. S., and McCloskey, J. A.,
 Chemical Ionization Mass Spectrometry of Nucleosides.
 Mechanisms of Ion Formation and Estimations of
 Proton Affinity,
 J. Am. Chem. Soc. 97, 3436 (1975).

Method used : MS
 Bracketing relative to (CH3)3N.

C13H21N	CAS REGN : 585-48-8
<pre> C . . . C C C C * * * * * * * * C**C**C C**C**C * * * * * * * * C N C </pre>	
Pyridine, 2,6-bis(1,1-dimethylethyl)- (9CI) Pyridine, 2,6-di-tert-butyl- (8CI)	
GB (exp) : 220.6+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 228.4+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C14H12	CAS REGN : 530-48-3
<pre> C C C . . C . . C . . C C . . C*****C**C C . . + ... + . . C C C </pre>	
Benzene, 1,1'-ethenylidenebis- (9CI) Ethylene, 1,1-diphenyl- (8CI)	
GB (exp) : 201.4+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 209.2+/-2 kcal/mol	Temperature : 300 K
Unpublished work by Wolf, Staley, Koppel, Taagepera, McIver, Beauchamp, and Taft.	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

C14H150P	CAS REGN : 1733-57-9
<pre> C C . . C . C O C . C . . + ... + . . C . CxxxxPxxx C . C . . x x x C x x C . . </pre>	
Phosphine oxide, ethyldiphenyl- (8CI9CI)	
PA (exp) : 220+/-3 kcal/mol	Temperature : ~400 K
Goff, S. D., Jelus, B. L., and Schweizer, E. E. Electron Impact and Chemical Ionization Mass Spectra of Alkyldiphenylphosphine Oxides, Org. Mass Spectrom. 12, 33 (1977).	
Method used : MS Bracketing relative to pyridine and Me3N; scale based on NH3	

C15H17OP	CAS REGN : 2959-75-3
<pre> C C . . C . . C O C . . C . . . + ... + . . C . . C * * * * P * * C . . . * * C * * C * * C </pre>	
Phosphine oxide, (1-methylethyl)diphenyl- (9CI) Phosphine oxide, isopropyldiphenyl- (8CI)	
PA (exp) : 220+/-3 kcal/mol	Temperature : ~400 K
Goff, S. D., Jelus, B. L., and Schweizer, E. E. Electron Impact and Chemical Ionization Mass Spectra of Alkyldiphenylphosphine Oxides, Org. Mass Spectrom. 12, 33 (1977).	
Method used : MS Bracketing relative to pyridine and Me3N; scale based on NH3	

C15H17OP	CAS REGN : 4252-88-4
The diagram illustrates the chemical structure of diphenylpropyl phosphine oxide. It features two phenyl rings (C6H5) linked by a three-carbon propyl chain. The middle carbon of this chain is attached to a phosphorus atom (P), which is further bonded to an oxygen atom (O) via a double bond. Additionally, the phosphorus atom is bonded to two methyl groups (CH3) and one hydrogen atom (H).	
Phosphine oxide, diphenylpropyl	
PA (exp) : 220+/-3 kcal/mol	Temperature : ~400 K
Goff, S. D., Jelus, B. L., and Schweizer, E. E. Electron Impact and Chemical Ionization Mass Spectra of Alkyldiphenylphosphine Oxides, Org. Mass Spectrom. 12, 33 (1977).	
Method used : MS Bracketing relative to pyridine and Me3N; scale based on NH3	

C17H21OP	CAS REGN : Not available
<pre> C O C . + . . + . C . . . C**P**C . . . C . * . . . * . . C . . . C . C . C . C . * . . . * . . C . . . C**C**C . C . . * . . * . C </pre>	
Phosphine oxide, neopentyldiphenyl-	
PA (exp) : 220+/-3 kcal/mol	Temperature : ~400 K
Goff, S. D., Jelus, B. L., and Schweizer, E. E. Electron Impact and Chemical Ionization Mass Spectra of Alkyldiphenylphosphine Oxides, Org. Mass Spectrom. 12, 33 (1977).	
Method used : MS Bracketing relative to pyridine and Me3N scale based on NH3	

C18H15N	CAS REGN : 603-34-9
<pre> C C C C C C C C C C C C C C C C C C C C C*****N**C * * * * * C C C C C C C C C C C C C C </pre>	
Benzenamine, N,N-diphenyl- (9CI) Triphenylamine (8CI)	
GB (exp) : >202.9 kcal/mol	Temperature : 423-473 K
Dzidic, I., Relative Gas-Phase Basicities of Some Amines, Anilines, and Pyridines. An Application of Some Bronsted Acids as Reactants in Chemical Ionization Mass Spectrometry, J. Am. Chem. Soc. 94, 8333 (1972).	
Method used : MS Bracketing relative to CH3NH2	

C22H310P	CAS REGN : Not available
<pre> C O C . + . . + . C C**P**C C . * . . * . C C C C C . * . . * . C C C . * . . * . C**C**C**C**C**C**C**C**C**C </pre>	
Phosphine oxide, decanyldiphenyl-	
PA (exp) : 220+/-3 kcal/mol	Temperature : ~400 K
Goff, S. D., Jelus, B. L., and Schweizer, E. E. Electron Impact and Chemical Ionization Mass Spectra of Alkyldiphenylphosphine Oxides, Org. Mass Spectrom. 12, 33 (1977).	
Method used : MS Bracketing relative to pyridine and Me3N scale based on NH3	

Ca	CAS REGN : 7440-70-2
CA	
Calcium (8CI9CI)	
PA (exp) : >211 kcal/mol	Temperature : ~800 K
Eslava, L. A., and Porter, R. F. A Kinetic Study of Proton Transfer Reactions of NH ₄ ⁺ , CH ₃ NH ₃ ⁺ , and PH ₄ ⁺ with Calcium Atoms, Chem. Phys. Letters 52 368 (1977).	
Method used : MS Bracketing relative to NH ₃ , PH ₃ and MeNH ₂	

ClH	CAS REGN : 7647-01-0
CL	
Hydrochloric acid (8CI9CI)	
PA (exp) : 140+/-3 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Absolute PA value estimated by assuming that for reaction $RH^+ + RH \rightarrow R + RH_2^+$ the $\Delta H \sim 5X$ TC. TC = measured translational energy of the products.	
PA (exp) : >120 kcal/mol	Temperature : 297 K
Fehsenfeld, F. C., and Ferguson, E. E. Rate Constants for the Reactions $Cl^+ + H_2 \rightarrow HCl^+ + H$ and $ClH^+ + H_2 \rightarrow ClH_2^+ + H$, J. Chem. Phys. 60, 5132 (1974).	
Method used : FAG Found reaction $ClH^+ + H_2 \rightarrow ClH_2^+ + H$ react rapidly. Since reaction $Cl^+ + H_2 \rightarrow HCl^+ + H + .22eV$ has a low exothermicity, they assumed HCl^+ to be in ground vibrational state.	

CsHO	CAS REGN : 21351-79-1
CS*O	
Cesium hydroxide (8CI9CI)	
PA (exp) : 270.1 kcal/mol	Temperature : 300 K
Searles, S. K., Dzidic, I., and Kebarle, P., Proton Affinities of the Alkali Hydroxides, J. Am. Chem. Soc. 91, 2810 (1969).	
Method used : MS PA derived from delta H of reaction M+ + H2O -> (MOH2)+.	

D2	CAS REGN : 7782-39-0
H**H	
Deuterium (8CI9CI)	
PA (exp) : 98+/-4 kcal/mol	Temperature :
Harris, H. H., Crowley, M. G., Grossheim, T. R., Woessner, P. J., and Leventhal, J. J., Binding Energy of H3+, J. Chem. Phys. 59, 6181 (1973).	
Method used : MS Threshold measurement of CID of D3+ + HE->D2 + D+ + HE. Correction for collision energies distribution.	

Continued on next page

PA (exp) : 103.8+/-1.1 kcal/mol	Temperature :
Cotter, R. J., Rozett, R. W., and Koski, W. S., Reactions of H2O+ and D2O+ with Molecular Hydrogen. I. Proton Affinity of Hydrogen, J. Chem. Phys. 57, 4100 (1972).	
Method used : MS Threshold measurement of D2O+ + D2 -> D3+ + OD. Correction for Doppler effect.	

D2O	CAS REGN : 7789-20-0
0	
Water-d2 (9CI) Water, heavy (D2O) (8CI)	
PA (exp) : 167+/-3 kcal/mol	Temperature :
Cotter, R. J., and Koski, W. S., Reaction of D3O+ with D2: Proton Affinity of Water, J. Chem. Phys. 59, 784 (1973).	
Method used : MS Correction for Doppler effects. Deuterium affinity. Threshold measurement of D2O+ + D2O -> D3O+ + D0.	

FH	CAS REGN : 7664-39-3
F	
Hydrofluoric acid (8CI9CI)	
PA (exp) : 112+/-2 kcal/mol	Temperature : 300 K
Foster, M. S., and Beauchamp, J. L., Proton Affinity and Gas-Phase Ion Chemistry of Hydrogen Fluoride, Inorg. Chem. 14, 1229 (1975).	
Method used : ICR Calculated entropy. Equilibrium relative to N2.	
PA (exp) : 94.3+/-1.4 kcal/mol	Temperature : 0 K
Ng, C. Y., Trevor, D. J., Tiedemann, P. W., Ceyer, S. T., Kronebusch, P. L., Mahan, B. H., and Lee, Y. T. Photoionization of Dimeric Polyatomic Molecules: Proton Affinities of H2O and HF, J. Chem. Phys. 67, 4235 (1977).	
Method used : PI AP of H2F+ from (HF)2. No temperature correction was applied in this supersonic expansion	

F0	CAS REGN : 12061-70-0
F+0	
Oxygen fluoride	
PA (exp) : 134+/-2 kcal/mol	Temperature : 0 K
Berkowitz, J., Appelman, E. H., and Chupka, W. A. Photoionization of HOF with Mass Analysis, J. Chem. Phys. 58, 1950 (1973).	
Method used : MS Threshold measurement of reaction HOF -> HOF+	

F3N	CAS REGN : 7783-54-2
F**N**F * * F	
Nitrogen fluoride (NF3) (8CI9CI)	
PA (exp) : 151+/-10 kcal/mol	Temperature : 300 K
Holtz, D., Beauchamp, J. L., Henderson, W. G., and Taft, R. W., Basicity of Nitrogen Trifluoride in the Gas Phase by Ion Cyclotron Resonance, Inorg. Chem. 10, 201 (1971).	
Method used : ICR Bracketing relative to C2H4 AND HCL.	

F6S	CAS REGN : 2551-62-4
<pre> F F * * ** F**S**F ** * * F F </pre>	
Sulfur fluoride (SF6), (OC-6-11)- (9CI) Sulfur fluoride (SF6) (8CI)	
PA (exp) : <144 kcal/mol	Temperature : 298 K
Gorden, R., Jr., and Sieck, L. W., Ion-Molecule Reactions in Isopropanol: Implications for its Radiolysis, J. Res. NBS 76A, 655 (1972).	
Method used : MS Proton transfer from C2H6+ to SF6 not observed.	

HI	CAS REGN : 10034-85-2
I	
Hydriodic acid (8CI9CI)	
PA (exp) : 145+/-3 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Absolute PA value estimated by assuming that for reaction $RH^+ + RH \rightarrow R + RH_2^+$; the ΔH ~5X TC. TC = measured translational energy of the products.	

HK0	CAS REGN : 1310-58-3
K*0	
Potassium hydroxide (8CI9CI)	
PA (exp) : 263.5 kcal/mol	Temperature : 300 K
Searles, S. K., Dzidic, I., and Kebarle, P., Proton Affinities of the Alkali Hydroxides, J. Am. Chem. Soc. 91, 2810 (1969).	
Method used : MS PA value derived from delta H of reaction: $M^+ + H_2O \rightarrow (MOH_2)^+$	

HLi0	CAS REGN : 1310-65-2
LI*0	
Lithium hydroxide (8CI9CI)	
PA (exp) : 241.6 kcal/mol	Temperature : 300 K
Searles, S. K., Dzidic, I., and Kebarle, P., Proton Affinities of the Alkali Hydroxides, J. Am. Chem. Soc. 91, 2810 (1969).	
Method used : MS PA value derived from delta H of reaction $M^+ + H_2O \rightarrow (MOH_2)^+$	

HN ₂ O	CAS REGN : 1310-73-2
NA ₂ O	
Sodium hydroxide (8CI9CI)	
PA (exp) : 248.5 kcal/mol	Temperature : 300 K
Searles, S. K., Dzidic, I., and Kebarle, P., Proton Affinities of the Alkali Hydroxides, J. Am. Chem. Soc. 91, 2810 (1969).	
Method used : MS PA value derived from delta H of reaction M ⁺ + H ₂ O = (MOH ₂) ⁺	

H ₂	CAS REGN : 1333-74-0
H ⁺ ×H	
Hydrogen (8CI9CI)	
PA (exp) : 99.9+/-1 kcal/mol	Temperature :
Duben, A. J., and Lowe, J. P., Correlation Studies on H ₃ ⁺ . I. The Wavefunctions, J. Chem. Phys. 55, 4270 (1971).	
Method used : TC SCF-MO-CI calculation.	

Continued on next page

PA (exp) : 97 kcal/mol	Temperature : 298 K
Schwartz, M. E., and Schaad, J. J., Ab Initio Studies of Small Molecules Using 1s Gaussian Basis Function. II. H ₃ ⁺ , J. Chem. Physics 47, 5325 (1967).	
Method used : TC SCF-MO-CI calculation.	
PA (exp) : 101.5 kcal/mol	Temperature :
Burt, J. A., Dunn, J. L., McEwan, M. J., Sutton, M. M., Rocke, A. E., and Schiff, H. I., Some Ion-Molecule Reactions of H ₃ ⁺ and the Proton Affinity of H ₂ , J. Chem. Phys. 52, 6062 (1970).	
Method used : FAG Bracketing relative to O ₂ AND N ₂ .	
PA (exp) : 101+/-2 kcal/mol	Temperature :
Cotter, R. J., Rozett, R. W., and Koski, W. S., Reactions of H ₂ O ⁺ and D ₂ O ⁺ with Molecular Hydrogen. I. Proton Affinity of Hydrogen, J. Chem. Phys. 57, 4100 (1972).	
Method used : MS Correction for Doppler effect. Threshold measurement of H ₂ O ⁺ + H ₂ -> H ₃ ⁺ + OH.	

Continued on next page

PA (exp) : 96+/-4 kcal/mol	Temperature :
Harris, H. H., Crowley, M. G., Grossheim, T. R., Woessner, P. J., and Leventhal, J. J., Binding Energy of H₃⁺, J. Chem. Phys. 59, 6181 (1973).	
Method used : MS Threshold measurement of CID of H₃⁺ + He -> H₂ + H⁺ + He. Correction for collision energies distribution.	

H ₂ O	CAS REGN : 7732-18-5
0	
Water (8CI9CI)	
PA (exp) : 167+/-7 kcal/mol	Temperature : 323-373 K
Long, J., and Munson, B., Proton Affinities of Some Oxygenated Compounds, J. Am. Chem. Soc. 95, 2427 (1973).	
Method used : MS Bracketing relative to CH₂O and C₂H₄	

Continued on next page

PA (exp) : 182 kcal/mol	Temperature :
Sherman, J., Crystal Energies of Ionic Compounds and Thermochemical Applications, Chem. Rev. 11, 93 (1932).	
Method used : CLE Using thermochemical cycle the value of PA(H ₂ O) was calculated from PA(NH ₃).	
PA (exp) : 171+/-2 kcal/mol	Temperature : 296,355,426 K
Hopkins, J. M., and Bone, L. I., Relative Proton Affinities of Hydrogen Sulfide and Water, J. Chem. Phys. 58, 1473 (1973).	
Method used : MS Van't Hoff plot Equilibrium relative to H ₂ S. Assume steady state kinetics.	
PA (exp) : 165.8+/-1.8 kcal/mol	Temperature : 0 K
Ng, C. Y., Trevor, D. J., Tiedemann, P. W., Ceyer, S. T., Kronebusch, P. L., Mahan, B. H., and Lee, Y. T. Photoionization of Dimeric Polyatomic Molecules: Proton Affinities of H ₂ O and HF, J. Chem. Phys. 67, 4235 (1977).	
Method used : PI AP of H ₃ O ⁺ from (H ₂ O) ₂ . No temperature correction was applied in this supersonic expansion	

Continued on next page

GB (exp) : 153.3+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 169.3+/-2 kcal/mol	Temperature : 600 K
Yamdaqni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 162.3+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 170.3+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

H2S	CAS REGN : 7783-06-4
S	
Hydrogen sulfide (H2S) (8CI9CI)	
GB (exp) : 165.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 173.9+/-2 kcal/mol	Temperature : 300 K
Freeman, C. G., Harland, P. W., and McEwan, M. J. The Equilibrium $\text{H}_3\text{S}^+ + \text{HCN} = \text{H}_2\text{CN}^+ + \text{H}_2\text{S}$ and the Relative Proton Affinities of HCN and H2S, Intern. J. Mass Spectrom. Ion Phys. 27, 77 (1978).	
Method used : FAG Equilibrium relative to HCN; scale based on NH3	
PA (exp) : 173.3+/-2 kcal/mol	Temperature : 296,355,426 K
Hopkins, J. M., and Bone, L. I., Relative Proton Affinities of Hydrogen Sulfide and Water, J. Chem. Phys. 58, 1473 (1973).	
Method used : MS Equilibrium relative to H2O Van't Hoff plot	

Continued on next page

PA (exp) : 177.1+/-2 kcal/mol	Temperature : 299,329,360 K
Wei, L. Y., and Bone, L. I., Ion-Molecule Reactions in Methanol and Hydrogen Sulfide, J. Phys. Chem. 78, 2527 (1974).	
Method used : MS Equilibrium; scale based on CH ₃ OH Van't Hoff plot	
PA (exp) : 171 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Heats of Formation of H ₃ O ⁺ , H ₃ S ⁺ , and NH ₄ ⁺ by Electron Impact, J. Chem. Phys. 50, 2028 (1969).	
Method used : EI Appearance potential by electron impact. Correction for excess energy.	
GB (exp) : 165.9+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 173.9+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH ₃	

Continued on next page

GB (exp) : 156.4+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 172.4+/-2 kcal/mol	Temperature : 600 K
Yamdagni, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : MS Calculated entropy Equilibrium; scale based on NH3	
GB (exp) : 172.4+/-2.5 kcal/mol	Temperature : 300 K
PA (exp) : 175.3+/-2.5 kcal/mol	Temperature : 300 K
Meot-Ner (Mautner), M., and Field, F. H. Proton Affinities and Cluster Ion Stabilities in CO2 and CS2. Application in Martian Ionospheric Chemistry, J. Chem. Phys. 66, 4527 (1977).	
Method used : MS Calculated entropy The measurements done in the temperature range of 400-650K. Using the Hoff plot the H and S at 300 K were derived. Equilibrium; scale based on NH3	

H ₂ Se	CAS REGN : 7783-07-5
SE	
Hydrogen selenide (H ₂ Se) (8CI9CI)	
GB (exp) : 166.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 174.7+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH₃	

H3N	CAS REGN : 7664-41-7
N	
Ammonia (8CI9CI)	
PA (exp) : >197 kcal/mol	Temperature :
Chupka, W. A., and Russell, M. E., Ion-Molecule Reactions of NH3+ by Photoionization, J. Chem. Phys. 48, 1527 (1968).	
Method used : PI NH3+ formed by photoionization and react with H2O.	
PA (exp) : 206.4 kcal/mol	Temperature :
Bleick, W. E., The Lattice Energies of the Ammonium Halides and the Proton Affinity of Ammonia, J. Chem. Phys. PA = 2, 160 (1934).	
Method used : CLE Using thermochemical cycle the PA is derived from crystal lattice energy.	

Continued on next page

PA (exp) : 208 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Heats of Formation of H_3O^+, H_3S^+, and NH_4^+ by Electron Impact, J. Chem. Phys. 50, 2028 (1969).	
Method used : EI Appearance potential by electron impact. Correction for excess energy.	
PA (exp) : 194+/-7 kcal/mol	Temperature :
Vetchinkin, S. I., Pshenichnov, E. A., and Sokolov, N. D., The Effect of the Hydrogen Bond on the Energy of the Ionic Lattice of Ammonium Chloride and Estimation of the Proton Affinity of an Ammonia Molecule, Zh. Fiz. Khim. 33, 1269 (1959).	
Method used : CLE PA from crystal lattice energy, correction for hydrogen bond.	
PA (exp) : 206+/-8 kcal/mol	Temperature :
Sherman, J., Crystal Energies of Ionic Compounds and Thermochemical Applications, Chem. Rev. 11, 93 (1932).	
Method used : CLE Using thermochemical cycle the PA values 209, 208.6 and 202.7 are derived from crystal lattice energy of NH_4Cl, NH_4Br and NH_4I respectively.	

Continued on next page

GB (exp) : 185.1+/-2 kcal/mol	Temperature : 600 K
PA (exp) : 202.3+/-2 kcal/mol	Temperature : 600 K
Yamdaqui, R., and Kebarle, P., Gas-Phase Basicities and Proton Affinities of Compounds between Water and Ammonia and Substituted Benzenes from a Continuous Ladder of Proton Transfer Equilibria Measurements, J. Am. Chem. Soc. 98, 1320 (1976).	
Method used : ICR Calculated entropy Equilibrium; scale based on PA(iso-C4H8)	
GB (exp) : 193.7+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 202.3+/-2 kcal/mol	Temperature : 300 K
Staley, R. H., Taagepera, M., Henderson, W. G., Koppel, I., Beauchamp, J. L., and Taft, R. W., Effects of Alkyl and Fluoroalkyl Substitution on the Heterolytic and Homolytic Bond Dissociation Energies of Protonated Amines, J. Am. Chem. Soc. 99, 326 (1977).	
Method used : ICR Calculated entropy Equilibrium.	

Continued on next page

PA (exp) : 205.7+/-1 kcal/mol	Temperature : 298.15 K
Meot-Ner (Mautner), M., and Field, F. H. Proton Affinities and Cluster Ion Stabilities in CO₂ and CS₂. Application in Martian Ionspheric Chemistry, J. Chem. Phys. 66, 4527 (1977).	
Method used : CLE Using thermochemical cycle the PA value is derived from crystal lattice energy. Error due to some of the thermochemical data and the similarity of crystal structure.	
PA (exp) : 205.3 kcal/mol	Temperature :
Goodliffe, A. L., Jenkins, H. D. B., Martin, S. V., and Waddington, T. C., The Proton Affinity of Gaseous Ammonia, the Charge Distribution on the NH₄⁺ Ion and the Lattice Energies of NH₄Cl, NH₄Br, and NH₄I, Mol. Phys. 21, 761 (1971).	
Method used : CLE Using thermochemical cycle the PA is derived from crystal lattice energy and the distributed charge on NH₄⁺.	

H3P	CAS REGN : 7803-51-2
P	
Phosphine (8CI9CI)	
PA (exp) : 200+/-10 kcal/mol	Temperature : 298 K
Wendlendt, W., Proton Affinity of Phosphine in the Phosphonium Halides, Science 122, 831 (1955).	
Method used : CLE Using thermochemical cycle calculate the PA value from crystal lattice energy.	
PA (exp) : 156.2+/-3 kcal/mol	Temperature :
Wada, Y., and Kiser, R. W., A Mass Spectrometric Study of Some Alkyl-Substituted Phosphines, J. Phys. Chem. 68, 2290 (1964).	
Method used : EI Appearance potential by electron impact of (CH3)3P and (C2H5)3P	

Continued on next page

PA (exp) : 194.5+/-5 kcal/mol	Temperature :
Waddington, T. C., Lattice Energies of Phosphonium Bromide and Iodide and the Proton Affinity of Phosphine, Trans. Faraday Soc. 61, 2652 (1965).	
Method used : CLE Using thermochemical cycle they calculated the PA from crystal lattice energy and heats of formation.	
PA (exp) : 186 kcal/mol	Temperature :
Haney, M. A., and Franklin, J. L., Mass Spectrometric Determination of the Proton Affinities of Various Molecules, J. Phys. Chem. 73, 4328 (1969).	
Method used : MS Delta H estimated = 5X total translational energy of the products of $RH^+ + RH \rightarrow RH_2^+ + R$.	

Continued on next page

GB (exp) : 178.8+/-2 kcal/mol	Temperature : 300 K
PA (exp) : 187.4+/-2 kcal/mol	Temperature : 300 K
Wolf, J. F., Staley, R. H., Koppel, I., Taagepera, M., McIver, R. T., Jr., Beauchamp, J. L., and Taft, R. W., Gas Phase Basicities and Relative Proton Affinities of Compounds between Water and Ammonia from Pulsed Ion Cyclotron Resonance Thermal Equilibria Measurements, J. Am. Chem. Soc. 99, 5417 (1977).	
Method used : ICR Calculated entropy Equilibrium; scale based on NH3	

H4N2	CAS REGN : 302-01-2
N*N	
Hydrazine (8CI9CI)	
PA (exp) : 221 kcal/mol	Temperature :
Jen, J. A., and Thomas, T. D., Core Ionization Potentials in Dimethyl Ether and Methyl Amine, J. Electron Spectrosc. 4, 43 (1974).	
Method used : CIP Assuming that ΔH of $CH_4 + NH_2NH_3^+ \rightarrow CH_3NH_2 + NH_4^+$ is equal to the difference in carbon 1s ionization potentials of CH_4 and CH_3NH_2. CIP = core ionization potential.	

H4Si	CAS REGN : 7803-62-5
SI	
Silane (8CI9CI)	
PA (exp) : 155+/-3 kcal/mol	Temperature :
Cheng, T. M. H., and Lampe, F. W., SiH5+ and the Proton Affinity of Monosilane, Chem. Phys. Letters 19, 532 (1973).	
Method used : MS Bracketing relative to C2H2 AND C3H7.	

He	CAS REGN : 7440-59-7
HE	
Helium (8CI9CI)	
PA (exp) : 41.5 kcal/mol	Temperature :
Bel'skii, V. E., and Izmailov, R. I., Proton Affinity of Atoms and Ions, Zh. Obshch. Khim. 44, 2297 (1974).	
Method used : TC Using thermochemical cycles.	

Continued on next page

PA (exp) : 42 kcal/mol	Temperature :
Beauchamp, J. L., Interactions between Ions and Molecules, Ed. P. Ausloos, (Plenum Press, New York and London, 1975) p. 413.	
Method used : HE Literature Survey.	

Kr	CAS REGN : 7439-90-9
KR	
Krypton (8CI9CI)	
PA (exp) : 115+/-15 kcal/mol	Temperature :
Munson, M. S. B., and Field, F. H., Reactions of Gaseous Ions. XVI. Effects of Additives on Ionic Reactions in Methane, J. Am. Chem. Soc. 87, 4243 (1965).	
Method used : MS Bracketing relative to H2 and CH4.	

Continued on next page

PA (exp) : 100.3+/-0.5 kcal/mol	Temperature :
Payzant, J. D., Schiff, H. I., and Bohme, D. K., Determination of the Proton Affinity from the Kinetics of Proton Transfer Reactions. V. The Equilibrium $H_3^+ + Kr = KrH^+ + H_2$ and the Relative Proton Affinity of Kr and H_2 , J. Chem. Phys. 63, 149 (1975).	
Method used : FAG Equilibrium relative to H_2 . Delta S estimated from the entropies of the particles.	

Mg	CAS REGN : 7439-95-4
MG	
Magnesium (8CI9CI)	
PA (exp) : ~198 kcal/mol	Temperature : 350 K
Po, P. L., and Porter, R. F. High-Temperature Ion-Molecule Chemistry. A Kinetic Study of Gas-Phase Reactions of Magnesium Atoms with D_3^+ , Methanium, Ammonium, and $tert-C_4H_9^+$ Ions, J. Am. Chem. Soc. 99, 4922 (1977).	
Method used : MS Bracketing relative to $(CH_3)_2C=CH_2$ and NH_3	

Mg ₂	CAS REGN : 29904-79-8
Mg+Mg	
Magnesium molecule	
PA (exp) : 219+/-7 kcal/mol	Temperature : 700 K
Po, P. L., and Porter, R. F. A Thermodynamic Study of the Reactions of Mg+(g) and MgH+(g) with Magnesium, J. Phys. Chem. 81, 2233 (1977).	
Method used : MS Calculated entropy From the K equilibrium of reaction MgH+ + Mg(s) = Mg ₂ H+ and using PA (Mg) = 196 and delta Hf(H+) = 365.7	

NO	CAS REGN : 10102-43-9
N++0	
Nitrogen oxide (NO) (8CI9CI)	
PA (exp) : <127 kcal/mol	Temperature :
Munson, M. S. B., and Field, F. H., Reactions of Gaseous Ions. XVI. Effects of Additives on Ionic Reactions in Methane, J. Am. Chem. Soc. 87, 4243 (1965).	
Method used : MS Bracketing relative to CH4.	
PA (exp) : 126.9+/-1.1 kcal/mol	Temperature : 300 K
Hemsworth, R. S., Payzant, J. D., Schiff, H. I., and Bohme, D. K., Rate Constants at 297 K for Proton Transfer Reactions with NH3 Comparisons with Classical Theories and Exothermicity, Chem. Phys. Letters 26, 417 (1974).	
Method used : FAG Equilibrium relative to H2. Use PA (H2) from J. Chem. Phys. 55, 4270 (1971).	

N2	CAS REGN : 7727-37-9
N##N	
Nitrogen (8CI9CI)	
PA (exp) : 112.6+/-1 kcal/mol	Temperature : 298 K
Schiff, H. I., and Bohme, D. K., Flowing Afterglow Studies at York University, Int. J. Mass Spectrom. Ion Phys. 16, 167 (1975).	
Method used : FAG Equilibrium scale based on H2. Delta S derived from theoretical entropies and entropies of iso-electronic particles.	
PA (exp) : 130+/-2 kcal/mol	Temperature :
Willis, C., Lossing, F. P., and Back, R. A. The Heat of Formation of N2H2 and the Proton Affinity of N2, Can. J. Chem. 54, 1 (1976).	
Method used : EM The PA is derived from the appearance potentials of N2+ and N2H+	

N2O	CAS REGN : 10024-97-2
O++N++N	
Nitrogen oxide (N2O) (8CI9CI)	
PA (exp) : 138.4 kcal/mol	Temperature : 298 K
Schiff, H. I., and Bohme, D. K., Flowing Afterglow Studies at York University, Int. J. Mass Spectrom. Ion Phys. 16, 167 (1975).	
Method used : FAG Calculated entropy Equilibrium; scale based on CO.	

O2	CAS REGN : 7782-44-7
O++O	
Oxygen (8CI9CI)	
PA (exp) : 100.9+/- .5 kcal/mol	Temperature : 298 K
McCulloh, K. E., To be published.	
Method used : PI Photoionization of H2O2.	

Continued on next page

PA (exp) : 100.4+/-1.5 kcal/mol	Temperature : 298 K
Kim, J. K., Theard, L. P., and Huntress, W. T., Jr., Proton Transfer Reactions from H3+ Ions to N2, O2, and CO Molecules, Chem. Phys. Letters 32, 610 (1975).	
Method used : ICR Equilibrium relative to H2. Entropy change estimated from the known theoretical iso-electronic counterpart entropies of the species.	
PA (exp) : 100+/-1 kcal/mol	Temperature : 298 K
Fennelly, P. F., Hemsworth, R. S., Schiff, H. I., and Bohme, D. K., Determination of the Proton Affinity from the Kinetics of Proton Transfer Reactions. IV. The Equilibrium $O_2H^+ + H_2 = H_3^+ + O_2$ and the Relative Proton Affinity of O2 and H2, J. Chem. Phys. 59, 6405 (1973).	
Method used : FAG Equilibrium relative to H2. Entropy change estimated from the known theoretical iso-electronic counterpart entropies of the species.	

03S	CAS REGN : 7446-11-9
0++S++0 + + 0	
Sulfur trioxide (8CI9CI)	
PA (exp) : 142+/-2 kcal/mol	Temperature :
Munson, B., Smith, D., and Polley, C. The Mass Spectrum, Proton Affinity and Ion-Molecule Reactions of SO ₃ , Intern. J. Mass Spectrom. Ion Phys. 25, 323 (1977).	
Method used : MS Bracketing relative to HBr and CO. Use $\Delta H_f(H^+) = 366$.	

U	CAS REGN : 7440-61-1
U	
Uranium (8CI9CI)	
PA (exp) : 238+/-4 kcal/mol	Temperature : 400 K
Armentrout, P., Hodges, R., and Beauchamp, J. L. Metal Atoms as Superbases: the Gas Phase Proton Affinity of Uranium, J. Am. Chem. Soc. 99, 3162 (1977).	
Method used : MS Threshold measurement of UD+. Correction for D(UD+) - D(UH+) assuming $\Delta H_{400} = \Delta H_{298}$ Entropy change assumed zero	

Xe	CAS REGN : 7440-63-3
XE	
Xenon (8CI9CI)	
PA (exp) : 118+/-11 kcal/mol	Temperature :
Munson, M. S. B., and Field, F. H., Reactions of Gaseous Ions. XVI. Effects of Additives on Ionic Reactions in Methane, J. Am. Chem. Soc. 87, 4243 (1965).	
Method used : MS Bracketing relative to CH4 and reaction XE+(2P1/2) + CH4.	
PA (exp) : 113.2+/-1.5 kcal/mol	Temperature : 800 K
Fehsenfeld, F. C., Lindinger, W., Schiff, H. I., Hemsworth, R. S., and Bohme, D. K. Determination of the Proton Affinity from the Kinetics of Proton Transfer Reactions. VI. The Relative Proton Affinities of N2, Xe, and CO2, J. Chem. Phys. 64, 4887 (1976).	
Method used : FAG Van't Hoff plot Equilibrium relative to N2; scale based on PA (N2) = 112.6	

123.0	41
153.3	453
154.7	46
156.4	456
162.3	453
162.5	24
164.4	55
165.1	47
165.4	108,253
165.7	42,454
165.9	256,455
166.5	33
166.7	20,457
166.8	51,265
166.9	19,267
167.0	267
167.2	22
167.3	54
167.8	270
169.5	96
169.6	13
169.7	43
169.8	65
170.1	60
171.2	52,83,246
171.8	57,66
171.9	25
172.0	269
172.2	69
172.3	2,116,144
172.4	456
172.5	97
172.7	247
173.0	98
173.1	47,154,248
173.4	249-250
173.7	251
173.9	119,257
174.1	252,334
174.2	264
174.3	254
174.4	34,255
174.5	85,266
174.7	269
175.1	370
175.4	151
175.6	123
176.1	174
176.4	217
176.6	294
176.7	268
177.1	157
177.2	64
177.4	162
177.5	34
177.6	361
178.1	35

178.3	103
178.4	82
178.8	464
178.9	83
179.0	68, 114
179.1	276
179.2	57, 329
179.6	333
179.8	67, 121
179.9	68
180.1	99, 336
180.2	116
180.8	88, 335
180.9	326, 335, 370
181.0	132
181.1	277
181.3	85, 360
181.4	191
181.8	369, 371, 391
181.9	134, 172
182.0	372-373
182.2	389, 413
182.4	104, 177, 411
182.5	190
182.6	390-391
182.7	86
182.8	169, 227
182.9	230, 359, 372
183.0	367, 388, 412-413
183.2	330, 410
183.4	123
183.5	337
183.6	159
183.7	122, 368
183.9	175
184.1	222
184.2	134
184.4	158
184.5	174
184.7	173, 232
185.1	162, 460
185.4	153
185.8	179
186.0	44
186.2	133
186.3	327
186.5	114
186.7	120
186.8	193, 295
187.2	373
187.4	218
187.6	121
188.4	352-353
188.5	75
188.6	176
188.7	189
188.9	61, 328

189.0	171
189.2	170
189.3	402
189.7	228
189.8	124, 178, 325
190.0	190, 300
190.2	89
190.3	177
190.4	330
190.5	180
190.7	296
190.8	229
191.1	225, 231
191.3	392
191.5	362
191.7	237
191.8	308
191.9	62, 313
192.1	181
192.2	152, 309
192.3	221
192.5	226
192.9	135, 236
193.1	365
193.4	38, 278
193.7	279, 291-292, 421, 460
193.9	298
194.0	279, 364
194.4	219
195.0	76
195.2	194, 384
195.3	363
195.5	314
195.6	312
195.9	37
196.1	111
196.6	315
196.9	385
197.0	274, 387
197.2	81, 220
197.5	72, 272
197.7	423
197.8	263
197.9	152
198.0	273
198.4	210
198.5	271
198.9	297
199.0	147, 262
199.5	316
199.6	110
199.8	261
199.9	280
200.2	339, 347
200.4	278
200.6	105, 108, 208
200.7	426

200.8	404
200.9	87,168
201.2	129,146,258
201.4	430
201.8	80
202.2	259
202.5	260
202.9	36,293,437
203.0	366,377
203.2	340-341
203.4	92,235
203.6	374
203.7	207,213
203.9	167,348
204.0	204
204.5	205
204.6	74
204.7	166
204.8	311
204.9	74
205.3	126
205.4	185
205.6	90,92,284
205.7	37
205.9	90
206.1	186,338
206.4	145,209
206.6	331
206.8	332,376,406
207.1	139
207.2	206
207.3	140
207.6	197
207.7	196
207.8	197
208.0	238
208.1	137,318
208.2	137,355
208.3	196
208.4	139
208.6	184
208.8	242
208.9	200
209.0	242
209.3	141,200,386
209.4	395
209.5	91,93
209.6	234
209.7	91,212,351,355,376,417
209.8	213
210.1	310
210.2	195
210.3	212
210.4	195
210.9	188
211.1	239
211.4	287

211.7	239
211.9	127
212.0	141
212.1	415
212.2	127, 140
212.3	375
213.0	282
213.3	182, 289
213.6	94, 300, 407
213.7	138, 182
213.8	283
214.0	95
214.1	198
214.3	138, 281
214.4	233
214.5	193
214.9	142, 233
215.5	94
215.9	401
216.0	199, 243
216.1	320, 420
216.5	320
216.7	199
217.0	285
217.2	307, 379
217.3	156, 235
217.7	378
217.8	234
217.9	307, 317
218.0	424
218.3	312
218.4	240, 317
218.6	357
218.8	311
218.9	240-241
219.0	398
219.1	143
219.9	324, 380
220.0	245, 323
220.5	419
220.6	319, 321, 429
220.8	354
221.1	143
221.2	319, 323
221.3	356
221.4	245
221.7	201, 382
221.9	400
222.8	399
223.1	399
223.6	381
224.0	322
224.1	425
224.6	322
224.9	351
225.4	244
226.9	418

227.7	358
229.3	383

41.5	465
42.0	466
94.3	443
96.0	451
97.0	450
98.0	441
99.9	449
100.0	1,472
100.3	467
100.4	472
100.9	471
101.0	450
101.5	450
103.8	442
112.0	443
112.6	470
113.2	475
115.0	3-4,466
118.0	475
120.0	440
121.0	17
123.0	42
124.0	41
126.0	31
126.9	469
127.0	30,469
128.0	79
129.1	40
130.0	470
130.5	31
130.7	40
131.4	31
131.8	80
134.0	444
138.4	471
139.0	19
139.6	79
140.0	440
141.0	11
142.0	473
142.7	39
143.0	39,130
144.0	9,446
145.0	447
147.0	5,18,20
148.2	130
151.0	27,445
152.8	131
155.0	465
156.2	462
160.0	27
160.6	59
162.0	21
163.0	25
164.0	26,187
165.0	71

165.8	452
166.0	49
166.8	187
167.0	49,442,451
168.0	42
169.0	22
169.3	453
170.0	22,70
170.3	26,453
171.0	69,452,455
171.6	255
171.9	46
172.0	10,45-46
172.2	55
172.4	456
173.3	47,454
173.6	53
173.9	454-455
174.0	28,53
174.5	19-20
174.6	22
174.7	457
174.8	48
175.0	73
175.3	456
176.0	100-101
177.0	13
177.1	455
177.4	13
177.5	43
177.7	96
177.9	60
178.0	23,48
178.5	16,246
178.9	24
179.0	12,24,32,52
179.8	253
180.0	107,276
180.1	25,144,269
180.3	12,97,247-248
180.4	249
180.5	250
180.6	106,257
180.8	98
180.9	2,47,154,165
181.0	78,164
181.1	103,252
181.2	270
181.3	251,254
181.4	269
181.6	256
181.9	163,264
182.0	29,32-33,452
182.1	33
182.2	34,107,266
182.3	265
182.5	265,267

182.6	108,267,334
183.2	151,160
183.8	334
184.0	51,63,87
184.2	217
184.4	294
184.5	54
184.9	157
185.0	64-65
185.3	34
185.4	65,361
185.9	35
186.0	56,63,77,82,463
186.1	103
186.2	82
186.7	83
186.8	23,83
187.0	8,57,68,115
187.4	57,333,464
187.5	333
187.7	68
187.8	69
187.9	99,116,336
188.0	66-67,113,116
188.2	66
188.3	99
188.5	277
188.6	335
188.7	326,335
188.8	132
189.0	58,84,133,372
189.1	85,360,370
189.3	85,88,191
189.4	64
189.5	56
189.6	334,371
189.7	134,172,373
189.8	150,359,372
190.0	84,115,132,367,391
190.1	86
190.2	104
190.3	30
190.4	389,391,413
190.6	169,227,370,411
190.8	413
190.9	175
191.1	119,122,368
191.2	123,159
191.4	159
191.7	174
191.9	222
192.0	109,131,134,192
192.1	173
192.2	158,268
192.3	174
192.5	232
192.6	161

193.0	6
193.2	153, 179
193.3	86
193.5	162, 224
193.6	112
193.7	161, 170
193.8	114
193.9	114
194.0	102, 117, 120, 122, 133, 173, 192, 459
194.1	50, 120, 327
194.2	162
194.4	223
194.5	463
194.6	193, 276, 373
194.7	14, 329
195.0	295
195.2	218
195.4	121
195.7	112
195.9	50, 225
196.0	113, 176
196.1	189, 304
196.4	171, 305
196.6	352
196.7	328
197.0	62, 118, 125, 170, 176, 178, 458
197.1	228, 402
197.2	124
197.3	190, 224
197.4	190, 300
197.6	89, 178, 325
197.7	303
197.9	61
198.0	177, 189, 467
198.1	177
198.2	306, 330
198.3	180
198.4	392
198.5	225
198.6	229
198.7	330
198.8	350
198.9	231
199.0	7, 35, 337
199.3	313, 362
199.5	237
199.6	308
199.9	181
200.0	202, 230, 309, 462
200.1	221, 230
200.3	62, 226
200.4	14
200.7	135, 236
200.9	365
201.0	210
201.7	298
201.8	33, 364

202.2	219
202.3	460
202.6	194
202.8.	76
203.0	312,384
203.1	363
203.3	314
204.0	299,315
204.3	111
204.5	81,111
204.6	220
204.7	385
205.0	423
205.2	337
205.3	461
205.4	274
205.6	263
205.7	75,461
205.9	72,152,272
206.0	459
206.2	210
206.4	147,273,458
206.8	81,262
206.9	271,316
207.0	21,148-149,203,216,393,397,414,422
207.1	260,297
207.6	261
207.8	110
208.0	403,459
208.1	426
208.3	15,87
208.4	208
208.6	146,404
208.7	168
208.8	278
208.9	278,280
209.0	15,105,108,129,258
209.2	430
209.4	152
210.0	259
210.2	80
210.3	260
210.4	377
210.6	136
210.8	366
211.0	36,439
211.2	235,279
211.3	36
211.5	207
211.6	340-342
211.8	204
212.3	167,205,348
212.4	166
212.8	74
213.0	75
213.1	37
213.7	37,126

213.8	145,284
213.9	186
214.0	90,349
214.2	209
214.3	90
214.4	331
214.5	338
214.6	332,406
215.0	206,211
215.5	139
215.7	140
216.0	197
216.2	197
216.4	238
216.5	94,137,318,355
216.6	137
216.7	196
216.8	184,301
217.1	386
217.2	242,286
217.3	93
217.4	242,339,347
217.6	213
217.7	91,200
217.9	91
218.1	212
218.5	310
218.6	195
218.8	195
218.9	188
219.0	468
219.2	287
219.4	128
219.5	239
219.8	288
220.0	127,155,214-215,275,375,427,431-436,438
220.1	127,239,375
220.2	141
220.4	140
220.5	183
220.6	92
220.8	282,374
220.9	213
221.0	141,394,464
221.1	289
221.4	407
221.5	138,182
221.6	283
221.8	300
221.9	182
222.0	290,311,396,405,409,428
222.1	138,281,408
222.3	193
222.6	233
222.7	198
223.1	233
223.3	142,401

223.5	346
223.8	199
223.9	420
224.0	376
224.2	243
224.3	320
224.5	199, 345
224.7	320
224.8	285
225.1	156, 235, 343
225.4	379
225.5	308
225.6	139, 234
225.8	424
225.9	307, 378
226.1	312, 317, 344
226.2	240
226.3	302
226.4	357
226.6	311, 317, 395
226.7	240-241
226.8	234, 398
227.5	94
227.7	324
227.9	416
228.1	380
228.3	419
228.4	319, 321, 429
228.6	354
229.0	319
229.1	356
229.3	415
229.9	382
230.1	400
230.6	399
230.9	399
231.4	381
231.6	143
232.0	425
232.4	322
232.7	351
234.6	245
235.0	323
235.1	143, 245
235.4	323
236.6	201
238.0	474
241.6	448
248.5	449
263.5	448
270.1	441

50-00-0	21-22
50-89-5	414
57-14-7	93
58-61-7	409
58-96-8	393
60-29-7	189-190
60-35-5	75
62-53-3	278-280
64-17-5	82-83
64-18-6	23-25
64-19-7	66-67
65-46-3	396
65-71-4	216
66-22-8	149
67-56-1	32-34
67-63-0	131-132
67-64-1	112-114
67-68-5	87
68-12-2	129
68-94-0	211
71-23-8	132-133
71-30-7	155
71-36-3	191
71-43-2	269-270
73-24-5	214
73-40-5	215
74-82-8	30-31
74-83-9	25
74-84-0	79-80
74-85-1	59
74-87-3	26-27
74-88-4	28
74-89-5	36-37
74-90-8	19-20
74-93-1	34-35
74-96-4	69
74-98-6	130-131
74-99-7	100
75-00-3	70
75-02-5	53
75-03-6	73
75-04-7	90
75-05-8	56-57
75-07-0	63-65
75-08-1	88
75-10-5	20
75-15-0	42
75-18-3	89
75-19-4	106
75-21-8	65
75-31-0	137
75-33-2	134
75-38-7	48
75-46-7	18
75-50-3	138-139
75-52-5	29
75-64-9	195

75-65-0	192
75-66-1	193
75-73-0	17
75-89-8	55
75-97-8	308
76-05-1	46-47
78-81-9	196
78-82-0	158
78-84-2	169
78-93-3	170
79-09-4	119
79-11-8	51
79-20-9	120-121
79-24-3	77
87-42-3	203
87-85-4	423
90-04-0	347
91-66-7	415
93-58-3	365
95-47-6	367-368
95-54-5	291
96-22-0	225
96-47-9	226
98-06-6	410-411
98-82-8	388-389
98-83-9	387
98-86-2	363
98-95-3	268
100-41-4	369-371
100-42-5	362
100-46-9	338
100-47-0	329
100-48-1	261
100-52-7	330
100-54-9	262
100-61-8	339
100-66-3	337
100-70-9	263
100-76-5	354-355
102-54-5	403
102-69-2	399
102-70-5	398
102-82-9	425
103-65-1	390-391
103-69-5	374
104-51-8	412-413
104-87-0	364
104-88-1	325
104-94-9	348
105-07-7	361
106-42-3	372
106-47-8	271
106-49-0	340
106-50-3	292
106-97-8	187
106-98-9	160
107-03-9	134

107-10-8	139-140
107-11-9	126
107-12-0	104
107-13-1	99
107-14-2	47
107-15-3	94
107-31-3	67-69
107-47-1	377
108-18-9	317
108-20-3	312
108-38-3	373
108-42-9	272
108-44-1	341-342
108-45-2	293
108-47-4	343
108-48-5	344
108-67-8	392
108-88-3	333-336
108-89-4	281
108-90-7	264-265
108-91-8	310-311
108-94-1	299
108-95-2	276
108-99-6	282
109-06-8	283
109-09-1	204
109-55-7	244
109-60-4	230
109-73-9	197
109-74-0	159
109-76-2	143
109-77-3	96
109-85-3	141
109-89-7	198
109-94-4	122-123
109-97-7	152
109-99-9	171
110-00-9	150
110-18-9	322
110-58-7	238
110-59-8	222
110-60-1	201
110-62-3	227
110-74-7	173-174
110-85-0	188
110-86-1	212-213
110-89-4	233-234
110-91-8	184
110-95-2	358
110-96-3	378
111-18-2	418
111-26-2	318
111-43-3	313
111-47-7	315
111-51-3	383
111-68-2	355
111-92-2	379

115-07-1	107-108
115-10-6	84-86
115-11-7	161-162
115-20-8	52
119-61-9	426
120-94-5	234-235
121-44-8	319
121-69-7	375-376
121-72-2	394
122-39-4	421
123-11-5	366
123-38-6	115-116
123-39-7	76
123-54-6	220
123-72-8	172
123-75-1	182
123-91-1	175
124-02-7	300
124-09-4	323
124-38-9	40-42
124-40-3	91-92
127-19-5	185
141-78-6	176-177
142-68-7	228
142-84-7	320
144-49-0	54
151-18-8	111
151-56-4	74-75
275-51-4	401
280-57-9	307-308
289-80-5	145
289-95-2	146
290-37-9	147
302-01-2	464
327-54-8	248
334-88-3	21
350-03-8	331
350-40-3	385
352-93-2	194
353-36-6	71
359-11-5	45
359-13-7	60
362-43-6	422
362-76-5	428
363-72-4	247
367-11-3	254
367-23-7	251
367-64-6	294
368-48-9	258
371-40-4	273
372-18-9	255-256
372-19-0	274
372-38-3	252-253
372-47-4	207
372-48-5	208
383-63-1	151
383-66-4	217

392-56-3	246
400-59-9	298
406-34-8	80
430-67-1	72
431-47-0	97
455-19-6	360
456-48-4	327
459-57-4	328
460-39-9	108
461-24-5	157
462-06-6	266-267
462-94-2	245
463-49-0	101
463-51-4	50
493-66-8	350
503-29-7	127
503-30-0	117
506-68-3	12
506-77-4	13
513-35-9	223
530-48-3	430
536-90-3	349
540-36-3	257
540-61-4	62
540-67-0	133
542-76-7	103
545-06-2	43
547-63-7	231
550-33-4	405
551-62-2	249
554-12-1	178
556-61-6	58
556-64-9	58
557-01-7	148
557-40-4	300
563-46-2	224-225
563-80-4	229
563-83-7	186
585-48-8	429
589-93-5	345
590-18-1	163
591-22-0	346
591-49-1	352
592-84-7	232
593-53-3	27
593-54-4	38
593-88-4	136
594-09-2	142
594-11-6	164
594-39-8	239
598-56-1	199
598-98-1	309
603-34-9	437
613-97-8	395
616-12-6	304
616-38-6	124
616-39-7	240

616-47-7	156
620-08-6	285
623-49-4	154
623-53-0	181
624-64-6	165
624-78-2	140-141
624-89-5	135
624-91-9	30
625-27-4	305
625-54-7	236
625-55-8	179
625-58-1	78
625-80-9	316
626-23-3	380
626-60-8	205
626-61-9	206
626-67-5	311-312
630-08-0	39
637-92-3	314
676-59-5	92
677-41-8	109
687-48-9	235
693-86-7	218
693-89-0	295
694-52-0	209
694-85-9	286
715-91-3	408
753-90-2	61-62
763-29-1	306
765-43-5	219
765-47-9	353
819-06-7	166
819-46-5	167
918-02-5	321
926-64-7	168
931-20-4	301
996-35-0	241
1072-44-2	128
1122-54-9	332
1122-58-3	351
1122-61-8	210
1124-18-1	326
1190-79-0	183
1195-32-0	404
1310-58-3	448
1310-65-2	448
1310-73-2	449
1333-74-0	449-451
1528-30-9	296
1534-08-3	118
1605-72-7	14-15
1628-89-3	287-288
1630-77-9	49
1630-78-0	49
1634-04-4	237
1683-88-1	144
1712-69-2	403

1712-70-5	384
1733-57-9	431
2004-03-7	275
2016-57-1	417
2085-66-7	400
2129-89-7	427
2154-59-8	16
2168-84-5	125
2367-82-0	250
2450-71-7	105
2551-62-4	446
2730-67-8	110
2781-85-3	102
2868-37-3	221
2959-75-3	432
3796-23-4	259
3796-24-5	260
4096-20-2	420
4096-21-3	407
4143-41-3	81
4233-13-0	435
4252-88-4	433
4363-25-1	419
4458-31-5	356
4663-22-3	297
5500-21-0	153
5616-32-0	111
5627-05-4	397
5813-64-9	242
6569-51-3	7
6921-27-3	277
6921-28-4	284
6921-29-5	386
7087-68-5	381
7295-76-3	289
7439-90-9	466-467
7439-95-4	467
7440-37-1	1
7440-59-7	465-466
7440-61-1	474
7440-63-3	475
7440-70-2	439
7446-11-9	473
7637-07-2	4
7647-01-0	440
7664-39-3	443
7664-41-7	458-461
7727-37-9	470
7732-18-5	451-453
7782-39-0	441-442
7782-44-7	471-472
7783-06-4	454-456
7783-07-5	457
7783-54-2	445
7784-42-1	2
7789-20-0	442
7803-51-2	462-464

7803-62-5	465
10024-97-2	471
10034-85-2	447
10035-10-6	11
10076-31-0	357
10102-43-9	469
12007-71-5	8
12061-70-0	444
13463-40-6	202
13709-83-6	3
13952-84-6	200
13960-30-0	95
17222-37-6	44
18182-40-6	324
18283-93-7	9
19237-45-7	5
19624-22-7	10
19961-27-4	243
21351-79-1	441
21981-37-3	382
22025-87-2	424
23662-75-1	290
25493-13-4	359
29904-79-8	468
31023-92-4	416
32042-38-9	98
38103-96-7	180
53687-79-9	302
55186-75-9	402

Ar	1
AsH3	2
BF2H	3
BF3	4
B2H6	5
B3H5N3	6
B3H6N3	7
B4H8	8
B4H10	9
B5H9	10
BrH	11
CBrN	12
CClN	13
CCl2	14-15
CF2	16
CF4	17
CHF3	18
CHN	19-20
CH2F2	20
CH2N2	21
CH2O	21-22
CH2O2	23-25
CH3Br	25
CH3Cl	26-27
CH3F	27
CH3I	28
CH3NO2	29-30
CH4	30-31
CH4O	32-34
CH4S	34-35
CH5N	36-37
CH5P	38
CO	39
CO2	40-42
CS2	42
C2C13N	43
C2D6O	44
C2HF3	45
C2HF3O2	46-47
C2H2C1N	47
C2H2F2	48-49
C2H2O	50
C2H3C1O2	51
C2H3C13O	52
C2H3F	53
C2H3FO2	54
C2H3F3O	55
C2H3N	56-57
C2H3NS	58
C2H4	59
C2H4F2O	60
C2H4F3N	61-62
C2H4N2	62
C2H4O	63-65
C2H4O2	66-69
C2H5Br	69
C2H5Cl	70

C2H5F	71
C2H5F2N	72
C2H5I	73
C2H5N	74-75
C2H5NO	75-76
C2H5NO2	77
C2H5NO3	78
C2H6	79-80
C2H6FN	80
C2H6N2	81
C2H6O	82-86
C2H6OS	87
C2H6S	88-89
C2H7N	90-92
C2H7P	92
C2H8N2	93-94
C3D9N	95
C3H2N2	96
C3H3F3O2	97-98
C3H3N	99
C3H4	100-102
C3H4ClN	103
C3H5N	104-105
C3H6	106-108
C3H6F3N	108-110
C3H6N2	111
C3H6O	112-117
C3H6OS	118
C3H6O2	119-123
C3H6O3	124
C3H6S2	125
C3H7N	126-128
C3H7NO	129
C3H8	130-131
C3H8O	131-133
C3H8S	134-135
C3H9As	136
C3H9N	137-141
C3H9NO	141
C3H9P	142
C3H10N2	143
C4H4F4O2	144
C4H4N2	145-147
C4H4N2O	148
C4H4N2O2	149
C4H4O	150
C4H5F3O2	151
C4H5N	152-153
C4H5NO2	154
C4H5N3O	155
C4H6N2	156
C4H7F3O	157
C4H7N	158-159
C4H8	160-165
C4H8F3N	166-167
C4H8N2	168
C4H8O	169-172

C4H8O2	173-179
C4H8O2S	180
C4H8O3	181
C4H9N	182-183
C4H9NO	184-186
C4H10	187
C4H10N2	188
C4H10O	189-192
C4H10S	193-194
C4H11N	195-200
C4H12N2	201
C5FeO5	202
C5H3ClN4	203
C5H4ClN	204-206
C5H4FN	207-209
C5H4N2O2	210
C5H4N4O	211
C5H5N	212-213
C5H5N5	214
C5H5N5O	215
C5H6N2O2	216
C5H7F3O2	217
C5H8	218
C5H8O	219
C5H8O2	220-221
C5H9N	222
C5H10	223-225
C5H10O	225-229
C5H10O2	230-232
C5H11N	233-235
C5H11NO2	235
C5H12O	236-237
C5H13N	238-243
C5H14N2	244-245
C6F6	246
C6HF5	247
C6H2F4	248-250
C6H3F3	251-253
C6H4F2	254-257
C6H4F3N	258-260
C6H4N2	261-263
C6H5Cl	264-265
C6H5F	266-267
C6H5NO2	268
C6H6	269-270
C6H6ClN	271-272
C6H6FN	273-274
C6H6N4	275
C6H6O	276-277
C6H7N	278-284
C6H7NO	285-289
C6H7N5O	290
C6H8N2	291-293
C6H9F3O2	294
C6H10	295-297
C6H10F3NO	298
C6H10O	299-300

C6H11N	300
C6H11NO	301-302
C6H12	303-306
C6H12N2	307-308
C6H12O	308
C6H12O2	309
C6H13N	310-312
C6H14O	312-314
C6H14S	315-316
C6H15N	317-321
C6H16N2	322-323
C6H17NSi	324
C7H5ClO	325
C7H5D3	326
C7H5FO	327-328
C7H5N	329
C7H6O	330
C7H7NO	331-332
C7H8	333-336
C7H8O	337
C7H9N	338-346
C7H9NO	347-349
C7H10	350
C7H10N2	351
C7H12	352-353
C7H13N	354-355
C7H17N	355-357
C7H18N2	358
C8H4D6	359
C8H5F3O	360
C8H5NO	361
C8H8	362
C8H3O	363-364
C8H6O2	365-366
C8H10	367-373
C8H11N	374-376
C8H18S	377
C8H19N	378-382
C8H20N2	383
C9H9Cl	384
C9H9F	385
C9H9N	386
C9H10	387
C9H12	388-392
C9H12N2O6	393
C9H13N	394-395
C9H13N3O5	396
C9H14N2O6	397
C9H15N	398
C9H21N	399-400
C10H8	401
C10H9F3	402
C10H10Fe	403
C10H12	404
C10H12N4O4	405
C10H12O	406
C10H13N	407

C10H13N5	408
C10H13N5O4	409
C10H14	410-413
C10H14N2O5	414
C10H15N	415
C10H19N	416
C10H23N	417
C10H24N2	418
C11H13N	419
C11H15N	420
C12H11N	421
C12H16N2O6	422
C12H18	423
C12H19N	424
C12H27N	425
C13H10O	426
C13H13OP	427
C13H17N5O5	428
C13H21N	429
C14H12	430
C14H15OP	431
C15H17OP	432-433
C16H19OP	434-435
C17H21OP	436
C18H15N	437
C22H31OP	438
Ca	439
ClH	440
CsHO	441
D2	441-442
D2O	442
FH	443
FO	444
F3N	445
F6S	446
HI	447
HKO	448
HLiO	448
HNaO	449
H2	449-451
H2O	451-453
H2S	454-456
H2Se	457
H3N	458-461
H3P	462-464
H4N2	464
H4Si	465
He	465-466
Kr	466-467
Mg	467
Mg2	468
NO	469
N2	470
N2O	471
O2	471-472
O3S	473
U	474

Xe

475

Acetaldehyde (8CI9CI)	63-65	
Acetamide (8CI9CI)	75	
Acetamide, N-butyl-2,2,2-trifluoro- (8CI9CI)		298
Acetamide, N,N-dimethyl- (8CI9CI)	185	
Acetdimethylamide	185	
Acetene	59	
Acetic acid (8CI9CI)	66-67	
Acetic acid amide	75	
Acetic acid ethyl ester (8CI9CI)	176-177	
Acetic acid, chloro- (8CI9CI)	51	
Acetic acid, ethyl ester	176-177	
Acetic acid, fluoro- (8CI9CI)	54	
Acetic acid, methyl ester (8CI9CI)	120-121	
Acetic acid, propyl ester (8CI9CI)	230	
Acetic acid, thio-, S-methyl ester (8CI)	118	
Acetic acid, trifluoro- (8CI9CI)	46-47	
Acetic acid, trifluoro-, butyl ester (8CI9CI)		294
Acetic acid, trifluoro-, ethyl ester (8CI9CI)		151
Acetic acid, trifluoro-, methyl ester (8CI9CI)		97
Acetic acid, trifluoro-, propyl ester (8CI9CI)		217
Acetic acid, trifluoro-, 2-fluoroethyl ester (8CI)		144
Acetic aldehyde	63-65	
Acetic ether	176-177	
Acetidin	176-177	
Acetimidic acid	75	
Acetoacetone	220	
Acetone (8CI)	112-114	
Acetonitrile (8CI9CI)	56-57	
Acetonitrile, (dimethylamino)- (8CI9CI)		168
Acetonitrile, (methylamino)- (8CI9CI)		111
Acetonitrile, amino- (8CI9CI)	62	
Acetonitrile, chloro- (8CI9CI)	47	
Acetonitrile, trichloro- (8CI9CI)		43
Acetophenon	363	
Acetophenone (8CI)	363	
3-Acetopyridine	331	
Acetoxyethane	176-177	
1-Acetoxypropane	230	
Acetylacetone	220	
Acetylbenzene	363	
Acetylcyclopropane	219	
.beta.-Acetylpyridine		331
3-Acetylpyridine	331	
4-Acetylpyridine	332	
Acrylon	99	
Acrylonitrile (8CI)	99	
Ad	214	
Adenine (8CI)	214	
Adenine riboside	409	
Adenine, 9-cyclopentyl- (8CI)		408
Adeninimine	214	
.beta.-Adenosine	409	
.beta.-D-Adenosine	409	
Adenosine (8CI9CI)	409	
Aethylis	70	
Aethylis chloridum	70	
Aetznatron	449	

.beta.-Alaninenitrile	111	
Alcohol	82-83	
Alcohol anhydrous	82-83	
Alcojel	131-132	
Alcosolve 2	131-132	
Algrain	82-83	
Allene (8CI)	101	
Allyl ether (8CI)	300	
Allylamine (8CI)	126	
sym-Allylene	101	
Allylene	100	
Aminic acid	23-25	
3-Amino-1-(dimethylamino)propane	244	
3-Amino-1-propene	126	
6-Amino-1H-purine	214	
4-Amino-2(1H)-pyrimidinone	155	
4-Amino-2-hydroxypyrimidine	155	
1-Amino-2-methoxybenzene	347	
2-Amino-2-methylbutane	239	
1-Amino-2-methylpropane	196	
2-Amino-2-methylpropane	195	
1-Amino-3-(dimethylamino)propane	244	
1-Amino-3-dimethylaminopropane	244	
1-Amino-3-methoxybenzene	349	
1-Amino-3-methylbenzene	341-342	
6-Amino-3H-purine	214	
1-Amino-4-chlorobenzene	271	
1-Amino-4-methylbenzene	340	
2-Amino-6-hydroxypurine	215	
6-Amino-9-cyclopentylpurine	408	
6-Amino-9H-purine	214	
Aminoacetonitrile	62	
m-Aminoaniline	293	
p-Aminoaniline	292	
2-Aminoaniline	291	
3-Aminoaniline	293	
4-Aminoaniline	292	
m-Aminoanisole	349	
o-Aminoanisole	347	
p-Aminoanisole	348	
2-Aminoanisole	347	
3-Aminoanisole	349	
4-Aminoanisole	348	
Aminobenzene	278-280	
1-Aminobutane	197	
2-Aminobutane	200	
m-Aminochlorobenzene	272	
p-Aminochlorobenzene	271	
Aminocyclohexane	310-311	
1-Aminodecane	417	
Aminoethane	90	
1-Aminoethane	90	
2-Aminoethyl methyl ether	141	
.beta.-Aminoethylamine	94	
1-Aminoheptane	355	
Aminohexahydrobenzene	310-311	
1-Aminohexane	318	

2-Aminohypoxanthine	215	
2-Aminoisobutane	195	
Aminomethane	36-37	
1-Aminopentane	238	
Aminophen	278-280	
3-Aminophenylmethane	341-342	
1-Aminopropane	139-140	
2-Aminopropane	137	
3-Aminopropanenitrile	111	
3-Aminopropene	126	
.beta.-Aminopropionitrile	111	
Aminopropionitrile	111	
3-Aminopropionitrile	111	
3-Aminopropylene	126	
6-Aminopurine	214	
.alpha.-Aminotoluene	338	
.omega.-Aminotoluene	338	
m-Aminotoluene	341-342	
p-Aminotoluene	340	
3-Aminotoluene	341-342	
4-Aminotoluene	340	
Ammonia (8CI9CI)	458-461	
Ammonia gas	458-461	
n-Amylamine	238	
tert-Amylamine	239	
Amylamine	238	
Amylene	223	
Anaesthetic ether	189-190	
Anesthesia ether	189-190	
Anesthetic ether	189-190	
Anhydrol	82-83	
Anhydrous hydriodic acid	447	
Anhydrous hydrobromic acid	11	
Anhydrous hydrochloric acid	440	
Anhydrous hydrofluoric acid	443	
Aniline (8CI)	278-280	
Aniline, m-chloro- (8CI)	272	
Aniline, m-fluoro- (8CI)	274	
Aniline, o-tert-butyl-N,N-dimethyl- (8CI)	424	
Aniline, p-chloro- (8CI)	271	
Aniline, p-fluoro- (8CI)	273	
Aniline, N-ethyl- (8CI)	374	
Aniline, N-ethyl-N-methyl- (8CI)	395	
Aniline, N-methyl- (8CI)	339	
Aniline, N,N-diethyl- (8CI)	415	
Aniline, N,N-dimethyl- (8CI)	375-376	
Anilinobenzene	421	
Anilinoethane	374	
Anilinomethane	339	
Animal conine	245	
4-Anisaldehyde	366	
p-Anisaldehyde (8CI)	366	
p-Anisic aldehyde	366	
4-Anisidine	348	
m-Anisidine (8CI)	349	
o-Anisidine (8CI)	347	
p-Anisidine (8CI)	348	

Anisole (8CI)	337	
Anisole, p-isopropenyl- (8CI)		406
m-Anisylamine	349	
p-Anisylamine	348	
Anizol	337	
[6]Annulene	269-270	
Anodynol	70	
Anon	299	
Anone	299	
Antiren	188	
Antisal 1a	333-336	
Antisal 2b	443	
Anyvim	278-280	
Arcton 0	17	
Arcton 1	18	
Areginal	122-123	
Argon (8CI9CI)	1	
Argon-40	1	
Arsenic hydride	2	
Arsenic hydride (AsH ₃)		2
Arsenic trihydride	2	
Arseniuretted hydrogen		2
Arsenous hydride	2	
Arsine (8CI9CI)	2	
Arsine, trimethyl- (8CI9CI)		136
Artic	26-27	
Artificial Almond Oil		330
Ascarite	449	
Atomic helium	465-466	
Aubepine	366	
Avantin	131-132	
Avantine	131-132	
1-Aza-2,4-cyclopentadiene		152
Azabenzene	212-213	
1-Azabicyclo[2.2.2]octane (9CI)		354-355
1-Azabicyclo[3.3.3]undecane (9CI)		416
Azacyclobutane	127	
Azacyclohexane	233-234	
Azacyclopentane	182	
Azacyclopropane	74-75	
Azete, tetrahydro-	127	
Azetidine (8CI9CI)	127	
Azimethylene	21	
Azine	212-213	
Aziran	74-75	
Aziridine (9CI)	74-75	
Aziridine, 1-methyl- (8CI9CI)		128
Azole	152	
Azolidine	182	
trans-Azomethane	81	
Azomethane, (E)- (8CI)		81
Azulene (8CI9CI)	401	
AAN	62	
2-AB	200	
Ba 35846	281	
Benzaldehyde (8CI9CI)		330
Benzaldehyde FFC	330	

Benzaldehyde, m-fluoro- (8CI)	327	
Benzaldehyde, p-chloro- (8CI)	325	
Benzaldehyde, p-cyano- (8CI)	361	
Benzaldehyde, p-fluoro- (8CI)	328	
Benzaldehyde, 3-fluoro- (9CI)	327	
Benzaldehyde, 4-(trifluoromethyl)- (9CI)	360	
Benzaldehyde, 4-chloro- (9CI)	325	
Benzaldehyde, 4-fluoro- (9CI)	328	
Benzaldehyde, 4-methoxy- (9CI)	366	
Benzaldehyde, 4-methyl- (9CI)	364	
Benzenamine (9CI)	278-280	
Benzenamine, hexahydro- (8CI)	310-311	
Benzenamine, N-ethyl- (9CI)	374	
Benzenamine, N-ethyl-N-methyl- (9CI)	395	
Benzenamine, N-methyl- (9CI)	339	
Benzenamine, N-phenyl- (9CI)	421	
Benzenamine, N,N-diethyl- (9CI)	415	
Benzenamine, N,N-dimethyl- (9CI)	375-376	
Benzenamine, N,N-diphenyl- (9CI)	437	
Benzenamine, N,N,3-trimethyl- (9CI)	394	
Benzenamine, 2-(1,1-dimethylethyl)-N,N-dimethyl- (9CI)	424	
Benzenamine, 2-methoxy- (9CI)	347	
Benzenamine, 3-chloro- (9CI)	272	
Benzenamine, 3-fluoro- (9CI)	274	
Benzenamine, 3-methoxy- (9CI)	349	
Benzenamine, 3-methyl- (9CI)	341-342	
Benzenamine, 4-chloro- (9CI)	271	
Benzenamine, 4-fluoro- (9CI)	273	
Benzenamine, 4-methoxy- (9CI)	348	
Benzenamine, 4-methyl- (9CI)	340	
Benzene (8CI9CI)	269-270	
Benzene, (phenylamino)- (8CI)	421	
Benzene, (1-methylethenyl)- (9CI)	387	
Benzene, (1-methylethyl)- (9CI)	388-389	
Benzene, (1,1-dimethylethyl)- (9CI)	410-411	
Benzene, amino- (8CI)	278-280	
Benzene, benzoyl- (8CI)	426	
Benzene, butyl- (8CI9CI)	412-413	
Benzene, chloro- (8CI9CI)	264-265	
Benzene, cyano- (8CI)	329	
Benzene, ethenyl- (9CI)	362	
Benzene, ethyl- (8CI9CI)	369-371	
Benzene, fluoro- (8CI9CI)	266-267	
Benzene, hexafluoro- (8CI9CI)	246	
Benzene, hexamethyl- (8CI9CI)	423	
Benzene, m-difluoro- (8CI)	255-256	
Benzene, methoxy- (9CI)	337	
Benzene, methyl- (9CI)	333-336	
Benzene, methyl-d3- (8CI)	326	
Benzene, nitro- (8CI9CI)	268	
Benzene, o-difluoro- (8CI)	254	
Benzene, p-difluoro- (8CI)	257	
Benzene, pentafluoro- (8CI9CI)	247	
Benzene, propyl- (8CI9CI)	390-391	
Benzene, tert-butyl- (8CI)	410-411	
Benzene, 1-(dimethylamino)-3-methyl- (8CI)	394	
Benzene, 1-(1-methylethenyl)-4-(trifluoromethyl)- (9CI)	402	

Benzene, 1-chloro-4-(1-methylethenyl)- (9CI)	384	
Benzene, 1-fluoro-4-(1-methylethenyl)- (9CI)	385	
Benzene, 1-methoxy-4-(1-methylethenyl)- (9CI)		406
Benzene, 1-methyl-4-(1-methylethenyl)- (9CI)	404	
Benzene, 1,1'-ethenylidenebis- (9CI)	430	
Benzene, 1,2-difluoro- (9CI)	254	
Benzene, 1,2-dimethyl- (9CI)	367-368	
Benzene, 1,2,3,4-tetrafluoro- (8CI9CI)	249	
Benzene, 1,2,3,5-tetrafluoro- (8CI9CI)	250	
Benzene, 1,2,4-trifluoro- (8CI9CI)	251	
Benzene, 1,2,4,5-tetrafluoro- (8CI9CI)	248	
Benzene, 1,3-difluoro- (9CI)	255-256	
Benzene, 1,3-dimethyl- (9CI)	373	
Benzene, 1,3,5-trifluoro- (8CI9CI)	252-253	
Benzene, 1,3,5-trimethyl- (9CI)	392	
Benzene, 1,4-(dimethyl-d3)-	359	
Benzene, 1,4-difluoro- (9CI)	257	
Benzene, 1,4-dimethyl- (9CI)	372	
Benzenecarbonal	330	
Benzenecarboxaldehyde	330	
m-Benzenediamine	293	
o-Benzenediamine	291	
p-Benzenediamine	292	
1,2-Benzenediamine (9CI)	291	
1,3-Benzenediamine (9CI)	293	
1,4-Benzenediamine (9CI)	292	
Benzenemethanamine (9CI)	338	
Benzenol	276	
Benzofur D	292	
Benzoic acid nitrile	329	
Benzoic acid, methyl ester (8CI9CI)	365	
Benzoic aldehyde	330	
Benzol	269-270	
Benzole	269-270	
Benzonitrile (8CI9CI)	329	
Benzonitrile, p-formyl-	361	
Benzonitrile, 4-formyl- (9CI)	361	
Benzophenone (8CI)	426	
Benzoquinuclidine	419	
Benzoyl methide	363	
Benzoylbenzene	426	
Benzylamine (8CI)	338	
Bicarburretted hydrogen	59	
Bicyclo[2.2.1]hept-2-ene (9CI)	350	
Bicyclo[2.2.1]heptene	350	
Bicyclo[2.2.2]-1,4-diazaoctane	307-308	
Bicyclo[5.3.0]decapentaene	401	
Bilorin	23-25	
Bimethyl	79-80	
Bis(.eta.-cyclopentadienyl)iron	403	
Bis(cyclopentadienyl)iron	403	
1,4-Bis(dimethylamino)butane	383	
1,2-Bis(dimethylamino)ethane	322	
1,6-Bis(dimethylamino)hexane	418	
1,3-Bis(dimethylamino)propane	358	
Bis(methyl-d3) ether	44	
Bis(1-methylpropyl)amine	380	

N,N-Bis(2-methylpropyl)amine	378	
Bis[(dimethylamino)methyl]methane		358
Blausäure	19-20	
Blood-coagulation factor IV	439	
Blue Oil	278-280	
Boniton	409	
Borane, difluoro- (8CI9CI)	3	
Borane, trifluoro- (9CI)	4	
Borazine (8CI9CI)	7	
.beta.-Borazinyl radical	6	
Borazole	7	
Borazyne, cyclic trimer	7	
Boron fluoride	4	
Boron fluoride (BF ₃) (8CI)	4	
Boron fluoride hydride (BHF ₂)		3
Boron hydride (B ₂ H ₆)	5	
Boron hydride (B ₄ H ₁₀)	9	
Boron trifluoride	4	
Bromic ether	69	
Bromine cyanide	12	
Bromine cyanide (BrCN)	12	
Bromocyan	12	
Bromocyanide	12	
Bromocyanide (BrCN)	12	
Bromocyanogen	12	
Bromoethane	69	
Bromomethane	25	
Butafume	200	
Butal	172	
Butaldehyde	172	
n-Butan-1-ol	191	
n-Butanal	172	
Butanal (9CI)	172	
Butanaldehyde	172	
1-Butanamine (9CI)	197	
2-Butanamine (9CI)	200	
2-Butanamine, N-(1-methylpropyl)- (9CI)		380
1-Butanamine, N-butyl- (9CI)	379	
1-Butanamine, N,N-dibutyl- (9CI)	425	
2-Butanamine, 2-methyl- (9CI)	239	
1-Butanamine, 4,4,4-trifluoro- (9CI)		167
n-Butane	187	
Butane .alpha.,.delta.-oxide	171	
Butane (8CI9CI)	187	
Butane, 1-cyano-	222	
Butane, 1,4-epoxy-	171	
1,4-Butanediamine (8CI9CI)	201	
1,4-Butanediamine, N,N,N',N'-tetramethyl- (8CI9CI)		383
Butanenitrile (9CI)	159	
tert-Butanethiol	193	
n-Butanol	191	
t-Butanol	192	
tert-Butanol	192	
Butanol	191	
1-Butanol (9CI)	191	
Butanone	170	
2-Butanone (8CI9CI)	170	

2-Butanone, 3-methyl- (8CI9CI)	229	
2-Butanone, 3,3-dimethyl- (8CI9CI)		308
.alpha.-Butene	160	
(E)-2-Butene	165	
(Z)-2-Butene	163	
cis-Butene	163	
cis-2-Butene	163	
iso-Butene	161-162	
trans-Butene	165	
trans-2-Butene	165	
2-trans-Butene	165	
1-Butene (8CI9CI)		160
Butene-1	160	
2-Butene, (E)- (8CI9CI)	165	
2-Butene, (Z)- (8CI9CI)	163	
1-Butene, 2-ethyl-	303	
1-Butene, 2-methyl- (8CI9CI)		224-225
2-Butene, 2-methyl- (8CI9CI)	223	
n-Butyl alcohol	191	
tert-Butyl alcohol (8CI)		192
Butyl alcohol (8CI)	191	
n-Butyl aldehyde	172	
Butyl aldehyde	172	
Butyl cyanide	222	
tert-Butyl ethyl ether		314
n-Butyl formate	232	
Butyl formate	232	
Butyl hydroxide	191	
tert-Butyl mercaptan	193	
tert-Butyl methyl ether		237
tert-Butyl methyl ketone		308
tert-Butyl sulfide (8CI)		377
Butyl trifluoroacetate		294
i-Butylamine	196	
iso-Butylamine	196	
n-Butylamine	197	
t-Butylamine	195	
2-Butylamine	200	
sec-Butylamine (8CI)		200
tert-Butylamine (8CI)		195
Butylamine (8CI)	197	
Butylamine, 4,4,4-trifluoro- (8CI)		167
n-Butylbenzene	412-413	
t-Butylbenzene	410-411	
tert-Butylbenzene	410-411	
Butylbenzene	412-413	
tert-Butyldimethylamine		321
.alpha.-Butylene	160	
.gamma.-Butylene	161-162	
cis-Butylene	163	
1-Butylene	160	
Butylenediamine	201	
1,4-Butylenediamine		201
Butylenimine	182	
t-Butylmercaptan	193	
tert-Butylthiol	193	
N-Butyltrifluoroacetamide		298

Butyral	172	
iso-Butyraldehyde		169
n-Butyraldehyde	172	
Butyraldehyde (8CI)		172
Butyric aldehyde	172	
n-Butyronitrile	159	
Butyronitrile (8CI)		159
Butyrylaldehyde	172	
Butyrylonitrile	159	
BAPN	111	
BASF Ursol D	292	
BASF 238	184	
BFV	21-22	
C.I. Azoic Coupling Component	107	340
C.I. Developer 11	293	
C.I. Developer 13	292	
C.I. Natural White 1	215	
C.I. Oxidation Base 10		292
C.I. Oxidation Base 16		291
C.I. 37107	340	
C.I. 75170	215	
C.I. 76000	278-280	
C.I. 76010	291	
C.I. 76060	292	
Cadaverine	245	
Calcium (8CI9CI)	439	
Campilit	12	
Carbacryl	99	
Carbamic acid, dimethyl-, ethyl ester (8CI9CI)		235
Carbinamine	36-37	
Carbinol	32-34	
Carbolic acid	276	
Carbomethene	50	
Carbon bisulfide	42	
Carbon bisulphide	42	
Carbon chloride (CCl ₂)		14-15
Carbon difluoride	16	
Carbon dioxide (8CI9CI)		40-42
Carbon disulfide (8CI9CI)		42
Carbon disulphide	42	
Carbon fluoride (CF ₂)	16	
Carbon fluoride (CF ₄)	17	
Carbon fluoride hydride (CF ₂ H ₂)		20
Carbon hydride nitride (CHN)		19-20
Carbon monoxide (8CI9CI)	39	
Carbon oxide (CO)	39	
Carbon oxide (CO ₂)	40-42	
Carbon sulfide (CS ₂)	42	
Carbon tetrafluoride (8CI)		17
Carbon trifluoride	18	
Carbonic acid gas	40-42	
Carbonic acid, dimethyl ester (8CI9CI)		124
Carbonic acid, ethyl methyl ester (8CI9CI)		181
Carbonic anhydride	40-42	
Carbonocyanidic acid, ethyl ester (9CI)		154
Carbonythioic acid, S-ethyl O-methyl ester (9CI)		180
Carboxyethane	119	

Caustic potash	448	
Caustic soda	449	
Cesium hydroxide (8CI9CI)		441
Chelen	70	
Chloracetic acid	51	
Chloracetonitrile		47
Chlorene	70	
Chlorethyl	70	
Chloridum	70	
Chlorine cyanide	13	
Chlorine cyanide (ClCN)		13
p-Chloro-.alpha.-methylstyrene		384
4-Chloro-.alpha.-methylstyrene		384
4-Chloro-1-aminobenzene	271	
1-Chloro-2-cyanoethane	103	
1-Chloro-4-isoproprenylbenzene		384
6-Chloro-9H-purine	203	
.alpha.-Chloroacetic acid		51
Chloroacetic acid	51	
.alpha.-Chloroacetoneitrile		47
Chloroacetoneitrile	47	
2-Chloroacetoneitrile	47	
m-Chloroaniline	272	
p-Chloroaniline	271	
3-Chloroaniline	272	
4-Chloroaniline	271	
p-Chlorobenzaldehyde	325	
4-Chlorobenzaldehyde	325	
3-Chlorobenzenamine	272	
4-Chlorobenzenamine	271	
Chlorobenzene	264-265	
Chlorocyan	13	
Chlorocyanide	13	
Chlorocyanide (ClCN)		13
Chlorocyanogen	13	
Chloroethane	70	
Chloroethanoic acid	51	
Chlorohydric acid	440	
Chloromethane	26-27	
Chloromethyl cyanide	47	
2-(p-Chlorophenyl)propene		384
2-(4-Chlorophenyl)propene		384
m-Chlorophenylamine	272	
p-Chlorophenylamine	271	
3-Chloropropanenitrile		103
3-Chloropropanonitrile		103
.beta.-Chloropropionitrile		103
3-Chloropropionitrile		103
6-Chloropurine	203	
.alpha.-Chloropyridine		204
.gamma.-Chloropyridine		206
m-Chloropyridine	205	
o-Chloropyridine	204	
2-Chloropyridine	204	
3-Chloropyridine	205	
4-Chloropyridine	206	
Chloryl	70	

Chloryl Anesthetic	70	
Cinnamene	362	
Cloretilo	70	
Clorius	365	
Coal naphtha	269-270	
Collo-Bueglatt	23-25	
Collo-Didax	23-25	
Collo-Grillrein	449	
Collo-Tapetta	449	
Combi-Schutz	131-132	
Cratagine	366	
Cumene (8CI)	388-389	
Cumol	388-389	
Curafume	25	
1-Cyano-1-methylethane		158
Cyanoacetonitrile	96	
p-Cyanobenzaldehyde	361	
4-Cyanobenzaldehyde	361	
Cyanobenzene	329	
Cyanobromide	12	
1-Cyanobutane	222	
Cyanocyclopropane		153
Cyanoethane	104	
.beta.-Cyanoethylamine		111
2-Cyanoethylamine	111	
Cyanoethylene	99	
Cyanogen bromide (BrCN)	12	
Cyanogen bromide (8CI9CI)		12
Cyanogen chloride (ClCN)	13	
Cyanogen chloride (8CI9CI)		13
Cyanogen monobromide	12	
Cyanomethane	56-57	
N-(Cyanomethyl)dimethylamine		168
Cyanomethylamine	62	
1-Cyanopropane	159	
2-Cyanopropane	158	
.gamma.-Cyanopyridine		261
2-Cyanopyridine	263	
3-Cyanopyridine	262	
4-Cyanopyridine	261	
Cyanotrichloromethane	43	
Cyclohexanamine (9CI)	310-311	
Cyclohexanone (8CI9CI)	299	
Cyclohexatriene	269-270	
Cyclohexene, 1-methyl- (8CI9CI)		352
Cyclohexylamine (8CI)	310-311	
Cyclooxabutane	117	
Cyclopentacycloheptene	401	
Cyclopentane, methylene- (8CI9CI)		296
Cyclopentene, 1-methyl- (8CI9CI)	295	
Cyclopentene, 1,2-dimethyl- (8CI9CI)		353
Cyclopentimine	233-234	
9-Cyclopentyladenine	408	
Cyclopropane (8CI9CI)	106	
Cyclopropane, (1-methylethenyl)- (9CI)		297
Cyclopropane, ethenyl- (9CI)	218	
Cyclopropane, methyl- (8CI9CI)		164

Cyclopropane, vinyl- (8CI)	218	
Cyclopropanecarbonitrile (8CI9CI)	153	
Cyclopropanecarboxylic acid, methyl ester (8CI9CI)		221
Cyclopropanenitrile	153	
Cyclopropene (8CI9CI)	102	
Cyclopropyl cyanide	153	
Cyclopropyl methyl ketone	219	
2-Cyclopropyl-1-propene	297	
Cyclopropylethylene	218	
Cyclopropylnitrile	153	
2-Cyclopropylpropene	297	
Cyclotetramethylene oxide	171	
Cyd	396	
Cymonic acid	54	
Cypentil	233-234	
Cyt	155	
Cytidine (9CI)	396	
Cytosine (8CI)	155	
Cytosine riboside	396	
Cytosinimine	155	
p-CA	271	
CCS 203	191	
CP 28	246	
CP 32	212-213	
D 33LV	307-308	
Dabco	307-308	
Dabco 33LV	307-308	
Decahydrotetraborane	9	
Decanamine	417	
1-Decanamine (9CI)	417	
n-Decylamine	417	
1-Decylamine	417	
Decylamine (8CI)	417	
Dehydro-p-cymene	404	
Demsodrox	87	
Denatured ethanol	82-83	
Deoxythymidine	414	
2'-Deoxythymidine	414	
Deuterium (8CI9CI)	441-442	
Deuterium oxide	442	
Deuterium oxide (D20)	442	
Developer C	293	
Developer H	293	
Developer M	293	
Developer PF	292	
Devoton	120-121	
Dew Pearl	215	
Di-.pi.-cyclopentadienyl iron	403	
Di-n-butylamine	379	
Di-n-propyl ether	313	
Di-n-propylamine	320	
Di-sec-butylamine (8CI)	380	
Di-t-Butylamine	382	
Di-tert-butyl sulfide	377	
Di-tert-butylamine (8CI)	382	
2,6-Di-tert-butylpyridine	429	
Di-2-propenylamine	300	

Di-2-propynylamine (8CI)	284	
Di-2,4-cyclopentadien-1-yliron		403
Diacetylmethane	220	
Diallyl ether	300	
Diallylamine (8CI)	300	
1,6-Diamino-n-hexane	323	
m-Diaminobenzene	293	
o-Diaminobenzene	291	
p-Diaminobenzene	292	
1,2-Diaminobenzene	291	
1,3-Diaminobenzene	293	
1,4-Diaminobenzene	292	
1,4-Diaminobutane	201	
1,2-Diaminoethane	94	
1,6-Diaminohexane	323	
1,5-Diaminopentane	245	
1,3-Diaminopropane	143	
1,2-Diazabenzene	145	
1,3-Diazabenzene	146	
1,4-Diazabenzene	147	
1,4-Diazabicyclo[2.2.2]octane (8CI9CI)		307-308
1,4-Diazacyclohexane	188	
Diazene, dimethyl-, (E)- (9CI)	81	
m-Diazine	146	
p-Diazine	147	
1,2-Diazine	145	
1,3-Diazine	146	
1,4-Diazine	147	
Diazirine	21	
Diazomethane	21	
Diborane	5	
Diborane (B ₂ H ₆)	5	
Diborane(6) (8CI9CI)	5	
Diboron hexahydride	5	
Dibutylamine (8CI)	379	
Dichlorocarbene	14-15	
Dichloromethylene	14-15	
Dicyanomethane	96	
Dicyclopentadienyliron		403
Dideuterium oxide	442	
Dietary calcium	439	
Diethyl	187	
Diethyl ether	189-190	
Diethyl ketone	225	
Diethyl oxide	189-190	
Diethyl sulfide	194	
N,N-Diethylamine	198	
Diethylamine (8CI)	198	
Diethylamine, N-methyl- (8CI)		240
Diethylamine, 1-methyl- (8CI)		243
N,N-Diethylaminobenzene	415	
Diethylaniline	415	
N,N-Diethylaniline	415	
Diethylene dioxide	175	
1,4-Diethylene dioxide	175	
Diethylene ether	175	
Diethylene imidoxide	184	

Diethylene oxide	175	
Diethylene oximide	184	
Diethylenediamine	188	
Diethyleneimine	188	
Diethylenimide oxide	184	
Diethylmethylaniline	240	
N,N-Diethylmethylaniline	240	
Diethylphenylamine	415	
Diethylpropylamine	356	
Diethylthioether	194	
m-Difluorobenzene	255-256	
o-Difluorobenzene	254	
p-Difluorobenzene	257	
1,2-Difluorobenzene	254	
1,3-Difluorobenzene	255-256	
1,4-Difluorobenzene	257	
Difluoroborane	3	
Difluorocarbene	16	
2,2-Difluoroethanol	60	
cis-1,2-Difluoroethene	49	
trans-1,2-Difluoroethene	49	
1,1-Difluoroethene	48	
2,2-Difluoroethylamine	72	
(E)-1,2-Difluoroethylene	49	
cis-Difluoroethylene	49	
cis-1,2-Difluoroethylene	49	
trans-Difluoroethylene	49	
trans-1,2-Difluoroethylene	49	
1,1-Difluoroethylene	48	
Difluoromethane	20	
Difluoromethylene	16	
Difluoromethylene radical	16	
Difluorocarbene	16	
3,4-Dihydro-2H-1,4-ethanoquinoline	419	
1,6-Dihydro-6-iminopurine	214	
3,6-Dihydro-6-iminopurine	214	
Dihydrogen monosulfide	454-456	
Dihydrogen selenide	457	
Dihydrogen sulfide	454-456	
Dihydrooxirene	65	
Dihydrouridine	397	
5,6-Dihydrouridine	397	
2,4-Dihydroxy-5-methylpyrimidine	216	
2,4-Dihydroxypyrimidine	149	
Diisobutylamine (8CI)	378	
Diisopropyl ether	312	
Diisopropyl oxide	312	
Diisopropyl sulfide	316	
Diisopropylamine (8CI)	317	
Diisopropylethylamine	381	
N,N-Diisopropylethylamine	381	
Dilute hydrochloric acid	440	
Dimazin	93	
Dimazine	93	
Dimethyl	79-80	
Dimethyl carbonate	124	
Dimethyl ether	84-86	

Dimethyl ketone	112-114	
Dimethyl monosulfide	89	
Dimethyl oxide	84-86	
Dimethyl sulfide	89	
Dimethyl sulfoxide	87	
Dimethyl sulphoxide	87	
Dimethyl thioether	89	
N,N-Dimethyl-m-methylaniline	394	
Dimethyl-m-toluidine	394	
N,N-Dimethyl-m-toluidine	394	
Dimethyl-tert-butylamine	321	
N,N-Dimethyl-tert-butylamine	321	
N,N-Dimethyl-N-(3-aminopropyl)amine		244
1,2-Dimethyl-1-cyclopentene	353	
N,N-Dimethyl-1,3-diaminopropane		244
N,N-Dimethyl-1,3-propanediamine		244
N,N-Dimethyl-1,3-propylenediamine		244
3,3-Dimethyl-2-butanone	308	
2,2-Dimethyl-3-butanone	308	
N,N-Dimethyl-3-methylaniline	394	
2,4-Dimethyl-3-thiapentane	316	
N,N-Dimethyl-4-aminopyridine	351	
N,N-Dimethyl-4-pyridinamine	351	
Dimethylacetamide	185	
N,N-Dimethylacetamide	185	
Dimethylacetone	225	
Dimethylamide acetate	185	
Dimethylamine (8CI)	91-92	
4-Dimethylaminepyridine	351	
3-(Dimethylamino)-1-propanamine		244
3-(Dimethylamino)-1-propylamine		244
(Dimethylamino)acetonitrile	168	
(Dimethylamino)benzene	375-376	
[(Dimethylamino)methyl]trimethylsilane		324
3-(Dimethylamino)propylamine	244	
.gamma.-(Dimethylamino)pyridine		351
4-(Dimethylamino)pyridine	351	
1-Dimethylamino-3-aminopropane		244
2-Dimethylaminopropane	241	
.gamma.-Dimethylaminopropylamine		244
Dimethylaniline	375-376	
N,N-Dimethylaniline	375-376	
m-Dimethylbenzene	373	
o-Dimethylbenzene	367-368	
p-Dimethylbenzene	372	
1,2-Dimethylbenzene	367-368	
1,3-Dimethylbenzene	373	
1,4-Dimethylbenzene	372	
2,2-Dimethylbutanone	308	
3,3-Dimethylbutanone	308	
Dimethylcarbinol	131-132	
1,2-Dimethylcyclopentene	353	
(E)-Dimethyldiazene	81	
Dimethylene oxide	65	
Dimethylenediamine	94	
Dimethylenemethane	101	
Dimethylenimine	74-75	

1,1-Dimethylethanethiol	193	
1,1-Dimethylethanol	192	
Dimethylethoxyformamide	235	
N,N-Dimethylethylamine	199	
1,1-Dimethylethylamine	195	
Dimethylethylbenzene	410-411	
cis-1,2-Dimethylethylene	163	
trans-1,2-Dimethylethylene	165	
1,1-Dimethylethylene	161-162	
Dimethylformaldehyde	112-114	
Dimethylformamide	129	
N,N-Dimethylformamide	129	
u-Dimethylhydrazine	93	
unsym-Dimethylhydrazine	93	
N,N-Dimethylhydrazine	93	
1,1-Dimethylhydrazine	93	
Dimethylisopropylamine	241	
N,N-Dimethylisopropylamine	241	
Dimethylmethane	130-131	
N,N-Dimethylneopentylamine	357	
Dimethylphenylamine	375-376	
Dimethylphosphine	92	
.alpha.,.alpha.-Dimethylpropylamine	239	239
1,1-Dimethylpropylamine	239	
2,2-Dimethylpropylamine	242	
N,N-Dimethylpropylenediamine	244	
.alpha.,.alpha.'-Dimethylpyridine		344
.alpha.,.gamma.-Dimethylpyridine	343	
2,4-Dimethylpyridine	343	
2,5-Dimethylpyridine	345	
2,6-Dimethylpyridine	344	
3,5-Dimethylpyridine	346	
.alpha.,p-Dimethylstyrene	404	
.alpha.,4-Dimethylstyrene	404	
p,.alpha.-Dimethylstyrene	404	
1,1'-Dimethyltriethylamine	381	
N,N-Dimethyltrimethylenediamine	244	
Dimethyl[2-(dimethylamino)ethyl]amine		322
Dimexide	87	
Dinitrogen monoxide	471	
Dinitrogen oxide	471	
1,4-Dioxacyclohexane	175	
p-Dioxan	175	
Dioxan	175	
1,4-Dioxan	175	
Dioxane	175	
p-Dioxane (8CI)	175	
1,4-Dioxane (9CI)	175	
1,4-Dioxin, tetrahydro-	175	
2,4-Dioxopentane	220	
2,4-Dioxypyrimidine	149	
Dioxyethylene ether	175	
Diphenyl ketone	426	
Diphenyl(ethyl)phosphine oxide		431
Diphenylamine (8CI)	421	
N,N-Diphenylaniline	437	
1,1-Diphenylethene	430	

.alpha.,.alpha.-Diphenylethylene	430
unsym.-Diphenylethylene	430
Diphenylethylene	430
1,1-Diphenylethylene	430
Diphenylmethanone	426
Diphenylmethylphosphine oxide	427
Dipirartril-tropico	87
Diplogen	441-442
Dipropargyl ether	277
Dipropargyl oxide	277
Dipropargylamine	284
Dipropyl ether	313
Dipropyl oxide	313
Dipropyl sulfide	315
Dipropyl thioether	315
n-Dipropylamine	320
Dipropylamine (8CI)	320
Direct Brown BR	293
Direct Brown GG	293
Dispermine	188
Distilled water	451-453
Dithiocarbonic anhydride	42
Divinylene oxide	150
Divinylenimine	152
Dolicur	87
Dromisol	87
Dry ice	40-42
Dublofix	70
Durafur Black R	292
Durasorb	87
DDNU	430
DEK	225
DFA	421
DMA	185
DMF	129
DMF (amide)	129
DMFA	129
DMS	89
DMS 70	87
DMS 90	87
DMSO	87
DPA	421
DT	414
DThyd	414
Elayl	59
Elegas	446
Embafume	25
3,6-Endomethylenecyclohexene	350
Epoxyethane	65
1,2-Epoxyethane	65
1,3-Epoxypropane	117
Eraverm	188
Essence of Mirbane	268
Essence of Myrbane	268
ps-Estragole	406
Ethanal	63-65
Ethanamide	75

Ethanamine (9CI)	90	
Ethanamine, N-ethyl- (9CI)	198	
Ethanamine, N-ethyl-N-methyl- (9CI)	240	
Ethanamine, N-ethyl-N-propyl-	356	
Ethanamine, N-ethylidene- (9CI)	183	
Ethanamine, N-methyl- (9CI)	140-141	
Ethanamine, N,N-diethyl- (9CI)	319	
Ethanamine, N,N-dimethyl- (9CI)	199	
Ethanamine, 2-fluoro- (9CI)	80	
Ethanamine, 2-methoxy- (9CI)	141	
Ethanamine, 2,2-difluoro- (9CI)	72	
Ethanamine, 2,2,2-trifluoro- (9CI)	61-62	
Ethanamine, 2,2,2-trifluoro-N-methyl- (9CI)	110	
Ethanamine, 2,2,2-trifluoro-N,N-dimethyl- (9CI)	166	
Ethane (8CI9CI)	79-80	
Ethane(dithioic)acid, methyl ester	125	
Ethane, (methylthio)- (9CI)	135	
Ethane, bromo- (8CI9CI)	69	
Ethane, chloro- (8CI9CI)	70	
Ethane, fluoro- (8CI9CI)	71	
Ethane, iodo- (8CI9CI)	73	
Ethane, methoxy- (9CI)	133	
Ethane, nitro- (8CI9CI)	77	
Ethane, 1,1'-oxybis- (9CI)	189-190	
Ethane, 1,1'-thiobis- (9CI)	194	
Ethane, 2-ethoxy-1,1,1-trifluoro- (9CI)	157	
Ethancarboxylic acid	119	
1,2-Ethanediamine (9CI)	94	
1,2-Ethanediamine, N,N,N',N'-tetramethyl- (9CI)	322	
Ethanenitrile	56-57	
Ethanethioic acid, S-methyl ester (9CI)	118	
Ethanethiol (8CI9CI)	88	
Ethanoic acid	66-67	
Ethanol (9CI)	82-83	
Ethanol, 2,2-difluoro- (8CI9CI)	60	
Ethanol, 2,2,2-trichloro- (8CI9CI)	52	
Ethanol, 2,2,2-trifluoro- (8CI9CI)	55	
Ethanol, 2,2,2-trifluoro-, formate (9CI)	98	
Ethanone, 1-(3-pyridinyl)- (9CI)	331	
Ethanone, 1-(4-pyridinyl)- (9CI)	332	
Ethanone, 1-cyclopropyl- (9CI)	219	
Ethanone, 1-phenyl- (9CI)	363	
2H-1,4-Ethanoquinoline, 3,4-dihydro- (8CI9CI)	419	
Ethene (9CI)	59	
Ethene oxide	65	
Ethene, cyclopropyl-	218	
Ethene, fluoro- (9CI)	53	
Ethene, trifluoro- (9CI)	45	
Ethene, 1,1-difluoro- (9CI)	48	
Ethene, 1,2-difluoro-, (E)- (9CI)	49	
Ethene, 1,2-difluoro-, (Z)- (9CI)	49	
Ethenone (9CI)	50	
Ether	189-190	
Ether chloratus	70	
Ether cyanatus	104	
Ether hydrochloric	70	
Ether muriatic	70	

Ether, ethyl isopropyl (8CI)	236	
Ether, ethyl methyl (8CI)	133	
Ether, ethyl 2,2,2-trifluoroethyl (8CI)		157
Ether, tert-butyl ethyl (8CI)	314	
Ether, tert-butyl methyl (8CI)	237	
Ethiridine	74-75	
Ethoxyethane	189-190	
Ethyl acetate	176-177	
Ethyl acetic ester	176-177	
Ethyl alcohol (8CI)	82-83	
Ethyl aldehyde	63-65	
Ethyl bromide	69	
Ethyl carbinol	132-133	
Ethyl chloride	70	
Ethyl cyanide	104	
Ethyl cyanoformate	154	
Ethyl dimethylcarbamate	235	
Ethyl ethanoate	176-177	
Ethyl ether (8CI)	189-190	
Ethyl fluoride	71	
Ethyl formate	122-123	
Ethyl hydrate	82-83	
Ethyl hydride	79-80	
Ethyl hydrosulfide	88	
Ethyl hydroxide	82-83	
Ethyl iodide	73	
Ethyl isopropyl ether	236	
Ethyl ketone	225	
Ethyl mercaptan	88	
Ethyl methanoate	122-123	
Ethyl methyl carbonate	181	
Ethyl methyl ether	133	
Ethyl methyl ketone	170	
Ethyl methyl sulfide	135	
Ethyl monosulfide	194	
Ethyl nitrate	78	
Ethyl nitrile	56-57	
Ethyl sulphydrate	88	
Ethyl sulfide (8CI)	194	
Ethyl tert-butyl ether	314	
Ethyl tert-butyl oxide	314	
Ethyl thioalcohol	88	
Ethyl thioether	194	
Ethyl trifluoroacetate	151	
Ethyl N,N-dimethylcarbamate	235	
S-Ethyl O-methyl thiocarbonate		180
N-Ethyl-N-methylaniline	395	
N-Ethyl-N-phenylamine	374	
N-Ethyl-N,N-diisopropylamine	381	
Ethylamine (8CI)	90	
Ethylamine, N-ethylidene- (8CI)	183	
Ethylamine, N-methyl- (8CI)	140-141	
Ethylamine, N,N-dimethyl- (8CI)	199	
Ethylamine, N,N,1-trimethyl- (8CI)		241
Ethylamine, N,N,1,1-tetramethyl- (8CI)		321
Ethylamine, 2-fluoro- (8CI)	80	
Ethylamine, 2-methoxy- (8CI)	141	

Ethylamine, 2,2-difluoro- (8CI)	72	
Ethylamine, 2,2,2-trifluoro- (8CI)	61-62	
Ethylamine, 2,2,2-trifluoro-N-methyl- (8CI)	110	
Ethylamine, 2,2,2-trifluoro-N,N-dimethyl- (8CI)	166	
N-Ethylaminobenzene	374	
Ethylaniline	374	
N-Ethylaniline	374	
N-Ethylbenzenamine	374	
Ethylbenzene	369-371	
Ethylbenzol	369-371	
Ethyl-diisopropylamine	381	
N-Ethyl-diisopropylamine	381	
Ethyl-diphenylphosphine oxide	431	
Ethylene (8CI)	59	
Ethylene oxide (8CI)	65	
Ethylene trifluoride	45	
Ethylene, cyclopropyl-	218	
Ethylene, fluoro- (8CI)	53	
Ethylene, trifluoro- (8CI)	45	
Ethylene, 1,1-difluoro- (8CI)	48	
Ethylene, 1,1-diphenyl- (8CI)	430	
Ethylene, 1,2-difluoro-, (E)- (8CI)	49	
Ethylene, 1,2-difluoro-, (Z)- (8CI)	49	
1,2-Ethylenediamine	94	
Ethylenediamine (8CI)	94	
Ethylenediamine, N,N,N',N'-tetramethyl- (8CI)	322	
Ethyleneimine	74-75	
N,N'-endo-Ethylenepiperazine	307-308	
1,4-Ethylenepiperazine	307-308	
1,4-Ethylenepiperidine	354-355	
Ethylenimine (8CI)	74-75	
Ethylethylene	160	
Ethylformic acid	119	
Ethyllic acid	66-67	
Ethylideneethylamine	183	
N-Ethylideneethylamine	183	
Ethylisopropylamine	243	
N-Ethylisopropylamine	243	
Ethylmethylaniline	140-141	
Ethylphenylaniline	374	
Ethylthioethane	194	
Exhaust gas	39	
EB	369-371	
EI	74-75	
ETO	65	
Fannoform	21-22	
Fast Orange GC Base	272	
Ferrocene (8CI9CI)	403	
Ferrotsen	403	
Flue gas	39	
Fluorhydric acid	443	
p-Fluoro-.alpha.-methylstyrene	385	
4-Fluoro-.alpha.-methylstyrene	385	
1-Fluoro-4-isopropenylbenzene	385	
Fluoroacetic acid	54	
m-Fluoroaniline	274	
p-Fluoroaniline	273	

3-Fluoroaniline	274	
4-Fluoroaniline	273	
m-Fluorobenzaldehyde	327	
p-Fluorobenzaldehyde	328	
3-Fluorobenzaldehyde	327	
4-Fluorobenzaldehyde	328	
Fluorobenzene	266-267	
Fluoroethane	71	
Fluoroethanoic acid	54	
Fluoroethene	53	
.beta.-Fluoroethylamine	80	
2-Fluoroethylamine	80	
Fluoroethylene	53	
Fluoroform	18	
Fluoromethane	27	
2-(4-Fluorophenyl)propene	385	
p-Fluorophenylamine	273	
p-Fluoropyridine	209	
2-Fluoropyridine	208	
3-Fluoropyridine	207	
4-Fluoropyridine	209	
Fluoryl	18	
Formaldehyde (8CI9CI)	21-22	
Formaldehyde solution	21-22	
Formaldehyde, gas	21-22	
Formalin	21-22	
Formalith	21-22	
Formamide, N-methyl- (8CI9CI)	76	
Formamide, N,N-dimethyl- (8CI9CI)	129	
Formic acid (8CI9CI)	23-25	
Formic acid, butyl ester (8CI9CI)	232	
Formic acid, cyano-, ethyl ester (8CI)	154	
Formic acid, ethyl ester (8CI9CI)	122-123	
Formic acid, isopropyl ester (8CI)	179	
Formic acid, methyl ester (8CI9CI)	67-69	
Formic acid, propyl ester (8CI9CI)	173-174	
Formic acid, 1-methylethyl ester (9CI)	179	
Formic aldehyde	21-22	
Formic anammonide	19-20	
Formisoton	23-25	
Formol	21-22	
Formonitrile	19-20	
p-Formylanisole	366	
p-Formylbenzonitrile	361	
4-Formylbenzonitrile	361	
N-Formyldimethylamine	129	
Formylic acid	23-25	
p-Formyltoluene	364	
Fouramine D	292	
Fourrine D	292	
Fourrine 1	292	
Freon F-23	18	
Freon 14	17	
Freon 23	18	
Freon 290	130-131	
Freon 32	20	
Freon 40	26-27	

Freon 41	27		
Freon 600	187		
Fuers Rohr	449		
Fumigrain	99		
Fur Black 41867	292		
Fur Brown 41866	292		
Fur Yellow	292		
Furan (8CI9CI)	150		
Furan, tetrahydro- (8CI9CI)	171		
Furan, tetrahydro-2-methyl- (8CI9CI)	226		
Furanidine	171		
Furfuran	150		
Furro D	292		
Futramine D	292		
Fyde	21-22		
FC 14	17		
T-Gas	65		
Genetron 1132a	48		
Genetron 23	18		
Genetron 32	20		
Gifblaar poison	54		
Glacial acetic acid	66-67		
Glycinenitrile	62		
Glycinenitrile	62		
Glycinonitrile, N-methyl-	111		
Guanin	215		
Guanine (8CI)	215		
Guanine enol	215		
Guanine, 8-methyl- (8CI)	290		
Guanosine, 2',3'-O-(1-methylethylidene)- (9CI)	428		
Guanosine, 2',3'-O-isopropylidene- (8CI)	428		
Halon 1001	25		
Haltox	25		
Hartosol	131-132		
Heavy water	442		
Heavy water-d2	442		
o-Helium	465-466		
p-Helium	465-466		
Helium (8CI9CI)	465-466		
Helium-4	465-466		
Hemostyp	191		
1-Heptanamine (9CI)	355		
n-Heptylamine	355		
Heptylamine (8CI)	355		
Hexafluorobenzene	246		
Hexahydro-s-triazaborine	7		
Hexahydroaniline	310-311		
Hexahydropyrazine	188		
Hexahydropyridine	233-234		
Hexamethylbenzene	423		
Hexamethylenebis(dimethylamine)	418		
Hexamethylenediamine	323		
1-Hexanamine (9CI)	318		
1,6-Hexanediamine (8CI9CI)	323		
1,6-Hexanediamine, N,N,N',N'-tetramethyl- (8CI9CI)	418		
Hexanon	299		
Hexazane	233-234		

n-Hexylamine	318		
1-Hexylamine	318		
Hexylamine (8CI)	318		
1,6-Hexylenediamine	323		
Hyadur	87		
Hydrazine (8CI9CI)	464		
Hydrazine, 1,1-dimethyl- (8CI9CI)		93	
Hydriodic acid (8CI9CI)	447		
Hydriodic ether	73		
Hydrobromic acid (8CI9CI)		11	
Hydrobromic ether	69		
Hydrochloric acid (8CI9CI)		440	
Hydrochloric acid gas	440		
Hydrochloric ether	70		
Hydrochloride	440		
Hydrocyanic acid (8CI9CI)		19-20	
Hydrocyanic ether	104		
Hydrofluoric acid (8CI9CI)		443	
Hydrofluoric acid gas	443		
Hydrogen (8CI9CI)	449-451		
Hydrogen arsenide	2		
Hydrogen bromide	11		
Hydrogen bromide (HBr)	11		
Hydrogen carboxylic acid	23-25		
Hydrogen chloride	440		
Hydrogen chloride (HCl)	440		
Hydrogen cyanide	19-20		
Hydrogen fluoride	443		
Hydrogen fluoride (HF)	443		
Hydrogen iodide	447		
Hydrogen iodide (HI)	447		
Hydrogen phosphide	462-464		
Hydrogen selenide	457		
Hydrogen selenide (H ₂ Se) (8CI9CI)		457	
Hydrogen sulfide	454-456		
Hydrogen sulfide (H ₂ S) (8CI9CI)		454-456	
Hydrogen sulphide	454-456		
Hydrogen-2	441-442		
Hydrosulfuric acid	454-456		
Hydrouracil, 1-.beta.-D-ribofuranosyl- (8CI)		397	
6-Hydroxy-1H-purine	211		
Hydroxybenzene	276		
(Hydroxymethyl)trichloromethane		52	
6-Hydroxypurine	211		
2-Hydroxypyrimidine	148		
Hypnon	363		
Hypnone	363		
Hypoxanthine (8CI)	211		
Hypoxanthine enol	211		
Hypoxanthine, 2-amino-	215		
Hytrol 0	299		
HFA	54		
HMDA	323		
Ice	451-453		
Imidazole, 1-methyl- (8CI)		156	
1H-Imidazole, 1-methyl- (9CI)		156	
Imidole	152		

Imsol A	131-132		
Infiltrina	87		
Iodoethane	73		
Iodomethane	28		
Iron bis(cyclopentadienide)		403	
Iron carbonyl (Fe(CO) ₅) (8CI)		202	
Iron carbonyl (Fe(CO) ₅), (TB-5-11)- (9CI)			202
Iron dicyclopentadienyl	403		
Iron pentacarbonyl	202		
Iron, bis(.eta.5-2,4-cyclopentadien-1-yl)-			403
Isobrome	25		
.beta.-Isoamylene		223	
.gamma.-Isoamylene		224-225	
1-Isoamylene	224-225		
Isobutanal	169		
2-Isobutanethiol	193		
Isobutene	161-162		
Isobutylamine (8CI)		196	
Isobutylene	161-162		
Isobutyraldehyde (8CI)		169	
Isobutyramide (8CI)		186	
Isobutyric acid, methyl ester (8CI)			231
Isobutyrimidic acid	186		
Isobutyronitrile (8CI)		158	
Isocumene	390-391		
Isohol	131-132		
Isonicotinic acid nitrile		261	
Isonicotinonitrile (8CI)		261	
Isopropanethiol	134		
Isopropanol	131-132		
p-Isopropenylanisole	406		
Isopropenylbenzene	387		
Isopropenylcyclopropane		297	
p-Isopropenyltoluene	404		
4-Isopropenyltoluene	404		
2-Isopropoxypropane	312		
Isopropyl alcohol (8CI)		131-132	
Isopropyl aldehyde	169		
Isopropyl cyanide	158		
Isopropyl ether (8CI)		312	
Isopropyl formate	179		
Isopropyl mercaptan	134		
Isopropyl methyl ketone	229		
Isopropyl nitrile	158		
Isopropyl sulfide (8CI)		316	
Isopropylamine (8CI)	137		
Isopropylamine, N-ethyl-		243	
Isopropylbenzene	388-389		
Isopropyldimethylamine		241	
Isopropyldiphenylphosphine oxide		432	
Isopropyl formaldehyde	169		
Isopropylformamide	186		
Isopropylidene guanosine	428		
2',3'-Isopropylidene guanosine		428	
2',3'-O-Isopropylidene guanosine		428	
Isopropylidenemethylene	161-162		
2',3'-Isopropylideneuridine		422	

2',3'-O-Isopropylideneuridine	422	
Isopropylthiol	134	
Isopurine, ribosyl-	405	
Isothiocyanatomethane	58	
Isothiocyanic acid, methyl ester (8CI)	58	
p-Isoxazine, tetrahydro-	184	
Ivalon	21-22	
Izal	276	
Jaysol	82-83	
Jaysol S	82-83	
Kelene	70	
Ketene (8CI)	50	
Ketone, cyclopropyl methyl (8CI)	219	
Ketone, methyl 3-pyridyl (8CI)	331	
Ketone, methyl 4-pyridyl (8CI)	332	
.beta.-Ketopropane	112-114	
Krypton (8CI9CI)	466-467	
Laughing gas	471	
Leuco-4	214	
Levoxine	464	
Liquefied petroleum gas	130-131	
Liquefied petroleum gas	187	
Lithium hydroxide (8CI9CI)	448	
Lithium Hydrate	448	
Lumbrical	188	
Luprosil	119	
.alpha.,.alpha.'-Lutidine	344	
2,4-Lutidine (8CI)	343	
2,5-Lutidine (8CI)	345	
2,6-Lutidine (8CI)	344	
3,5-Lutidine (8CI)	346	
Lutosol	131-132	
Lysoform	21-22	
LPG	130-131	
LPG	187	
Magnesium (8CI9CI)	467	
Magnesium molecule	468	
Malonic acid dinitrile	96	
Malonodinitrile	96	
Malononitrile (8CI)	96	
Manxin	416	
Manxine	416	
Marsh gas	30-31	
Mearlmaid	215	
Mellitene	423	
Mercaptoethane	88	
Mercaptomethane	34-35	
2-Mercaptopropane	134	
Mesitylene (8CI)	392	
Metacetone	225	
Metacetic acid	119	
Metadiazine	146	
Methacetone	225	
Methacide	333-336	
Methan-d3-amine, N,N-di(methyl-d3)- (9CI)	95	
Methanal	21-22	
Methanamine (9CI)	36-37	

Methanamine, N-ethyl-N-methyl-	199	
Methanamine, N-methyl- (9CI)	91-92	
Methanamine, N,N-dimethyl- (9CI)	138-139	
Methanamine, N,N-dimethyl-1-(trimethylsilyl)- (9CI)		324
Methanamine, 1,1,1-trifluoro-N,N-dimethyl-	109	
Methane (8CI9CI)	30-31	
Methane-d3, oxybis- (9CI)	44	
Methane, bromo- (8CI9CI)	25	
Methane, chloro- (8CI9CI)	26-27	
Methane, cyano-	56-57	
Methane, diazo- (8CI9CI)	21	
Methane, dicyano-	96	
Methane, difluoro- (8CI9CI)	20	
Methane, ethoxy-	133	
Methane, fluoro- (8CI9CI)	27	
Methane, iodo- (8CI9CI)	28	
Methane, isothiocyanato- (9CI)	58	
Methane, nitro- (8CI9CI)	29	
Methane, oxybis- (9CI)	84-86	
Methane, sulfinylbis- (9CI)	87	
Methane, tetrafluoro- (9CI)	17	
Methane, thiobis- (9CI)	89	
Methane, trifluoro- (8CI9CI)	18	
Methanecarbonitrile	56-57	
Methanecarboxamide	75	
Methanecarboxylic acid	66-67	
Methanethiol (8CI9CI)	34-35	
Methanoic acid	23-25	
Methanol (8CI9CI)	32-34	
Methanone, diphenyl- (9CI)	426	
p-Methoxy-.alpha.-methylstyrene	406	
4-Methoxy-.alpha.-methylstyrene	406	
2-Methoxy-1-aminobenzene	347	
1-Methoxy-2-aminoethane	141	
2-Methoxy-2-methylpropane	237	
m-Methoxyaniline	349	
o-Methoxyaniline	347	
p-Methoxyaniline	348	
2-Methoxyaniline	347	
3-Methoxyaniline	349	
4-Methoxyaniline	348	
p-Methoxybenzaldehyde	366	
4-Methoxybenzaldehyde	366	
4-Methoxybenzenamine	348	
Methoxybenzene	337	
4-Methoxybenzeneamine	348	
2-Methoxyethanamine	141	
.beta.-Methoxyethylamine	141	
Methoxyethylamine	141	
2-Methoxyethylamine	141	
Methoxymethane	84-86	
2-(p-Methoxyphenyl)propene	406	
2-(p-Methoxyphenyl)propylene	406	
p-Methoxyphenylamine	348	
.beta.-Methoxypyridine	289	
.gamma.-Methoxypyridine	285	
2-Methoxypyridine	287-288	

3-Methoxypyridine	289	
4-Methoxypyridine	285	
Methyl .beta.-pyridyl ketone		331
Methyl acetate	120-121	
Methyl alcohol	32-34	
Methyl aldehyde	21-22	
Methyl benzenecarboxylate		365
Methyl benzoate	365	
Methyl bromide	25	
Methyl carbonate	124	
Methyl carbonate ((MeO)2CO)		124
Methyl chloride	26-27	
Methyl cyanide	56-57	
Methyl cyclopropanecarboxylate		221
Methyl cyclopropyl ketone		219
Methyl cyclopropylcarboxylate		221
Methyl ether (8CI)	84-86	
(Methyl ether)-d6 (8CI)	44	
Methyl ethyl ether	133	
Methyl ethyl ketone	170	
Methyl ethyl sulfide	135	
Methyl fluoride	27	
Methyl formate	67-69	
Methyl hydride	30-31	
Methyl hydroxide	32-34	
Methyl iodide	28	
Methyl isobutyrate	231	
Methyl isopropyl ketone		229
Methyl isothiocyanate		58
Methyl ketone	112-114	
Methyl mercaptan	34-35	
Methyl methanoate	67-69	
Methyl monosulfide	89	
Methyl mustard oil	58	
Methyl nitrite	30	
Methyl phenyl ether	337	
Methyl phenyl ketone	363	
Methyl pivalate	309	
Methyl propanoate	178	
Methyl propionate	178	
Methyl propylate	178	
Methyl sulfide (8CI)	89	
Methyl sulfocyanate	58	
Methyl sulfoxide (8CI)		87
Methyl tert-butyl ether		237
Methyl tert-butyl ketone		308
S-Methyl thioacetate	118	
Methyl thiocyanate	58	
Methyl thioisocyanate		58
Methyl trifluoride	18	
Methyl trifluoroacetate	97	
Methyl trimethylacetate	309	
Methyl 2-methylpropanoate		231
Methyl 2-methylpropionate		231
Methyl 3-pyridyl ketone	331	
Methyl 4-pyridyl ketone	332	
p-Methyl-.alpha.-methylstyrene		404

4-Methyl-.alpha.-methylstyrene	404
N-Methyl-.delta.-valerolactam	301
Methyl-d3 ether	44
N-Methyl-N-ethylamine	140-141
N-Methyl-N-ethylaniline	395
m-Methyl-N,N-dimethylaniline	394
2-Methyl-1-butene	224-225
1-Methyl-1-cyclopentene	295
2-Methyl-1-pentene	306
1-Methyl-1-phenylethylene	387
2-Methyl-1-propene	161-162
1-Methyl-1,2-dihydro-2-pyridinone	286
1-Methyl-2(1H)-pyridinone	286
1-Methyl-2(1H)-pyridone	286
2-Methyl-2-aminopropane	195
3-Methyl-2-butanone	229
2-Methyl-2-butene	223
3-Methyl-2-butene	223
2-Methyl-2-methoxypropane	237
(E)-3-Methyl-2-pentene	304
trans-3-Methyl-2-pentene	304
2-Methyl-2-pentene	305
2-Methyl-2-phenylpropane	410-411
N-Methyl-2-piperidinone	301
1-Methyl-2-piperidinone	301
N-Methyl-2-piperidone	301
1-Methyl-2-piperidone	301
2-Methyl-2-propanethiol	193
2-Methyl-2-propanol	192
3-Methyl-2-propylamine	196
1-Methyl-2-pyridinone	286
N-Methyl-2-pyridone	286
1-Methyl-2-pyridone	286
4-Methyl-3-pentene	305
1-Methyl-4-isopropenylbenzene	404
4-Methyl-4-pentene	306
Methylacetaldehyde	115-116
Methylacetylene	100
Methylamine (8CI)	36-37
(Methylamino)acetonitrile	111
(Methylamino)benzene	339
N-Methylaminoacetonitrile	111
Methylaminoethane	140-141
m-Methylaniline	341-342
p-Methylaniline	340
N-Methylaniline	339
3-Methylaniline	341-342
4-Methylaniline	340
N-Methylaziridine	128
1-Methylaziridine	128
p-Methylbenzaldehyde	364
4-Methylbenzaldehyde	364
m-Methylbenzenamine	341-342
p-Methylbenzenamine	340
N-Methylbenzenamine	339
3-Methylbenzenamine	341-342
4-Methylbenzenamine	340

Methylbenzene	333-336	
Methylbenzol	333-336	
Methylbutanone	229	
Methylcarbinol	82-83	
.alpha.-Methylcyclohexene		352
1-Methylcyclohexene	352	
1-Methylcyclopentene	295	
Methylcyclopropane	164	
5-Methyldeoxyuridine	414	
Methyldiethylamine	240	
N-Methyldiethylamine	240	
Methyldiphenylphosphine oxide		427
Methylene cyanide	96	
Methylene difluoride	20	
Methylene oxide	21-22	
.alpha.-Methylene-diphenylmethane		430
Methylene, dichloro- (8CI9CI)		14-15
Methylene, difluoro- (8CI9CI)		16
Methylenecyclopentane	296	
Methylenedinitrile	96	
1-Methylethanethiol	134	
Methylethylamine	140-141	
N-Methylethylamine	140-141	
1-Methylethylamine	137	
Methylethylene	107-108	
N-Methylethylenimine	128	
1-Methylethylenimine	128	
Methylformamide	76	
N-Methylformamide	76	
8-Methylguanine	290	
N-Methylimidazole	156	
1-Methylimidazole	156	
Methylmethane	79-80	
Methylol	32-34	
Methylolpropane	191	
2-(p-Methylphenyl)propene		404
Methylphenylamine	339	
N-Methylphenylamine	339	
Methylphosphine	38	
N-Methylpiperidine	311-312	
1-Methylpiperidine	311-312	
2-Methylpropanal	169	
2-Methylpropanamide	186	
1-Methylpropanamine	200	
2-Methylpropane nitrile		158
2-Methylpropene	161-162	
.alpha.-Methylpropionaldehyde		169
2-Methylpropionaldehyde		169
2-Methylpropionitrile		158
1-Methylpropylamine	200	
2-Methylpropylamine	196	
6-Methylpurine	275	
.alpha.-Methylpyridine		283
.beta.-Methylpyridine		282
.gamma.-Methylpyridine		281
p-Methylpyridine	281	
2-Methylpyridine	283	

3-Methylpyridine	282	
4-Methylpyridine	281	
N-Methylpyrrolidine	234-235	
1-Methylpyrrolidine	234-235	
.alpha.-Methylstyrene	387	
.alpha.-Methylstyrol	387	
2-Methyltetrahydrofuran	226	
N-Methyltetrahydropyrrole		234-235
(Methylthio)ethane	135	
o-Methyltoluene	367-368	
5-Methyluracil	216	
Miazine	146	
Mirbane oil	268	
Molecular hydrogen		449-451
Molecular oxygen	471-472	
Mono-n-butylamine	197	
Mono-n-hexylamine	318	
Mono-n-propylamine	139-140	
Monoallylamine	126	
Monoamylamine	238	
Monobenzylamine	338	
Monobromoethane	69	
Monobromomethane	25	
Monobutylamine	197	
Monocarbon difluoride		16
Monochlorethane	70	
Monochloroacetonitrile		47
Monochlorobenzene	264-265	
Monochloroethane	70	
Monochloroethanoic acid		51
Monochloromethane	26-27	
Monochloromethyl cyanide		47
Monoethylamine	90	
Monofluoroacetic acid		54
Monofluorobenzene	266-267	
Monofluoroethane	71	
Monofluoroethylene		53
Monohydroxybenzene		276
Monohydroxymethane		32-34
Monoiodoethane	73	
Monoisobutylamine		196
Monoisopropylamine		137
Monomethylamine	36-37	
N-Monomethylaniline		339
Monomethylformamide		76
Monophenol	276	
Monopropylamine	139-140	
Monopyrrole	152	
Morbicid	21-22	
Morpholine (8CI9CI)		184
Morton EP-161E	58	
Muriatic acid	440	
Muriatic ether	70	
Myocol	409	
Myrmicyl	23-25	
MCB	264-265	
MEK	170	

Nadone	299	
Nako H	292	
Naphtol AS-KG	340	
Naphtol AS-KGLL	340	
Narcotile	70	
Natural Pearl Essence	215	
Naturon	215	
Nebularine	405	
Neopentylamine (8CI)	242	
Nicotinic acid nitrile	262	
Nicotinonitrile (8CI)	262	
Niobe oil	365	
Nitric acid, ethyl ester (8CI9CI)	78	
Nitric oxide	469	
Nitric oxide (NO)	469	
Nitro-Sil	458-461	
Nitrobenzene	268	
Nitrobenzol	268	
Nitrocarbol	29	
Nitroethane	77	
Nitrogen (8CI9CI)	470	
Nitrogen fluoride	445	
Nitrogen fluoride (NF3) (8CI9CI)	445	
Nitrogen gas	470	
Nitrogen monoxide	469	
Nitrogen oxide (NO) (8CI9CI)	469	
Nitrogen oxide (N2O) (8CI9CI)	471	
Nitrogen trifluoride	445	
Nitrogen-14	470	
Nitromethane	29	
4-Nitropyridine	210	
Nitrosyl radical	469	
Nitrous acid, methyl ester (8CI9CI)	30	
Nitrous oxide	471	
Norbornene	350	
2-Norbornene (8CI)	350	
Norbornylene	350	
2-Norbornylene	350	
Norcamphene	350	
Norfenchene	350	
Norleucamine	238	
Norvalamine	197	
Nucleocardyl	409	
NSC 65423	405	
Obepin	366	
Oil of Mirbane	268	
Oil of Myrbane	268	
Olefiant gas	59	
Optal	132-133	
Orange GC base	272	
Orsin	292	
Orthamine	291	
Orthodiazine	145	
Orthohydrogen	449-451	
Osmosol extra	132-133	
1-Oxa-4-azacyclohexane	184	
Oxacyclobutane	117	

Oxacyclohexane	228	
Oxacyclopentadiene	150	
Oxacyclopentane	171	
Oxacyclopropane	65	
Oxane	65	
Oxane	228	
2H-1,4-Oxazine, tetrahydro-	184	
4H-1,4-Oxazine, tetrahydro-	184	
Oxetane (9CI)	117	
Oxidoethane	65	
Oxirane (9CI)	65	
Oxirene, dihydro-	65	
.alpha.-Oxodiphenylmethane	426	
.alpha.-Oxoditane	426	
Oxolane	171	
Oxole	150	
Oxomethane	21-22	
6-Oxopurine	211	
2-Oxopyrimidine	148	
Oxybenzene	276	
1,1'-Oxybispropane	313	
Oxyfume	65	
Oxyfume 12	65	
Oxygen (8CI9CI)	471-472	
Oxygen fluoride	444	
Oxygen molecule	471-472	
Oxygen-16	471-472	
Oxymethylene	21-22	
Paradiazine	147	
Paraform	21-22	
Parahydrogen	449-451	
Pearl Essence	215	
Pelagol D	292	
Pelagol DR	292	
Pelagol Grey D	292	
Peltol D	292	
Pentaborane	10	
Pentaborane (B5H9)	10	
Pentaborane(9) (8CI9CI)	10	
Pentacarbonyl iron	202	
2,4-Pentadione	220	
Pentafluorobenzene	247	
1,2,3,4,5-Pentafluorobenzene	247	
Pentamethylene oxide	228	
Pentamethylenediamine	245	
Pentamethylenimine	233-234	
n-Pentanal	227	
Pentanal (9CI)	227	
1-Pentanamine (9CI)	238	
1,5-Pentanediamine (8CI9CI)	245	
2,4-Pentanedione (8CI9CI)	220	
Pentanenitrile (9CI)	222	
3-Pentanone (8CI9CI)	225	
1-Pentene, 2-methyl- (8CI9CI)	306	
2-Pentene, 2-methyl- (8CI9CI)	305	
2-Pentene, 3-methyl-, (E)- (8CI9CI)	304	
n-Pentylamine	238	

1-Pentylamine	238	
tert-Pentylamine (8CI)		239
Pentylamine (8CI)	238	
Perfluoroacetic acid	46-47	
Perfluoroammonia	445	
Perfluorobenzene	246	
Perfluoromethane	17	
Petrohol .	131-132	
Phene	269-270	
Phenethylene	362	
Phenic acid	276	
Phenol (8CI9CI)	276	
Phenyl alcohol	276	
Phenyl chloride	264-265	
Phenyl cyanide	329	
Phenyl fluoride	266-267	
Phenyl hydrate	276	
Phenyl hydride	269-270	
Phenyl hydroxide	276	
Phenyl ketone	426	
Phenyl methyl ether	337	
Phenyl methyl ketone	363	
1-Phenyl-1-methylethylene		387
2-Phenyl-1-propene	387	
Phenylamine	278-280	
N-Phenylaniline	421	
1-Phenylbutane	412-413	
1,2-Phenylenediamine	291	
1,3-Phenylenediamine	293	
1,4-Phenylenediamine	292	
m-Phenylenediamine (8CI)	293	
o-Phenylenediamine (8CI)	291	
p-Phenylenediamine (8CI)	292	
Phenylethane	369-371	
1-Phenylethanone	363	
Phenylethene	362	
Phenylethylene	362	
Phenylic acid	276	
Phenylic alcohol	276	
Phenylmethanal	330	
Phenylmethane	333-336	
(Phenylmethyl)amine	338	
N-Phenylmethylaniline	339	
N-Phenylpiperidine	420	
1-Phenylpiperidine	420	
1-Phenylpropane	390-391	
2-Phenylpropane	388-389	
2-Phenylpropene	387	
.beta.-Phenylpropylene		387
N-Phenylpyrrolidine	407	
1-Phenylpyrrolidine	407	
.alpha.-Phenylstyrene		430
Phenyltrimethylmethane		410-411
Phosphine (PH3)	462-464	
Phosphine (8CI9CI)	462-464	
Phosphine oxide, (1-methylethyl)diphenyl- (9CI)		432
Phosphine oxide, butyldiphenyl-	435	

Phosphine oxide, decanyldiphenyl-	438	
Phosphine oxide, diphenylpropyl-	433	
Phosphine oxide, ethyldiphenyl- (8CI9CI)	431	
Phosphine oxide, isobutyldiphenyl-	434	
Phosphine oxide, isopropyldiphenyl- (8CI)		432
Phosphine oxide, methyldiphenyl- (8CI9CI)		427
Phosphine oxide, neopentyldiphenyl-	436	
Phosphine, dimethyl- (8CI9CI)	92	
Phosphine, methyl- (8CI9CI)	38	
Phosphine, trimethyl- (8CI9CI)	142	
Phosphorus trihydride	462-464	
Piazine	147	
.alpha.-Picoline	283	
.beta.-Picoline	282	
.gamma.-Picoline	281	
m-Picoline	282	
o-Picoline	283	
p-Picoline	281	
2-Picoline (8CI)	283	
3-Picoline (8CI)	282	
4-Picoline (8CI)	281	
Picolinic acid nitrile	263	
Picolinonitrile (8CI)	263	
Pimelic ketone	299	
Pimelin ketone	299	
Pinacolin	308	
Pinacoline	308	
Pinacolone	308	
Piperazidine	188	
1,4-Piperazine	188	
Piperazine (8CI9CI)	188	
Piperidine (8CI9CI)	233-234	
Piperidine, 1-methyl- (8CI9CI)	311-312	
Piperidine, 1-phenyl- (8CI9CI)	420	
Piperidine, 2-methoxy-	302	
2-Piperidinone, 1-methyl- (9CI)	301	
2-Piperidone, 1-methyl- (8CI)	301	
Pipersol	188	
Piroad	149	
Pivalic acid, methyl ester (8CI)	309	
Potassa	448	
Potassium hydrate	448	
Potassium hydroxide (8CI9CI)	448	
Praval	439	
Prolamine	182	
Pronarcol	189-190	
Propadiene	101	
1,2-Propadiene (9CI)	101	
Propaldehyde	115-116	
n-Propan-1-ol	132-133	
n-Propan-2-ol	131-132	
Propanal (9CI)	115-116	
Propanal, 2-methyl- (9CI)	169	
Propanamide, 2-methyl- (9CI)	186	
1-Propanamine (9CI)	139-140	
2-Propanamine (9CI)	137	
2-Propanamine, N-(1-methylethyl)- (9CI)	317	

2-Propanamine, N-(1,1-dimethylethyl)-2-methyl- (9CI)	382
2-Propanamine, N-ethyl- (9CI)	243
2-Propanamine, N-ethyl-N-(1-methylethyl)- (9CI)	381
1-Propanamine, N-propyl- (9CI)	320
1-Propanamine, N,N-diethyl- (9CI)	356
2-Propanamine, N,N-dimethyl- (9CI)	241
1-Propanamine, N,N-dipropyl- (9CI)	399
2-Propanamine, N,N,2-trimethyl- (9CI)	321
1-Propanamine, N,N,2,2-tetramethyl- (9CI)	357
1-Propanamine, 2-methyl- (9CI)	196
2-Propanamine, 2-methyl- (9CI)	195
1-Propanamine, 2-methyl-N-(2-methylpropyl)- (9CI)	378
1-Propanamine, 2,2-dimethyl- (9CI)	242
1-Propanamine, 3,3,3-trifluoro- (9CI)	108
n-Propane	130-131
Propane (8CI9CI)	130-131
.alpha.,.gamma.-Propane oxide	117
Propane, 1,1'-oxybis- (9CI)	313
Propane, 1,1'-thiobis- (9CI)	315
Propane, 1,3-epoxy-	117
Propane, 2-ethoxy- (9CI)	236
Propane, 2-ethoxy-2-methyl- (9CI)	314
Propane, 2-methoxy-2-methyl- (9CI)	237
Propane, 2,2'-oxybis- (9CI)	312
Propane, 2,2'-thiobis- (9CI)	316
Propane, 2,2'-thiobis[2-methyl- (9CI)]	377
1,3-Propanediamine (8CI9CI)	143
1,3-Propanediamine, N,N-dimethyl- (8CI9CI)	244
1,3-Propanediamine, N,N,N',N'-tetramethyl- (8CI9CI)	358
Propanedinitrile (9CI)	96
Propanenitrile (9CI)	104
Propanenitrile, 2-methyl- (9CI)	158
Propanenitrile, 3-amino- (9CI)	111
Propanenitrile, 3-chloro- (9CI)	103
Propanethiol	134
1-Propanethiol (8CI9CI)	134
2-Propanethiol (8CI9CI)	134
2-Propanethiol, 2-methyl- (8CI9CI)	193
Propanoic acid (9CI)	119
Propanoic acid, methyl ester (9CI)	178
Propanoic acid, 2-methyl-, methyl ester (9CI)	231
Propanoic acid, 2,2-dimethyl-, methyl ester (9CI)	309
n-Propanol	132-133
1-Propanol (9CI)	132-133
2-Propanol (9CI)	131-132
2-Propanol, 2-methyl- (9CI)	192
Propanone	112-114
2-Propanone (9CI)	112-114
2-Propanone, acetyl-	220
Propargyl ether	277
Propargylamine	105
2-Propen-1-amine (9CI)	126
2-Propen-1-amine, N-2-propenyl- (9CI)	300
2-Propen-1-amine, N,N-di-2-propenyl- (9CI)	398
Propene (8CI)	107-108
1-Propene (9CI)	107-108
Propene, 2-cyclopropyl- (8CI)	297

Propene, 2-methyl- (8CI)	161-162	
1-Propene, 2-methyl- (9CI)	161-162	
1-Propene, 2-phenyl-	387	
1-Propene, 3,3'-oxybis- (9CI)	300	
Propenenitrile	99	
2-Propenenitrile (9CI)	99	
Propine	100	
Propional	115-116	
Propionaldehyde (8CI)	115-116	
Propione	225	
Propionic acid (8CI)	119	
Propionic acid, methyl ester (8CI)	178	
Propionic aldehyde	115-116	
Propionic nitrile	104	
Propionitrile (8CI)	104	
Propionitrile, 3-amino- (8CI)	111	
Propionitrile, 3-chloro- (8CI)	103	
Propiononitrile	104	
Propol	131-132	
n-Propyl acetate	230	
Propyl acetate	230	
1-Propyl acetate	230	
n-Propyl alcohol	132-133	
sec-Propyl alcohol	131-132	
Propyl alcohol (8CI)	132-133	
Propyl carbinol	191	
Propyl cyanide	159	
Propyl ether (8CI)	313	
n-Propyl formate	173-174	
Propyl formate	173-174	
Propyl hydride	130-131	
n-Propyl mercaptan	134	
Propyl mercaptan	134	
Propyl methanoate	173-174	
Propyl monosulfide	315	
Propyl sulfide (8CI)	315	
Propyl trifluoroacetate	217	
Propylaldehyde	115-116	
n-Propylamine	139-140	
sec-Propylamine	137	
1-Propylamine	139-140	
2-Propylamine	137	
Propylamine (8CI)	139-140	
Propylamine, N-tert-butyl-1,1-dimethyl- (8CI)	400	
Propylamine, N,N-diethyl- (8CI)	356	
Propylamine, N,N,2,2-tetramethyl- (8CI)	357	
Propylamine, 3,3,3-trifluoro- (8CI)	108	
n-Propylbenzene	390-391	
Propylbenzene	390-391	
1-Propylbenzene	390-391	
Propylene	107-108	
1-Propylene	107-108	
1,3-Propylene oxide	117	
1,3-Propylenediamine	143	
1,3-Propylenimine	127	
Propylic alcohol	132-133	
Propylic aldehyde	115-116	

1-Propylmercaptan	134	
2-Propylmercaptan	134	
Propylnitrile	104	
2-Propyn-1-amine (9CI)	105	
2-Propyn-1-amine, N-2-propynyl- (9CI)		284
2-Propyn-1-amine, N,N-di-2-propynyl- (9CI)		386
Propyne (8CI)	100	
1-Propyne (9CI)	100	
1-Propyne, 3,3'-oxybis- (9CI)	277	
2-Propynyl ether (8CI)	277	
2-Propynylamine (8CI)	105	
Prozoin	119	
Prussic acid	19-20	
Pseudoacetic acid	119	
Pseudobutylbenzene	410-411	
Pseudoestragole	406	
Pure oxygen	471-472	
9H-Purin-1-amine, 9-cyclopentyl- (9CI)		408
Purin-6(1H)-one	211	
9H-Purin-6(1H)-one	211	
Purin-6(3H)-one	211	
1H-Purin-6-amine (9CI)	214	
9H-Purin-6-amine, 9-.beta.-D-ribofuranosyl-		409
Purin-6-ol	211	
3H-Purin-6-ol	211	
9H-Purin-6-ol	211	
6H-Purin-6-one, 1,7-dihydro- (9CI)	211	
6H-Purin-6-one, 2-amino-1,7-dihydro- (9CI)		215
6H-Purin-6-one, 2-amino-1,7-dihydro-8-methyl- (9CI)		290
Purine ribonucleoside	405	
Purine riboside	405	
Purine, 6-chloro- (8CI)	203	
1H-Purine, 6-chloro- (9CI)	203	
Purine, 6-methyl- (8CI)	275	
1H-Purine, 6-methyl- (9CI)	275	
9H-Purine, 9-.beta.-D-ribofuranosyl- (8CI9CI)		405
Purinosine	405	
Putrescin	201	
Putrescine	201	
2H-Pyran, tetrahydro- (8CI9CI)	228	
Pyrazine (8CI9CI)	147	
Pyrazine hexahydride	188	
Pyrazine, hexahydro-	188	
Pyridazine (8CI9CI)	145	
4-Pyridinamine, N,N-dimethyl- (9CI)	351	
Pyridine (8CI9CI)	212-213	
Pyridine, hexahydro-	233-234	
Pyridine, 2-(trifluoromethyl)- (8CI9CI)		258
Pyridine, 2-chloro- (8CI9CI)	204	
Pyridine, 2-fluoro- (8CI9CI)	208	
Pyridine, 2-methoxy- (8CI9CI)	287-288	
Pyridine, 2-methyl- (9CI)	283	
Pyridine, 2,4-dimethyl- (9CI)	343	
Pyridine, 2,5-dimethyl- (9CI)	345	
Pyridine, 2,6-bis(1,1-dimethylethyl)- (9CI)		429
Pyridine, 2,6-di-tert-butyl- (8CI)	429	
Pyridine, 2,6-dimethyl- (9CI)	344	

Pyridine, 3-(trifluoromethyl)- (8CI)	259
Pyridine, 3-chloro- (8CI9CI)	205
Pyridine, 3-fluoro- (8CI9CI)	207
Pyridine, 3-methoxy- (8CI9CI)	289
Pyridine, 3-methyl- (9CI)	282
Pyridine, 3,5-dimethyl- (9CI)	346
Pyridine, 4-(dimethylamino)- (8CI)	351
Pyridine, 4-(trifluoromethyl)- (8CI9CI)	260
Pyridine, 4-chloro- (8CI9CI)	206
Pyridine, 4-fluoro- (8CI9CI)	209
Pyridine, 4-methoxy- (8CI9CI)	285
Pyridine, 4-methyl- (9CI)	281
Pyridine, 4-nitro- (8CI9CI)	210
2-Pyridinecarbonitrile (9CI)	263
3-Pyridinecarbonitrile (9CI)	262
4-Pyridinecarbonitrile (9CI)	261
2(1H)-Pyridinone, 1-methyl- (9CI)	286
2(1H)-Pyridone, 1-methyl- (8CI)	286
3-Pyridyl methyl ketone	331
4-Pyridyl methyl ketone	332
2-Pyridyl nitrile	263
3-Pyridylcarbonitrile	262
Pyrimidine (8CI9CI)	146
2,4-Pyrimidinediol	149
2,4-Pyrimidinedione	149
2,4(1H,3H)-Pyrimidinedione (9CI)	149
2,4(1H,3H)-Pyrimidinedione, dihydro-1-.beta.-D-ribofuranosyl-	397
2,4(1H,3H)-Pyrimidinedione, 1-(2-deoxy-.beta.-D-erythro-pentofuranosyl-414	
2,4(1H,3H)-Pyrimidinedione, 5-methyl- (9CI)	216
2-Pyrimidinol	148
2-Pyrimidinone	148
2(1H)-Pyrimidinone (8CI9CI)	148
2(1H)-Pyrimidinone, 4-amino- (9CI)	155
2(1H)-Pyrimidinone, 4-amino-1-.beta.-D-ribofuranosyl-	396
2(1H)-Pyrimidone	148
2-Pyrimidone	148
Pyroacetic ether	112-114
Pyrobenzol	269-270
Pyrobenzole	269-270
Pyrod	149
Pyrrol	152
Pyrrole (8CI)	152
1H-Pyrrole (9CI)	152
Pyrrole, tetrahydro-	182
Pyrrolidine (8CI9CI)	182
Pyrrolidine ring	182
Pyrrolidine, 1-methyl- (8CI9CI)	234-235
Pyrrolidine, 1-phenyl- (8CI9CI)	407
PRO	131-132
Quinuclidine (8CI)	354-355
R 23	18
R 290	130-131
Renal PF	292
.beta.-D-Ribofuranose, 1-(6-amino-9H-purin-9-yl)-1-deoxy-	409
.beta.-D-Ribofuranoside, adenine-9	409
.beta.-D-Ribofuranoside, cytosine-1	396

.beta.-D-Ribofuranoside, thymine-1 2-deoxy-	414	
.beta.-D-Ribofuranoside, 2,4(1H,3H)-pyrimidinedione-1		393
9-(.beta.-D-Ribofuranosyl)purine	405	
9-.beta.-D-Ribofuranosyladenine	409	
1-.beta.-D-Ribofuranosylcytosine	396	
9-.beta.-D-Ribofuranosylpurine	405	
1-.beta.-D-Ribofuranosyluracil	393	
Ribosylpurine	405	
Rohrputz	449	
Rohrreiniger Rofix	449	
Salzsaeure	440	
Sandesin	409	
Santoflex LC	292	
Sarcine	211	
Sarcosinonitrile	111	
Sarkin	211	
Sarkine	211	
Selane	457	
Selenium hydride	457	
Sextone	299	
Silane (8CI9CI)	465	
Silicane	465	
Silicon hydride (SiH ₄)	465	
Silicon tetrahydride	465	
Sodium hydrate	449	
Sodium hydroxide (8CI9CI)	449	
Solvent ether	189-190	
Somipront	87	
Spirit of Hartshorn	458-461	
Stella Polaris	215	
Stink damp	454-456	
Styrene (8CI)	362	
Styrene, .alpha.-methyl- (8CI)	387	
Styrene, p-chloro-.alpha.-methyl- (8CI)	384	
Styrene, p-fluoro-.alpha.-methyl- (8CI)	385	
Styrene, p-methoxy-.alpha.-methyl-	406	
Styrene, p,.alpha.-dimethyl- (8CI)	404	
Styrol	362	
Styrole	362	
Styrolene	362	
Styropol S0	362	
Sulfide, ethyl methyl (8CI)	135	
Sulfinylbismethane	87	
Sulfur fluoride (SF ₆) (8CI)	446	
Sulfur fluoride (SF ₆), (OC-6-11)- (9CI)	446	
Sulfur hexafluoride	446	
Sulfur hydride	454-456	
Sulfur oxide (S03)	473	
Sulfur trioxide (8CI9CI)	473	
Sulfureted hydrogen	454-456	
Sulfuric anhydride	473	
Sulfuric ether	189-190	
Superlysoform	21-22	
SD Alcohol 23-hydrogen	82-83	
S03	473	
SQ 9453	87	
Takineocol	131-132	

Tecsol	82-83	
Terabol	25	
Terephthalaldehydonitrile (8CI)		361
Tereton	120-121	
Tertral D	292	
Tetraborane(10) (8CI9CI)	9	
Tetraborane(8)	8	
1,2,3,4-Tetrafluorobenzene		249
1,2,3,5-Tetrafluorobenzene		250
1,2,4,5-Tetrafluorobenzene		248
1,2,4,6-Tetrafluorobenzene		250
1,3,4,5-Tetrafluorobenzene		250
2,3,4,5-Tetrafluorobenzene		249
2,3,5,6-Tetrafluorobenzene		248
Tetrafluoromethane	17	
Tetrahydro-p-oxazine	184	
Tetrahydro-1,4-oxazine		184
Tetrahydro-2-methylfuran		226
Tetrahydro-2H-pyran	228	
Tetrahydro-2H-1,4-oxazine		184
Tetrahydrofuran	171	
Tetrahydropyran	228	
Tetrahydropyrrole		182
Tetrahydrosylvan	226	
N,N,N',N'-Tetramethyl-1,2-diaminoethane		322
N,N,N',N'-Tetramethyl-1,2-ethanediamine		322
N,N,N',N'-Tetramethyl-1,3-diaminopropane		358
N,N,N',N'-Tetramethyl-1,3-propanediamine		358
N,N,N',N'-Tetramethyl-1,4-butanediamine		383
N,N,N',N'-Tetramethyl-1,4-diaminobutane		383
N,N,N',N'-Tetramethyl-1,6-diaminohexane		418
N,N,N',N'-Tetramethyl-1,6-hexanediamine		418
2,2,4,4-Tetramethyl-3-thiapentane		377
N,N,N',N'-Tetramethylbutanediamine		383
.alpha.,.delta.-(Tetramethyldiamino)butane		383
Tetramethyldiaminobutane	383	
Tetramethyldiaminoethane	322	
Tetramethylene oxide	171	
Tetramethylenediamine		201
Tetramethylenimine	182	
N,N,N',N'-Tetramethylethanediamine		322
N,N,N',N'-Tetramethylethylenediamine		322
N,N,N',N'-Tetramethylhexamethylenediamine		418
Tetramethylpropanediamine	358	
Tetramethylputrescine	383	
N,N,N',N'-Tetramethyltetramethylenediamine		383
N,N,N',N'-Tetramethyltrimethylenediamine		358
Tetrole	150	
2-Thiabutane	135	
4-Thiaheptane	315	
3-Thiapentane	194	
2-Thiapropane	89	
Thiocyanic acid, methyl ester (8CI9CI)		58
Thioethanol	88	
Thioethyl alcohol	88	
Thioethyl ether	194	
Thymidin	414	

Thymidine (8CI9CI)	414	
Thymin (purine base)	216	
Thymine (8CI)	216	
Thymine-2-desoxyribose	414	
Thymine-2-desoxyribose	414	
4-Tolualdehyde	364	
p-Tolualdehyde (8CI)	364	
p-Tolualdehyde, .alpha.,.alpha.,.alpha.-trifluoro- (8CI)	360	
Toluene (8CI)	333-336	
3-Toluidine	341-342	
4-Toluidine	340	
m-Toluidine (8CI)	341-342	
p-Toluidine (8CI)	340	
m-Toluidine, N,N-dimethyl- (8CI)	394	
Toluol	333-336	
p-Toluyaldehyde	364	
p-Tolylaldehyde	364	
p-Tolylamine	340	
2-p-Tolylpropene	404	
Trapex	58	
Trapexide	58	
Tri(methyl-d3)amine	95	
Tri-n-butylamine	425	
Tri-n-propylamine	399	
Tri-2-propynylamine (8CI)	386	
Triallylamine (8CI)	398	
s-Triazaborane	7	
1,3,5,2,4,6-Triazatriborine, hexahydro-	7	
Triborinetriamine	7	
Triboron nitride	7	
Tributylamine (8CI)	425	
Trichlorethanol	52	
Trichloroacetonitrile	43	
.beta.-Trichloroethanol	52	
Trichloroethanol	52	
2,2,2-Trichloroethanol	52	
2,2,2-Trichloroethyl alcohol	52	
Trichloromethyl cyanide	43	
Triethylamine (8CI)	319	
Triethylamine, 1,1'-dimethyl- (8CI)	381	
Triethylenediamine	307-308	
.alpha.,.alpha.,.alpha.-Trifluoro-p-tolualdehyde	360	
Trifluoroacetic acid	46-47	
Trifluoroamine	445	
Trifluoroammonia	445	
sym-Trifluorobenzene	252-253	
1,2,4-Trifluorobenzene	251	
1,2,5-Trifluorobenzene	251	
1,3,4-Trifluorobenzene	251	
1,3,5-Trifluorobenzene	252-253	
Trifluoroborane	4	
Trifluoroboron	4	
4,4,4-Trifluorobutylamine	167	
Trifluoroethanoic acid	46-47	
2,2,2-Trifluoroethanol	55	
Trifluoroethene	45	
.beta.,.beta.,.beta.-Trifluoroethyl alcohol	55	

2,2,2-Trifluoroethyl alcohol	55	
2,2,2-Trifluoroethyl formate	98	
(2,2,2-Trifluoroethyl)dimethylamine	61-62	166
Trifluoroethylamine	61-62	
2,2,2-Trifluoroethylamine	45	61-62
Trifluoroethylene	18	
Trifluoromethane	360	
p-(Trifluoromethyl)benzaldehyde	360	
4-(Trifluoromethyl)benzaldehyde	258	
2-(Trifluoromethyl)pyridine	260	
4-(Trifluoromethyl)pyridine	108	
3,3,3-Trifluoropropylamine	324	
Trimethyl(dimethylaminomethyl)silane	138-139	
Trimethylamine (8CI)	95	
Trimethylamine-d ₉ (8CI)	324	
Trimethylamine, (trimethylsilyl)- (8CI)	195	
Trimethylaminomethane	394	
m,N,N-Trimethylaniline	394	
N,N,3-Trimethylaniline	136	
Trimethylarsenic	136	
Trimethylarsine	392	
sym-Trimethylbenzene	392	
1,3,5-Trimethylbenzene	192	
Trimethylcarbinol	106	
Trimethylene	106	
Trimethylene (cyclic)	117	
Trimethylene oxide (8CI)	143	
Trimethylenediamine	127	
Trimethylenimine	223	
Trimethylethylene	223	
1,1,2-Trimethylethylene	192	
Trimethylmethanol	410-411	
Trimethylphenylmethane	142	
Trimethylphosphine	142	
Trimethylphosphorus	437	
Triphenylamine (8CI)	386	
Tripropargylamine	399	
Tripropylamine (8CI)	425	
Tris-n-butylamine	43	
Tritox	58	
Tropex	200	
Tutane	171	
THF	228	
THP	143	
TMEDA	93	
Unsymmetrical dimethylhydrazine	149	
Ura	149	
Uracil (8CI)	393	
Uracil, 1-.beta.-D-ribofuranosyl-	474	
Uranium (8CI9CI)	474	
Uranium I (238U)	474	
Uranium-238	393	
Urd	393	
Uridin	393	
.beta.-Uridine	393	
Uridine (8CI9CI)	422	
Uridine 2',3'-acetonide		

Uridine, 2'-deoxy-5-methyl-	414	
Uridine, 2',3'-O-(1-methylethylidene)- (9CI)		422
Uridine, 2',3'-O-isopropylidene- (8CI)		422
Uridine, 5,6-dihydro- (9CI)	397	
Ursol D	292	
Uvilon	188	
UDMH	93	
Valamine	196	
Valeral	227	
n-Valeraldehyde	227	
Valeraldehyde (8CI)	227	
Valerianic aldehyde	227	
Valeric acid aldehyde		227
Valeric aldehyde	227	
Valeronitrile (8CI)	222	
Valerylaldehyde	227	
Ventox	99	
Versneller NL 63/10		375-376
Vinegar acid	66-67	
Vinegar naphtha		176-177
Vinyl cyanide	99	
Vinyl fluoride	53	
Vinylbenzene	362	
Vinylbenzol	362	
Vinylcyclopropane		218
Vinylcyclopropane	218	
Vinylidene difluoride		48
Vinylidene fluoride		48
Vitamin B4	214	
VCN	99	
Water (8CI9CI)		451-453
Water vapor	451-453	
Water-d2 (9CI)		442
Water-2H2	442	
Water, heavy (D2O) (8CI)		442
White caustic	449	
Wood alcohol	32-34	
Wood ether	84-86	
Worm-A-Ton	188	
Wurmrazin	188	
WN 12	58	
Xenon (8CI9CI)		475
1,2-Xylene	367-368	
1,3-Xylene	373	
1,4-Xylene	372	
m-Xylene (8CI)		373
o-Xylene (8CI)		367-368
p-Xylene (8CI)		372
p-Xylene-d6	359	
m-Xylol	373	
o-Xylol	367-368	
p-Xylol	372	
Zoba Black D	292	

U.S. DEPT. OF COMM. BIBLIOGRAPHIC DATA SHEET	1. PUBLICATION OR REPORT NO. NBSIR 79-1777	2. Gov't. Accession No.	3. Recipient's Accession No.
4. TITLE AND SUBTITLE A Compendium of Gas Phase Basicity and Proton Affinity Measurements		5. Publication Date July 1979	
		6. Performing Organization Code	
7. AUTHOR(S) K. N. Hartman, S. Lias, P. Ausloos, H. M. Rosenstock, S. S. Schroyer, C. Schmidt, D. Martinsen, & G. W. A. Milne		8. Performing Organ. Report No.	
9. PERFORMING ORGANIZATION NAME AND ADDRESS NATIONAL BUREAU OF STANDARDS DEPARTMENT OF COMMERCE WASHINGTON, DC 20234		10. Project/Task/Work Unit No.	
		11. Contract/Grant No.	
12. SPONSORING ORGANIZATION NAME AND COMPLETE ADDRESS (Street, City, State, ZIP) same as above.		13. Type of Report & Period Covered April 1979	
		14. Sponsoring Agency Code	
15. SUPPLEMENTARY NOTES ☐ Document describes a computer program; SF-185, FIPS Software Summary, is attached.			
16. ABSTRACT (A 200-word or less factual summary of most significant information. If document includes a significant bibliography or literature survey, mention it here.) Literature values of experimental measurements, calculations and estimations of gas phase basicities and proton affinities are presented. Where appropriate they are recalculated to common values for the proton affinity of ammonia and the heat of formation of the proton. The annotated information is ordered by molecular formula following the Hill System. Indices are given to permit location of information of by numerical value of the measurement(s), the compound formula, name and synonym, and CAS Registry Number. The literature coverage is essentially complete through 1977.			
17. KEY WORDS (six to twelve entries; alphabetical order; capitalize only the first letter of the first key word unless a proper name; separated by semicolons)			
18. AVAILABILITY ☒ Unlimited ☐ For Official Distribution. Do Not Release to NTIS ☐ Order From Sup. of Doc., U.S. Government Printing Office, Washington, DC 20402, SD Stock No. SN003-003- ☒ Order From National Technical Information Service (NTIS), Springfield, VA, 22161		19. SECURITY CLASS (THIS REPORT) UNCLASSIFIED	21. NO. OF PRINTED PAGES 562
		20. SECURITY CLASS (THIS PAGE) UNCLASSIFIED	22. Price \$16.25

